

Lecture 1: Variational formulation and Sobolev spaces

Pr. Ismail Merabet

Univ. of K-M-Ouargla

October 19, 2024

Contents

- 1 Variational formulation
- 2 Green's Formula
- 3 Sobolev spaces $H^1(\Omega)$, $H^m(\Omega)$
- 4 A Sobolev embedding theorem
- 5 Trace theorem
- 6 The space $H_0^1(\Omega)$

Model problem in 1d

Consider the following boundary value problem

$$\begin{cases} -u'' = f & \text{in }]0, 1[\\ u(0) = u(1) = 0. \end{cases} \quad (1)$$

If u is the solution of (1) and v is a sufficiently regular function satisfies $v(0) = v(1) = 0$, then multiplying the first equation of (1) by v integrating we get:

Model problem in 1d

Consider the following boundary value problem

$$\begin{cases} -u'' = f & \text{in }]0, 1[\\ u(0) = u(1) = 0. \end{cases} \quad (1)$$

If u is the solution of (1) and v is a sufficiently regular function satisfies $v(0) = v(1) = 0$, then multiplying the first equation of (1) by v integrating we get:

$$\int_0^1 -u''(x)v(x) \, dx = \int_0^1 f(x)v(x) \, dx =: (f, v) \quad (2)$$

Model problem in 1d

Consider the following boundary value problem

$$\begin{cases} -u'' = f & \text{in }]0, 1[\\ u(0) = u(1) = 0. \end{cases} \quad (1)$$

If u is the solution of (1) and v is a sufficiently regular function satisfies $v(0) = v(1) = 0$, then multiplying the first equation of (1) by v integrating we get:

$$\int_0^1 -u''(x)v(x) \, dx = \int_0^1 f(x)v(x) \, dx =: (f, v) \quad (2)$$

then integration by parts yields

$$(f, v) = \int_0^1 u'(x)v'(x) \, dx =: a(u, v). \quad (3)$$

Let us define formally¹

$$V = \{v \in L^2(]0, 1[); a(v, v) < +\infty \text{ and } v(0) = v(1) = 0\}.$$

¹for the moment since the notion of derivative to be used has not been made precise

Let us define formally¹

$$V = \{v \in L^2(]0, 1[); a(v, v) < +\infty \text{ and } v(0) = v(1) = 0\}.$$

Defintion

The space V is called the functional space

Then the solution of (1) is characterized by

¹for the moment since the notion of derivative to be used has not been made precise

Let us define formally¹

$$V = \{v \in L^2(]0, 1[); a(v, v) < +\infty \text{ and } v(0) = v(1) = 0\}.$$

Defintion

The space V is called the functional space

Then the solution of (1) is characterized by

$$\begin{cases} \text{Find } u \in V \text{ s.t} \\ a(u, v) = (f, v), \quad \forall v \in V. \end{cases} \quad (4)$$

¹for the moment since the notion of derivative to be used has not been made precise

Let us define formally¹

$$V = \{v \in L^2(]0, 1[); a(v, v) < +\infty \text{ and } v(0) = v(1) = 0\}.$$

Definition

The space V is called the functional space

Then the solution of (1) is characterized by

$$\begin{cases} \text{Find } u \in V \text{ s.t} \\ a(u, v) = (f, v), \quad \forall v \in V. \end{cases} \quad (4)$$

Definition

Problem (4) is called *the weak formulation* or the variational problem of problem (1).

¹for the moment since the notion of derivative to be used has not been made precise

Remark

By construction it is clear that any solution of (1) is a solution of (4). Of course, the central issue is that (4) gives a solution of (1).

Remark

By construction it is clear that any solution of (1) is a solution of (4). Of course, the central issue is that (4) gives a solution of (1). The following theorem verifies this under some simplifying assumptions.

Remark

By construction it is clear that any solution of (1) is a solution of (4). Of course, the central issue is that (4) gives a solution of (1). The following theorem verifies this under some simplifying assumptions.

Suppose that $f \in C^0([0, 1])$ and $u \in C^2([0, 1])$ satisfies (4). Then u is a solution of (1).

Proof

Let $v \in V \cap C^1([0, 1])$, then

$$(f, v) = \int_{\Omega} u' v' \, dx = - \int_{\Omega} u'' v \, dx + u' v|_0^1 = (-u'', v).$$

Proof

Let $v \in V \cap C^1([0, 1])$, then

$$(f, v) = \int_{\Omega} u' v' \, dx = - \int_{\Omega} u'' v \, dx + u' v|_0^1 = (-u'', v).$$

So,

$$(f + u'', v) = 0, \quad \forall v \in V \cap C^1([0, 1]). \quad (5)$$

To finish the proof, it suffices to prove that (5) implies $-u'' = f$.

Proof

Let $v \in V \cap C^1([0, 1])$, then

$$(f, v) = \int_{\Omega} u' v' \, dx = - \int_{\Omega} u'' v \, dx + u' v|_0^1 = (-u'', v).$$

So,

$$(f + u'', v) = 0, \quad \forall v \in V \cap C^1([0, 1]). \quad (5)$$

To finish the proof, it suffices to prove that (5) implies $-u'' = f$. Indeed, if not ($-u'' \neq f$) then (5) implies the existence of x_0 and x_1 , $0 < x_0 < x_1 < 1$ such that $(f + u'')$ has the same sign on $[x_0, x_1]$ i.e. $f + u'' > 0$ or $f + u'' < 0$.

Let $v \in V \cap C^1([0, 1])$, then

$$(f, v) = \int_{\Omega} u' v' \, dx = - \int_{\Omega} u'' v \, dx + u' v|_0^1 = (-u'', v).$$

So,

$$(f + u'', v) = 0, \quad \forall v \in V \cap C^1([0, 1]). \quad (5)$$

To finish the proof, it suffices to prove that (5) implies $-u'' = f$. Indeed, if not ($-u'' \neq f$) then (5) implies the existence of x_0 and x_1 ,

$0 < x_0 < x_1 < 1$ such that $(f + u'')$ has the same sign on $[x_0, x_1]$ i.e. $f + u'' > 0$ or $f + u'' < 0$.

take

$$v(x) = (x - x_0)^2(x - x_1)^2$$

this implies that $(f + u'', v) \neq 0$ which is in contradiction with (5).

Remark

- The choice of the functional space is crucial. It seems to be natural that the space where we seek the solution is the same as the space of the test functions v .

Remark

- The choice of the functional space is crucial. It seems to be natural that the space where we seek the solution is the same as the space of the test functions v .
- Observe that if the space is with finite dimension then the weak formulation (4) can be reformulated as a linear system $Ax = b$. in that case, the uniqueness means that A is injective thus it is an isomorphism this implies the existence also.

Remark

- The choice of the functional space is crucial. It seems to be natural that the space where we seek the solution is the same as the space of the test functions v .
- Observe that if the space is with finite dimension then the weak formulation (4) can be reformulated as a linear system $Ax = b$. in that case, the uniqueness means that A is injective thus it is an isomorphism this implies the existence also.
- But we work here in infinite dimensional case, and the previous theory does not work here. The existence of solution follows from the Lax-Milgram theorem which suppose that the space V must be a Hilbert space.

Let Ω be a bounded open subset of \mathbb{R}^N with boundary $\partial\Omega$ of class C^1 .

Let Ω be a bounded open subset of \mathbb{R}^N with boundary $\partial\Omega$ of class C^1 .

(Green's Formula I)

Let $v \in C^1(\bar{\Omega})$ with compact support in $\bar{\Omega}$. Then we have

$$\int_{\Omega} \frac{\partial v}{\partial x_i}(x) \, dx = \int_{\partial\Omega} v(x) n_i(x) \, ds \quad (6)$$

where n_i is the i -th component of the unit outward normal to Ω .

Let Ω be a bounded open subset of \mathbb{R}^N with boundary $\partial\Omega$ of class C^1 .

(Green's Formula I)

Let $v \in C^1(\bar{\Omega})$ with compact support in $\bar{\Omega}$. Then we have

$$\int_{\Omega} \frac{\partial v}{\partial x_i}(x) \, dx = \int_{\partial\Omega} v(x) n_i(x) \, ds \quad (6)$$

where n_i is the i -th component of the unit outward normal to Ω .

We can generalize Green's formula to elements of the space $H^1(\Omega)$.

Let Ω be a bounded open subset of \mathbb{R}^N with boundary $\partial\Omega$ of class C^1 .

(Green's Formula I)

Let $v \in C^1(\bar{\Omega})$ with compact support in $\bar{\Omega}$. Then we have

$$\int_{\Omega} \frac{\partial v}{\partial x_i}(x) \, dx = \int_{\partial\Omega} v(x) n_i(x) \, ds \quad (6)$$

where n_i is the i -th component of the unit outward normal to Ω .

We can generalize Green's formula to elements of the space $H^1(\Omega)$.

Theorem

Let Ω be a bounded open subset of \mathbb{R}^N with boundary $\partial\Omega$ of class C^1 . If u, v are two functions in $H^1(\omega)$.

Let Ω be a bounded open subset of \mathbb{R}^N with boundary $\partial\Omega$ of class C^1 .

(Green's Formula I)

Let $v \in C^1(\bar{\Omega})$ with compact support in $\bar{\Omega}$. Then we have

$$\int_{\Omega} \frac{\partial v}{\partial x_i}(x) \, dx = \int_{\partial\Omega} v(x) n_i(x) \, ds \quad (6)$$

where n_i is the i -th component of the unit outward normal to Ω .

We can generalize Green's formula to elements of the space $H^1(\Omega)$.

Theorem

Let Ω be a bounded open subset of \mathbb{R}^N with boundary $\partial\Omega$ of class C^1 . If u, v are two functions in $H^1(\omega)$. Then

$$\int_{\Omega} u \frac{\partial v}{\partial x_i}(x) \, dx = - \int_{\Omega} \frac{\partial u}{\partial x_i}(x) v(x) \, dx + \int_{\partial\Omega} u(x) v(x) n_i(x) \, ds \quad (7)$$

(Green's formula II)

Let Ω be a bounded open subset of \mathbb{R}^N with boundary $\partial\Omega$ of class C^1 .

(Green's formula II)

Let Ω be a bounded open subset of \mathbb{R}^N with boundary $\partial\Omega$ of class C^1 .
If $u \in H^2(\Omega)$ and $v \in H^1(\Omega)$.

(Green's formula II)

Let Ω be a bounded open subset of \mathbb{R}^N with boundary $\partial\Omega$ of class C^1 . If $u \in H^2(\Omega)$ and $v \in H^1(\Omega)$. Then

$$-\int_{\Omega} \Delta u \, v \, dx = \int_{\Omega} \nabla u \cdot \nabla v \, dx - \int_{\partial\Omega} \frac{\partial u}{\partial n} \, v \, ds \quad (8)$$

Proof

Sobolev Spaces

Let Ω be a bounded open subset of \mathbb{R}^N .

Sobolev Spaces

Let Ω be a bounded open subset of \mathbb{R}^N .

Definition

$C_0^\infty(\Omega)$ is the space of class C^∞ with compact support included in Ω and $\mathcal{D}'(\Omega)$ is the space of distributions defined on Ω .

Sobolev Spaces

Let Ω be a bounded open subset of \mathbb{R}^N .

Definition

$C_0^\infty(\Omega)$ is the space of class C^∞ with compact support included in Ω and $\mathcal{D}'(\Omega)$ is the space of distributions defined on Ω .

Definition

Given a function $v \in L^2(\Omega)$.

Sobolev Spaces

Let Ω be a bounded open subset of \mathbb{R}^N .

Definition

$C_0^\infty(\Omega)$ is the space of class C^∞ with compact support included in Ω and $\mathcal{D}'(\Omega)$ is the space of distributions defined on Ω .

Definition

Given a function $v \in L^2(\Omega)$. We say that v has a weak derivative if there exists $w_i \in L^2(\Omega)$, for all $i = 1, \dots, N$ such that for all $\phi \in C_0^\infty(\Omega)$ we have

$$\int_{\Omega} v \frac{\partial \phi}{\partial x_i} = - \int_{\Omega} w_i \phi \, dx$$

the functions w_i are called the weak derivatives and they are denoted by $\frac{\partial v}{\partial x_i}$.

The Sobolev space $H^1(\Omega)$

Definition

Let Ω be an open set of \mathbb{R}^N .

The Sobolev space $H^1(\Omega)$

Definition

Let Ω be an open set of \mathbb{R}^N . The Sobolev space $H^1(\Omega)$ is given by :

$$H^1(\Omega) =: \{u \in L^2(\Omega); \frac{\partial u}{\partial x_i} \in L^2(\Omega), i = 1, 2, \dots, N\} \quad (9)$$

where $\frac{\partial u}{\partial x_i}$ is the weak derivative of u

More, generally, Let

$$\alpha = (\alpha_1, \alpha_2, \dots, \alpha_i, \dots, \alpha_N), \alpha_i \in \mathbb{N}, \quad i = 1, 2, \dots, N$$

a multi index

More, generally, Let

$$\alpha = (\alpha_1, \alpha_2, \dots, \alpha_i, \dots, \alpha_N), \alpha_i \in \mathbb{N}, \quad i = 1, 2, \dots, N$$

a multi index We note

$$D^\alpha = \partial_1^{\alpha_1} \partial_2^{\alpha_2} \dots \partial_N^{\alpha_N}, \quad |\alpha| = \sum_{i=1}^N \alpha_i.$$

More, generally, Let

$$\alpha = (\alpha_1, \alpha_2, \dots, \alpha_i, \dots, \alpha_N), \alpha_i \in \mathbb{N}, \quad i = 1, 2, \dots, N$$

a multi index We note

$$D^\alpha = \partial_1^{\alpha_1} \partial_2^{\alpha_2} \dots \partial_N^{\alpha_N}, \quad |\alpha| = \sum_{i=1}^N \alpha_i.$$

Definition

for $m \in \mathbb{N}$

$$H^m(\Omega) =: \{u \in \mathcal{D}'(\Omega); D^\alpha u \in L^2(\Omega) \quad |\alpha| \leq m\} \quad (10)$$

For $m = 0$ we have $H^0(\Omega) = L^2(\Omega)$ and for $m = 1$, $H^1(\Omega)$.

Sobolev space $H^m(\Omega)$

We equip $H^m(\Omega)$ by the inner product:

$$(u, v)_m = \sum_{|\alpha| \leq m} \int_{\Omega} D^{\alpha} u D^{\alpha} v \, dx. \quad (11)$$

Sobolev space $H^m(\Omega)$

We equip $H^m(\Omega)$ by the inner product:

$$(u, v)_m = \sum_{|\alpha| \leq m} \int_{\Omega} D^{\alpha} u D^{\alpha} v \, dx. \quad (11)$$

The associated norm is

$$\|u\|_{m,\Omega} = \left(\sum_{|\alpha| \leq m} \int_{\Omega} |D^{\alpha} u|^2 \right)^{1/2}. \quad (12)$$

Sobolev space $H^m(\Omega)$

We equipped $H^m(\Omega)$ by the inner product:

$$(u, v)_m = \sum_{|\alpha| \leq m} \int_{\Omega} D^{\alpha} u D^{\alpha} v \, dx. \quad (11)$$

The associated norm is

$$\|u\|_{m,\Omega} = \left(\sum_{|\alpha| \leq m} \int_{\Omega} |D^{\alpha} u|^2 \right)^{1/2}. \quad (12)$$

Proposition

- ① If $m \geq m'$, $H^m(\Omega)$ is continuously embedded in $H^{m'}(\Omega)$.

Sobolev space $H^m(\Omega)$

We equip $H^m(\Omega)$ by the inner product:

$$(u, v)_m = \sum_{|\alpha| \leq m} \int_{\Omega} D^{\alpha} u D^{\alpha} v \, dx. \quad (11)$$

The associated norm is

$$\|u\|_{m,\Omega} = \left(\sum_{|\alpha| \leq m} \int_{\Omega} |D^{\alpha} u|^2 \right)^{1/2}. \quad (12)$$

Proposition

- 1 If $m \geq m'$, $H^m(\Omega)$ is continuously embedded in $H^{m'}(\Omega)$.
- 2 $H^m(\Omega)$ equipped with the inner product (11) is a Hilbert space.

Theorem

Let Ω be a bounded open subset of \mathbb{R}^2 with Lipschitz boundary m and k two integer numbers satisfying $k - m > 1$.

Theorem

Let Ω be a bounded open subset of \mathbb{R}^2 with Lipschitz boundary m and k two integer numbers satisfying $k - m > 1$. Then there exists a positive constant C such that for all $u \in H^k(\Omega)$ we have:

Theorem

Let Ω be a bounded open subset of \mathbb{R}^2 with Lipschitz boundary m and k two integer numbers satisfying $k - m > 1$. Then there exists a positive constant C such that for all $u \in H^k(\Omega)$ we have:

$$\|D^m u\|_{L^\infty(\Omega)} \leq C \|u\|_{k,\Omega} \quad (13)$$

In addition there exists a function of class C^m equal to u almost everywhere.

Trace theorem

Suppose that Ω is sufficiently regular (of class C^1 , for example).

Trace theorem

Suppose that Ω is sufficiently regular (of class C^1 , for example). Then we define the trace γ_0 by :

$$\begin{aligned}\gamma_0 : H^1(\Omega) \cap C^0(\bar{\Omega}) &\rightarrow L^2(\partial\Omega) \cap C^0(\overline{\partial\Omega}) \\ u &\mapsto \gamma_0(u) = u|_{\partial\Omega}\end{aligned}\tag{14}$$

Trace theorem

Suppose that Ω is sufficiently regular (of class C^1 , for example). Then we define the trace γ_0 by :

$$\begin{aligned}\gamma_0 : H^1(\Omega) \cap C^0(\bar{\Omega}) &\rightarrow L^2(\partial\Omega) \cap C^0(\overline{\partial\Omega}) \\ u &\mapsto \gamma_0(u) = u|_{\partial\Omega}\end{aligned}\tag{14}$$

Theorem

The linear map γ_0 given in (14) can be extended to a linear continuous map from $H^1(\Omega)$ to $L^2(\partial\Omega)$, i.e., there exists c such that:

Trace theorem

Suppose that Ω is sufficiently regular (of class C^1 , for example). Then we define the trace γ_0 by :

$$\begin{aligned}\gamma_0 : H^1(\Omega) \cap C^0(\bar{\Omega}) &\rightarrow L^2(\partial\Omega) \cap C^0(\overline{\partial\Omega}) \\ u &\mapsto \gamma_0(u) = u|_{\partial\Omega}\end{aligned}\tag{14}$$

Theorem

The linear map γ_0 given in (14) can be extended to a linear continuous map from $H^1(\Omega)$ to $L^2(\partial\Omega)$, i.e., there exists c such that:

$$\|u\|_{L^2(\partial\Omega)} \leq c \|u\|_{H^1(\Omega)}, \forall u \in H^1(\Omega).\tag{15}$$

The space $H_0^1(\Omega)$

Definition

Let $\mathcal{D}(\Omega)$ the space of C^∞ with compact support included in Ω .

The space $H_0^1(\Omega)$

Definition

Let $\mathcal{D}(\Omega)$ the space of C^∞ with compact support included in Ω . We define $H_0^1(\Omega)$ as the adherence of $\mathcal{D}(\Omega)$ in $H^1(\Omega)$, i.e.,

$$\overline{\mathcal{D}(\Omega)}^{\|\cdot\|_{1,\Omega}} =: H_0^1(\Omega).$$

The space $H_0^1(\Omega)$

Definition

Let $\mathcal{D}(\Omega)$ the space of C^∞ with compact support included in Ω . We define $H_0^1(\Omega)$ as the adherence of $\mathcal{D}(\Omega)$ in $H^1(\Omega)$, i.e.,

$$\overline{\mathcal{D}(\Omega)}^{\|\cdot\|_{1,\Omega}} =: H_0^1(\Omega).$$

Theorem

The $H_0^1(\Omega)$ is the kernel of γ_0 , i.e.,

$$H_0^1(\Omega) = \{u \in H^1(\Omega), u|_{\partial\Omega} = 0\}$$

Lemma

Let Ω a bounded set of \mathbb{R}^N . Then, there exists a positive C which depends only on Ω such that:

$$\forall v \in H_0^1(\Omega), \quad \|v\|_{L^2(\Omega)} \leq C \|\nabla v\|_{L^2(\Omega)}. \quad (16)$$