

Lecture 1: A posteriori error analysis by duality

Pr. Ismail Merabet

Univ. of K-M-Ouargla

October 3, 2025

Contents

- 1 Introduction
- 2 One-dimensional model problem
- 3 Numerical Test
- 4 Conclusion

Introduction

- In the previous year, MEF II, for some elliptic problems, we have seen how to show that

$$\|u - u_h\|_{H^1} \leq Ch^k |u|_{H^{k+1}}, \quad \|u - u_h\|_{L^2} \leq Ch^{k+1} |u|_{H^{k+1}}$$

Introduction

- In the previous year, MEF II, for some elliptic problems, we have seen how to show that

$$\|u - u_h\|_{H^1} \leq Ch^k |u|_{H^{k+1}}, \quad \|u - u_h\|_{L^2} \leq Ch^{k+1} |u|_{H^{k+1}}$$

- This means that the error can be arbitrarily reduced by reducing h .

But ! Higher regularity than that of the weak solution is needed.

This might be very **inefficient** for problems that involve:

Introduction

- In the previous year, MEF II, for some elliptic problems, we have seen how to show that

$$\|u - u_h\|_{H^1} \leq Ch^k |u|_{H^{k+1}}, \quad \|u - u_h\|_{L^2} \leq Ch^{k+1} |u|_{H^{k+1}}$$

- This means that the error can be arbitrarily reduced by reducing h .

But ! Higher regularity than that of the weak solution is needed.

This might be very **inefficient** for problems that involve:

- corner singularities

Introduction

- In the previous year, MEF II, for some elliptic problems, we have seen how to show that

$$\|u - u_h\|_{H^1} \leq Ch^k |u|_{H^{k+1}}, \quad \|u - u_h\|_{L^2} \leq Ch^{k+1} |u|_{H^{k+1}}$$

- This means that the error can be arbitrarily reduced by reducing h .

But ! Higher regularity than that of the weak solution is needed.

This might be very **inefficient** for problems that involve:

- corner singularities
- boundary or internal layers (singular perturbation problems)

Introduction

- In the previous year, MEF II, for some elliptic problems, we have seen how to show that

$$\|u - u_h\|_{H^1} \leq Ch^k |u|_{H^{k+1}}, \quad \|u - u_h\|_{L^2} \leq Ch^{k+1} |u|_{H^{k+1}}$$

- This means that the error can be arbitrarily reduced by reducing h .

But ! Higher regularity than that of the weak solution is needed.

This might be very **inefficient** for problems that involve:

- corner singularities
- boundary or internal layers (singular perturbation problems)

i.e, the error is mostly concentrated in one area of the grid.

Why adaptive? 1d Example

Question: given a continuous function $u : [0, 1] \rightarrow \mathbb{R}$, a partition $\mathcal{T}_h = \{x_n\}_{n=0}^N$ with $x_0 = 0, x_N = 1$, and a pw constant approximation u_h of u over \mathcal{T}_h , what is the best decay rate of $\|u - u_h\|_{L^\infty(0,1)}$?

Why adaptive? 1d Example

Question: given a continuous function $u : [0, 1] \rightarrow \mathbb{R}$, a partition $\mathcal{T}_h = \{x_n\}_{n=0}^N$ with $x_0 = 0, x_N = 1$, and a pw constant approximation u_h of u over \mathcal{T}_h , what is the best decay rate of $\|u - u_h\|_{L^\infty(0,1)}$?

Answer 1: W_∞^1 -Regularity. Let $u \in W_\infty^1(0, 1)$ and \mathcal{T}_h be quasi-uniform. Then $u_h^n(x) = u(x_{n-1})$ for $x_{n-1} \leq x < x_n$ satisfies

$$|u_h^n(x) - u(x)| = |u(x_{n-1}) - u(x)| \leq \int_x^{x_{n-1}} |u'(s)| ds \preceq h \|u'\|_{L^\infty(0,1)}$$

Why adaptive? 1d Example

Question: given a continuous function $u : [0, 1] \rightarrow \mathbb{R}$, a partition $\mathcal{T}_h = \{x_n\}_{n=0}^N$ with $x_0 = 0, x_N = 1$, and a pw constant approximation u_h of u over \mathcal{T}_h , what is the best decay rate of $\|u - u_h\|_{L^\infty(0,1)}$?

Answer 1: W_∞^1 -Regularity. Let $u \in W_\infty^1(0, 1)$ and \mathcal{T}_h be quasi-uniform. Then $u_h^n(x) = u(x_{n-1})$ for $x_{n-1} \leq x < x_n$ satisfies


$$|u_h^n(x) - u(x)| = |u(x_{n-1}) - u(x)| \leq \int_x^{x_{n-1}} |u'(s)| ds \preccurlyeq h \|u'\|_{L^\infty(0,1)}$$

Answer 2: W_1^1 -Regularity. Let $u \in W_1^1(0, 1)$. If x_n is defined by

$$\int_{x_{n-1}}^{x_n} |u'(s)| ds = \frac{1}{N} \|u'\|_{L^1(0,1)}$$

then

$$|u_h^n(x) - u(x)| = |u(x_{n-1}) - u(x)| \leq \int_x^{x_{n-1}} |u'(s)| ds \leq \frac{1}{N} \|u'\|_{L^1(0,1)}$$



$L2$ error (uniform partition) = $3.321646e - 02$

$L2$ error (adaptive partition) = $2.953768e - 02$

Sobolev Number

Definition

Let $\omega \subset \mathbb{R}^d$ be Lipschitz and bounded, $m \in \mathbb{N}$, $1 \leq p \leq \infty$. The Sobolev number of $W^{m,p}(\omega)$ is

$$\text{sob}(W^{m,p}) := m - \frac{d}{p}$$

Theorem (Quasi-local error estimate)

if $0 \leq t \leq s \leq k+1$ ($k \geq 1$ polynomial degree) and $1 \leq p, q \leq \infty$ satisfy $\text{sob}(W^{s,p}) > \text{sob}(W^{t,q})$, then for all $T \in \mathcal{T}$

$$\|D^t(v - I_{\mathcal{T}}v)\|_{L^q(T)} \lesssim h_T^{\text{sob}(W^{s,p}) - \text{sob}(W^{t,q})} \|D^s v\|_{L^p(\mathcal{N}_{\mathcal{T}}(T))}, \quad (1)$$

where $\mathcal{N}_{\mathcal{T}}(T)$ is a discrete neighborhood of T and $I_{\mathcal{T}}$ is a quasi interpolation operator (Clement or Scott-Zhang).

Piecewise Polynomial Interpolation

If $\text{sob}(W_p^s) > 0$, then v is Hölder continuous, $I_{\mathcal{T}}$ can be replaced by the Lagrange interpolation operator, and $\mathcal{N}_{\mathcal{T}}(T) = T$.

Quasi-uniform meshes: if $1 \leq s \leq k + 1$ and $u \in H^s(\Omega)$, then

$$\|\nabla(v - I_{\mathcal{T}}v)\|_{L^2(\Omega)} \lesssim |v|_{H^s(\Omega)} N^{-\frac{s-1}{d}}.$$

Optimal error decay: If $s = k + 1$, then

$$\|\nabla(v - I_{\mathcal{T}}v)\|_{L^2(\Omega)} \lesssim |v|_{H^{k+1}(\Omega)} N^{-\frac{k}{d}}.$$

Introduction

The aim of this lecture is,

Introduction

The aim of this lecture is,

- to derive a **computable** bound on the error,

Introduction

The aim of this lecture is,

- to derive a **computable** bound on the error,
- to demonstrate how such a bound may be implemented into an **adaptive mesh-refinement algorithm**, capable of reducing the error $u - u_h$ below a certain prescribed tolerance in an automated manner.

Introduction

The aim of this lecture is,

- to derive a **computable** bound on the error,
- to demonstrate how such a bound may be implemented into an **adaptive mesh-refinement algorithm**, capable of reducing the error $u - u_h$ below a certain prescribed tolerance in an automated manner.
- The approach is based on seeking a bound on $u - u_h$ in terms of the computed solution u_h rather than in terms of norms of the unknown analytical solution u .

Introduction

The aim of this lecture is,

- to derive a **computable** bound on the error,
- to demonstrate how such a bound may be implemented into an **adaptive mesh-refinement algorithm**, capable of reducing the error $u - u_h$ below a certain prescribed tolerance in an automated manner.
- The approach is based on seeking a bound on $u - u_h$ in terms of the computed solution u_h rather than in terms of norms of the unknown analytical solution u .
- A bound on the error in terms of u_h is referred to as an **a posteriori error bound**, due to the fact that it becomes **computable** only after the numerical solution u_h has been obtained.

Model problem in 1d

We consider the two-point boundary value problem

$$\begin{cases} -u'' + b(x)u' + c(x)u = f(x), & 0 < x < 1, \\ u(0) = 0, & u(1) = 0, \end{cases} \quad (2)$$

where

- $b \in W^{1,\infty}(0,1)$,
- $c \in L^\infty(0,1)$,
- $f \in L^2(0,1)$.

Letting

Model problem in 1d

We consider the two-point boundary value problem

$$\begin{cases} -u'' + b(x)u' + c(x)u = f(x), & 0 < x < 1, \\ u(0) = 0, & u(1) = 0, \end{cases} \quad (2)$$

where

- $b \in W^{1,\infty}(0,1)$,
- $c \in L^\infty(0,1)$,
- $f \in L^2(0,1)$.

Letting

$$a(w, v) = \int_0^1 [w'(x)v'(x) + b(x)w'(x)v(x) + c(x)w(x)v(x)] dx$$
$$\ell(v) = \int_0^1 f(x)v(x)dx$$

The weak formulation of this problem can be stated as follows:

$$\begin{cases} \text{find } u \in H_0^1(0, 1) \text{ such that} \\ a(u, v) = \ell(v), \quad \text{for all } v \in H_0^1(0, 1). \end{cases} \quad (3)$$

The weak formulation of this problem can be stated as follows:

$$\begin{cases} \text{find } u \in H_0^1(0, 1) \text{ such that} \\ a(u, v) = \ell(v), \quad \text{for all } v \in H_0^1(0, 1). \end{cases} \quad (3)$$

Lemma 1

Assuming that

$$c(x) - \frac{1}{2}b'(x) \geq 0, \quad \text{for } x \in (0, 1). \quad (4)$$

Then, there exists a unique weak solution, $u \in H_0^1(0, 1)$.

The finite element approximation

We consider a subdivision of the interval $[0, 1]$ by the points

$$0 = x_0 < x_1 < \dots < x_{N-1} < x_N = 1$$

We let $h_i = x_i - x_{i-1}$, $i = 1, \dots, N$, and put $h = \max_i h_i$ and defining the finite element space $V_h \subset H_0^1(0, 1)$ consisting of continuous piecewise linear function.

The finite element approximation

We consider a subdivision of the interval $[0, 1]$ by the points

$$0 = x_0 < x_1 < \dots < x_{N-1} < x_N = 1$$

We let $h_i = x_i - x_{i-1}$, $i = 1, \dots, N$, and put $h = \max_i h_i$ and defining the finite element space $V_h \subset H_0^1(0, 1)$ consisting of continuous piecewise linear function. The finite element approximation of the boundary value problem is:

$$\begin{cases} \text{find } u_h \in V_h \text{ such that} \\ a(u_h, v_h) = \ell(v_h), \quad \text{for all } v_h \in V_h. \end{cases} \quad (5)$$

We wish to derive an *a posteriori error* bound; that is, we aim to quantify the size of the global error $u - u_h$ in terms of the mesh parameter h and the computed solution u_h .

Theorem 1

Let u and u_h the solutions of the continuous problem (3) and the discrete problem (5). Then we have the computable a posteriori error bound,

$$\|u - u_h\|_{L_2(0,1)} \leq K_0 \left(\sum_{i=1}^N h_i^4 \|R(u_h)\|_{L_2(x_{i-1}, x_i)}^2 \right)^{1/2}, \quad (6)$$

where:

Theorem 1

Let u and u_h the solutions of the continuous problem (3) and the discrete problem (5). Then we have the computable a posteriori error bound,

$$\|u - u_h\|_{L_2(0,1)} \leq K_0 \left(\sum_{i=1}^N h_i^4 \|R(u_h)\|_{L_2(x_{i-1}, x_i)}^2 \right)^{1/2}, \quad (6)$$

where:

- K_0 is a positive constant depending only on the coefficients b and c .
- for $i = 1, \dots, N$,

Main theorem

Theorem 1

Let u and u_h the solutions of the continuous problem (3) and the discrete problem (5). Then we have the computable a posteriori error bound,

$$\|u - u_h\|_{L_2(0,1)} \leq K_0 \left(\sum_{i=1}^N h_i^4 \|R(u_h)\|_{L_2(x_{i-1}, x_i)}^2 \right)^{1/2}, \quad (6)$$

where:

- K_0 is a positive constant depending only on the coefficients b and c .
- for $i = 1, \dots, N$,

$$R(u_h)(x) = f(x) + u_h''(x) - b(x)u_h'(x) - c(x)u_h(x), \quad x \in (x_{i-1}, x_i).$$

The dual problem

To do so, we consider the following **auxiliary boundary value problem**:

$$\begin{cases} -z'' - (b(x)z)' + c(x)z = (u - u_h)(x), & 0 < x < 1, \\ z(0) = 0, \quad z(1) = 0, \end{cases} \quad (7)$$

called **the dual or adjoint problem**.

The dual problem

To do so, we consider the following **auxiliary boundary value problem**:

$$\begin{cases} -z'' - (b(x)z)' + c(x)z = (u - u_h)(x), & 0 < x < 1, \\ z(0) = 0, \quad z(1) = 0, \end{cases} \quad (7)$$

called **the dual or adjoint problem**.

Lemma 2

Suppose that z is the solution of the dual problem (7). Then, there exists a positive constant K , dependent only on b , and c , such that

$$\|z''\|_{L_2(0,1)} \leq K \|u - u_h\|_{L_2(0,1)}. \quad (8)$$

Proof of Lemma 2

We have

$$z'' = u_h - u - (bz)' + cz = u_h - u - bz' + (c - b')z,$$

and therefore,

Proof of Lemma 2

We have

$$z'' = u_h - u - (bz)'' + cz = u_h - u - bz' + (c - b')z,$$

and therefore,

$$\|z''\| \leq \|u - u_h\| + \|b\|_{L_\infty(0,1)} \|z'\| + \|c - b'\|_{L_\infty(0,1)} \|z\|. \quad (9)$$

We shall show that both $\|z'\|_{L_2(0,1)}$ and $\|z\|$ can be bounded in terms of $\|u - u_h\|$ and then, by virtue of (9), we shall deduce that the same is true of $\|z''\|$.

Proof of Lemma 2

We have

$$z'' = u_h - u - (bz)'' + cz = u_h - u - bz' + (c - b')z,$$

and therefore,

$$\|z''\| \leq \|u - u_h\| + \|b\|_{L_\infty(0,1)} \|z'\| + \|c - b'\|_{L_\infty(0,1)} \|z\|. \quad (9)$$

We shall show that both $\|z'\|_{L_2(0,1)}$ and $\|z\|$ can be bounded in terms of $\|u - u_h\|$ and then, by virtue of (9), we shall deduce that the same is true of $\|z''\|$.

Observe that

$$(-z'' - (bz)'' + cz, z) = (u - u_h, z).$$

Proof of Lemma 2

Integrating by parts and noting that $z(0) = 0$ and $z(1) = 0$ yields:

Proof of Lemma 2

Integrating by parts and noting that $z(0) = 0$ and $z(1) = 0$ yields:

$$\begin{aligned}(-z'' - (bz)') + cz, z) &= (z', z') + (bz, z') + (cz, z) \\&= \|z'\|^2 + \frac{1}{2} \int_0^1 b(x) [z^2(x)]' dx + \int_0^1 c(x) [z(x)]^2 dx\end{aligned}$$

Proof of Lemma 2

Integrating by parts and noting that $z(0) = 0$ and $z(1) = 0$ yields:

$$\begin{aligned}(-z'' - (bz)' + cz, z) &= (z', z') + (bz, z') + (cz, z) \\&= \|z'\|^2 + \frac{1}{2} \int_0^1 b(x) [z^2(x)]' dx + \int_0^1 c(x) [z(x)]^2 dx\end{aligned}$$

Integrating by parts, again, in the second term on the right gives

$$(-z'' - (bz)' + cz, z) = \|z'\|^2 - \frac{1}{2} \int_0^1 b'(x) [z^2(x)] dx + \int_0^1 c(x) [z(x)]^2 dx.$$

Hence,

Proof of Lemma 2

Integrating by parts and noting that $z(0) = 0$ and $z(1) = 0$ yields:

$$\begin{aligned}(-z'' - (bz)' + cz, z) &= (z', z') + (bz, z') + (cz, z) \\ &= \|z'\|^2 + \frac{1}{2} \int_0^1 b(x) [z^2(x)]' dx + \int_0^1 c(x) [z(x)]^2 dx\end{aligned}$$

Integrating by parts, again, in the second term on the right gives

$$(-z'' - (bz)' + cz, z) = \|z'\|^2 - \frac{1}{2} \int_0^1 b'(x) [z^2(x)] dx + \int_0^1 c(x) [z(x)]^2 dx.$$

Hence,

$$\|z'\|^2 + \int_0^1 \left(c(x) - \frac{1}{2} b'(x) \right) [z(x)]^2 dx = (u - u_h, z),$$

and thereby, noting (4),

Proof of Lemma 2

Integrating by parts and noting that $z(0) = 0$ and $z(1) = 0$ yields:

$$\begin{aligned}(-z'' - (bz)' + cz, z) &= (z', z') + (bz, z') + (cz, z) \\&= \|z'\|^2 + \frac{1}{2} \int_0^1 b(x) [z^2(x)]' dx + \int_0^1 c(x) [z(x)]^2 dx\end{aligned}$$

Integrating by parts, again, in the second term on the right gives

$$(-z'' - (bz)' + cz, z) = \|z'\|^2 - \frac{1}{2} \int_0^1 b'(x) [z^2(x)] dx + \int_0^1 c(x) [z(x)]^2 dx.$$

Hence,

$$\|z'\|^2 + \int_0^1 \left(c(x) - \frac{1}{2} b'(x) \right) [z(x)]^2 dx = (u - u_h, z),$$

and thereby, noting (4),

$$\|z'\|_{L^2(0,1)}^2 \leq (u - u_h, z) \leq \|u - u_h\|_{L_2(0,1)} \|z\|_{L_2(0,1)}. \quad (10)$$

By the Poincaré-Friedrichs inequality,

$$\|z\|_{L_2(0,1)}^2 \leq \frac{1}{2} \|z'\|_{L_2(0,1)}^2 .$$

By the Poincaré-Friedrichs inequality,

$$\|z\|_{L_2(0,1)}^2 \leq \frac{1}{2} \|z'\|_{L_2(0,1)}^2.$$

Thus, (10) gives

$$\|z\|_{L_2(0,1)} \leq \frac{1}{2} \|u - u_h\|_{L_2(0,1)} \tag{11}$$

By the Poincaré-Friedrichs inequality,

$$\|z\|_{L_2(0,1)}^2 \leq \frac{1}{2} \|z'\|_{L_2(0,1)}^2.$$

Thus, (10) gives

$$\|z\|_{L_2(0,1)} \leq \frac{1}{2} \|u - u_h\|_{L_2(0,1)} \quad (11)$$

Inserting this into the right-hand side of (10) yields

$$\|z'\|_{L_2(0,1)} \leq \frac{1}{\sqrt{2}} \|u - u_h\|_{L_2(0,1)} \quad (12)$$

By the Poincaré-Friedrichs inequality,

$$\|z\|_{L_2(0,1)}^2 \leq \frac{1}{2} \|z'\|_{L_2(0,1)}^2.$$

Thus, (10) gives

$$\|z\|_{L_2(0,1)} \leq \frac{1}{2} \|u - u_h\|_{L_2(0,1)} \quad (11)$$

Inserting this into the right-hand side of (10) yields

$$\|z'\|_{L_2(0,1)} \leq \frac{1}{\sqrt{2}} \|u - u_h\|_{L_2(0,1)} \quad (12)$$

Now we substitute (11) and (12) into (9) to deduce that

$$\|z''\|_{L_2(0,1)} \leq K \|u - u_h\|_{L_2(0,1)}. \quad (13)$$

By the Poincaré-Friedrichs inequality,

$$\|z\|_{L_2(0,1)}^2 \leq \frac{1}{2} \|z'\|_{L_2(0,1)}^2.$$

Thus, (10) gives

$$\|z\|_{L_2(0,1)} \leq \frac{1}{2} \|u - u_h\|_{L_2(0,1)} \quad (11)$$

Inserting this into the right-hand side of (10) yields

$$\|z'\|_{L_2(0,1)} \leq \frac{1}{\sqrt{2}} \|u - u_h\|_{L_2(0,1)} \quad (12)$$

Now we substitute (11) and (12) into (9) to deduce that

$$\|z''\|_{L_2(0,1)} \leq K \|u - u_h\|_{L_2(0,1)}. \quad (13)$$

Where

$$K = 1 + \frac{1}{\sqrt{2}} \|b\|_{L_\infty(0,1)} + \frac{1}{2} \|c - b'\|_{L_\infty(0,1)}.$$

Proof of Theorem 1

The L2-norm of the error is related to the solution of the dual problem via the relation:

$$\|u - u_h\|_{L_2(0,1)}^2 = a(u - u_h, z)$$

Proof of Theorem 1

The L2-norm of the error is related to the solution of the dual problem via the relation:

$$\|u - u_h\|_{L_2(0,1)}^2 = a(u - u_h, z)$$

Indeed,

$$\begin{aligned}\|u - u_h\|_{L_2(0,1)}^2 &= (u - u_h, u - u_h) = (u - u_h, -z'' - (bz)'+ cz) \\ &= a(u - u_h, z) .\end{aligned}$$

Proof of Theorem 1

The L2-norm of the error is related to the solution of the dual problem via the relation:

$$\|u - u_h\|_{L_2(0,1)}^2 = a(u - u_h, z)$$

Indeed,

$$\begin{aligned}\|u - u_h\|_{L_2(0,1)}^2 &= (u - u_h, u - u_h) = (u - u_h, -z'' - (bz)'+ cz) \\ &= a(u - u_h, z).\end{aligned}$$

By virtue of the Galerkin orthogonality property,

$$a(u - u_h, z_h) = 0 \quad \forall z_h \in V_h.$$

In particular, choosing $z_h = \mathcal{I}_h z \in V_h$, the continuous piecewise linear interpolant of the function z , we have that

$$a(u - u_h, \mathcal{I}_h z) = 0$$

Thus,

$$\begin{aligned}\|u - u_h\|_{L_2(0,1)}^2 &= a(u - u_h, z - \mathcal{I}_h z) = a(u, z - \mathcal{I}_h z) - a(u_h, z - \mathcal{I}_h z) \\ &= (f, z - \mathcal{I}_h z) - a(u_h, z - \mathcal{I}_h z).\end{aligned}\tag{14}$$

Thus,

$$\begin{aligned}\|u - u_h\|_{L_2(0,1)}^2 &= a(u - u_h, z - \mathcal{I}_h z) = a(u, z - \mathcal{I}_h z) - a(u_h, z - \mathcal{I}_h z) \\ &= (f, z - \mathcal{I}_h z) - a(u_h, z - \mathcal{I}_h z).\end{aligned}\tag{14}$$

Now,

$$\begin{aligned}a(u_h, z - \mathcal{I}_h z) &= \sum_{i=1}^N \int_{x_{i-1}}^{x_i} u_h'(x) (z - \mathcal{I}_h z)'(x) dx \\ &\quad + \sum_{i=1}^N \int_{x_{i-1}}^{x_i} b(x) u_h'(x) (z - \mathcal{I}_h z)(x) dx \\ &\quad + \sum_{i=1}^N \int_{x_{i-1}}^{x_i} c(x) u_h(x) (z - \mathcal{I}_h z)(x) dx.\end{aligned}$$

Integrating by parts in each of the $(N - 1)$ integrals in the first sum on the right-hand side, noting that

$$(z - \mathcal{I}_h z)(x_i) = 0, i = 0, \dots, N,$$

we deduce that

Integrating by parts in each of the $(N - 1)$ integrals in the first sum on the right-hand side, noting that

$$(z - \mathcal{I}_h z)(x_i) = 0, i = 0, \dots, N,$$

we deduce that

$$a(u_h, z - \mathcal{I}_h z) = \sum_{i=1}^N \int_{x_{i-1}}^{x_i} [-u_h''(x) + b(x)u_h'(x) + c(x)u_h(x)](z - \mathcal{I}_h z)(x)dx.$$

Integrating by parts in each of the $(N - 1)$ integrals in the first sum on the right-hand side, noting that

$$(z - \mathcal{I}_h z)(x_i) = 0, i = 0, \dots, N,$$

we deduce that

$$a(u_h, z - \mathcal{I}_h z) = \sum_{i=1}^N \int_{x_{i-1}}^{x_i} [-u_h''(x) + b(x)u_h'(x) + c(x)u_h(x)](z - \mathcal{I}_h z)(x)dx.$$

Further

$$(f, z - \mathcal{I}_h z) = \sum_{i=1}^N \int_{x_{i-1}}^{x_i} f(x)(z - \mathcal{I}_h z)(x)dx.$$

Integrating by parts in each of the $(N - 1)$ integrals in the first sum on the right-hand side, noting that

$$(z - \mathcal{I}_h z)(x_i) = 0, i = 0, \dots, N,$$

we deduce that

$$a(u_h, z - \mathcal{I}_h z) = \sum_{i=1}^N \int_{x_{i-1}}^{x_i} [-u_h''(x) + b(x)u_h'(x) + c(x)u_h(x)](z - \mathcal{I}_h z)(x)dx.$$

Further

$$(f, z - \mathcal{I}_h z) = \sum_{i=1}^N \int_{x_{i-1}}^{x_i} f(x)(z - \mathcal{I}_h z)(x)dx.$$

Substituting these two identities into (14), we deduce that

$$\|u - u_h\|_{L_2(0,1)}^2 = \sum_{i=1}^N \int_{x_{i-1}}^{x_i} R(u_h)(x)(z - \mathcal{I}_h z)(x)dx \quad (15)$$

where,

Integrating by parts in each of the $(N - 1)$ integrals in the first sum on the right-hand side, noting that

$$(z - \mathcal{I}_h z)(x_i) = 0, i = 0, \dots, N,$$

we deduce that

$$a(u_h, z - \mathcal{I}_h z) = \sum_{i=1}^N \int_{x_{i-1}}^{x_i} [-u_h''(x) + b(x)u_h'(x) + c(x)u_h(x)](z - \mathcal{I}_h z)(x)dx.$$

Further

$$(f, z - \mathcal{I}_h z) = \sum_{i=1}^N \int_{x_{i-1}}^{x_i} f(x)(z - \mathcal{I}_h z)(x)dx.$$

Substituting these two identities into (14), we deduce that

$$\|u - u_h\|_{L_2(0,1)}^2 = \sum_{i=1}^N \int_{x_{i-1}}^{x_i} R(u_h)(x)(z - \mathcal{I}_h z)(x)dx \quad (15)$$

where, for $i = 1, \dots, N$,

$$R(u_h)(x) = f(x) + u_h''(x) - b(x)u_h'(x) - c(x)u_h(x), \quad x \in (x_{i-1}, x_i).$$

The function $R(u_h)$ is called the finite element residual;

The function $R(u_h)$ is called the finite element residual;
it measures the extent to which u_h fails to satisfy the differential equation

$$-u'' + b(x)u' + c(x)u = f(x) \text{ on the interval } (0, 1).$$

The function $R(u_h)$ is called the finite element residual; it measures the extent to which u_h fails to satisfy the differential equation

$$-u'' + b(x)u' + c(x)u = f(x) \text{ on the interval } (0, 1).$$

Now, applying the Cauchy-Schwarz inequality on the right-hand side of (15) yields

$$\|u - u_h\|_{L_2(0,1)}^2 \leq \sum_{i=1}^N \|R(u_h)\|_{L_2(x_{i-1}, x_i)} \|z - \mathcal{I}_h z\|_{L_2(x_{i-1}, x_i)}.$$

The function $R(u_h)$ is called the finite element residual; it measures the extent to which u_h fails to satisfy the differential equation

$$-u'' + b(x)u' + c(x)u = f(x) \text{ on the interval } (0, 1).$$

Now, applying the Cauchy-Schwarz inequality on the right-hand side of (15) yields

$$\|u - u_h\|_{L_2(0,1)}^2 \leq \sum_{i=1}^N \|R(u_h)\|_{L_2(x_{i-1}, x_i)} \|z - \mathcal{I}_h z\|_{L_2(x_{i-1}, x_i)}.$$

Recalling from the proof of interpolation theorem that

$$\|z - \mathcal{I}_h z\|_{L_2(x_{i-1}, x_i)} \leq \left(\frac{h_i}{\pi}\right)^2 \|z''\|_{L_2(x_{i-1}, x_i)}, \quad i = 1, \dots, N,$$

The function $R(u_h)$ is called the finite element residual; it measures the extent to which u_h fails to satisfy the differential equation

$$-u'' + b(x)u' + c(x)u = f(x) \text{ on the interval } (0, 1).$$

Now, applying the Cauchy-Schwarz inequality on the right-hand side of (15) yields

$$\|u - u_h\|_{L_2(0,1)}^2 \leq \sum_{i=1}^N \|R(u_h)\|_{L_2(x_{i-1}, x_i)} \|z - \mathcal{I}_h z\|_{L_2(x_{i-1}, x_i)}.$$

Recalling from the proof of interpolation theorem that

$$\|z - \mathcal{I}_h z\|_{L_2(x_{i-1}, x_i)} \leq \left(\frac{h_i}{\pi}\right)^2 \|z''\|_{L_2(x_{i-1}, x_i)}, \quad i = 1, \dots, N,$$

and consequently,

$$\|u - u_h\|_{L_2(0,1)}^2 \leq \frac{1}{\pi^2} \left(\sum_{i=1}^N h_i^4 \|R(u_h)\|_{L_2(x_{i-1}, x_i)}^2 \right)^{1/2} \|z''\|_{L_2(0,1)}. \quad (16)$$

Inserting (13) into (16), we arrive at our final result, the computable a posteriori error bound,

$$\|u - u_h\|_{L_2(0,1)} \leq K_0 \left(\sum_{i=1}^N h_i^4 \|R(u_h)\|_{L_2(x_{i-1}, x_i)}^2 \right)^{1/2}, \quad (17)$$

where $K_0 = K/\pi^2$.

Inserting (13) into (16), we arrive at our final result, the computable a posteriori error bound,

$$\|u - u_h\|_{L_2(0,1)} \leq K_0 \left(\sum_{i=1}^N h_i^4 \|R(u_h)\|_{L_2(x_{i-1}, x_i)}^2 \right)^{1/2}, \quad (17)$$

where $K_0 = K/\pi^2$.

The name a posteriori stems from the fact that (17) can only be employed to quantify the size of the approximation error that has been committed in the course of the computation after u_h has been computed.

Inserting (13) into (16), we arrive at our final result, the computable a posteriori error bound,

$$\|u - u_h\|_{L_2(0,1)} \leq K_0 \left(\sum_{i=1}^N h_i^4 \|R(u_h)\|_{L_2(x_{i-1}, x_i)}^2 \right)^{1/2}, \quad (17)$$

where $K_0 = K/\pi^2$.

The name a posteriori stems from the fact that (17) can only be employed to quantify the size of the approximation error that has been committed in the course of the computation after u_h has been computed. In the next section we shall describe the construction of an adaptive mesh refinement algorithm based on the bound (17).

Adaptive method

Suppose that TOL is a prescribed tolerance and that our aim is to compute a finite element approximation u_h to the unknown solution u (with the same definition of u and u_h as in the previous section) so that

$$\|u - u_h\|_{L_2(0,1)} \leq TOL.$$

Adaptive method

Suppose that TOL is a prescribed tolerance and that our aim is to compute a finite element approximation u_h to the unknown solution u (with the same definition of u and u_h as in the previous section) so that

$$\|u - u_h\|_{L_2(0,1)} \leq TOL.$$

We shall use the a posteriori error bound (17) to achieve this goal by successively refining the subdivision, and computing a succession of numerical solutions u_h on these subdivisions, until the stopping criterion

Adaptive method

Suppose that TOL is a prescribed tolerance and that our aim is to compute a finite element approximation u_h to the unknown solution u (with the same definition of u and u_h as in the previous section) so that

$$\|u - u_h\|_{L_2(0,1)} \leq TOL.$$

We shall use the a posteriori error bound (17) to achieve this goal by successively refining the subdivision, and computing a succession of numerical solutions u_h on these subdivisions, until the stopping criterion

$$K_0 \left(\sum_{i=1}^N h_i^4 \|R(u_h)\|_{L_2(x_{i-1}, x_i)}^2 \right)^{1/2} \leq TOL$$

is satisfied.

- Choose an initial subdivision

$$\mathcal{T}_0 : \quad 0 = x_0^{(0)} < x_1^{(0)} < \dots < x_{N_0-1}^{(0)} < x_{N_0}^{(0)} = 1$$

of the interval $[0, 1]$, with $h_i^{(0)} = x_i^{(0)} - x_{i-1}^{(0)}$ for $i = 1, \dots, N_0$,

- Choose an initial subdivision

$$\mathcal{T}_0 : \quad 0 = x_0^{(0)} < x_1^{(0)} < \dots < x_{N_0-1}^{(0)} < x_{N_0}^{(0)} = 1$$

of the interval $[0, 1]$, with $h_i^{(0)} = x_i^{(0)} - x_{i-1}^{(0)}$ for $i = 1, \dots, N_0$,

- Compute the corresponding solution $u_{h^{(0)}} \in V_{h^{(0)}}$.

- Choose an initial subdivision

$$\mathcal{T}_0 : \quad 0 = x_0^{(0)} < x_1^{(0)} < \dots < x_{N_0-1}^{(0)} < x_{N_0}^{(0)} = 1$$

of the interval $[0, 1]$, with $h_i^{(0)} = x_i^{(0)} - x_{i-1}^{(0)}$ for $i = 1, \dots, N_0$,

- Compute the corresponding solution $u_{h^{(0)}} \in V_{h^{(0)}}$.
- Given a computed solution $u_{h^{(m)}} \in V_{h^{(m)}}$ for some $m \geq 0$, defined on a subdivision \mathcal{T}_m , stop if

$$K_0 \left(\sum_{i=1}^{N_m} \left(h_i^{(m)} \right)^4 \|R(u_{h^{(m)}})\|_{L_2(x_{i-1}^{(m)}, x_i^{(m)})}^2 \right)^{1/2} \leq TOL.$$

- Choose an initial subdivision

$$\mathcal{T}_0 : \quad 0 = x_0^{(0)} < x_1^{(0)} < \dots < x_{N_0-1}^{(0)} < x_{N_0}^{(0)} = 1$$

of the interval $[0, 1]$, with $h_i^{(0)} = x_i^{(0)} - x_{i-1}^{(0)}$ for $i = 1, \dots, N_0$,

- Compute the corresponding solution $u_{h^{(0)}} \in V_{h^{(0)}}$.
- Given a computed solution $u_{h^{(m)}} \in V_{h^{(m)}}$ for some $m \geq 0$, defined on a subdivision \mathcal{T}_m , stop if

$$K_0 \left(\sum_{i=1}^{N_m} \left(h_i^{(m)} \right)^4 \|R(u_{h^{(m)}})\|_{L_2(x_{i-1}^{(m)}, x_i^{(m)})}^2 \right)^{1/2} \leq TOL.$$

- If not, then determine a new subdivision

$$\mathcal{T}_{m+1} : \quad 0 = x_0^{(m+1)} < x_1^{(m+1)} < \dots < x_{N_{m+1}-1}^{(m+1)} < x_{N_{m+1}}^{(m+1)} = 1$$

and an associated finite element space $V_{h^{(m+1)}}$, such that

- Choose an initial subdivision

$$\mathcal{T}_0 : \quad 0 = x_0^{(0)} < x_1^{(0)} < \dots < x_{N_0-1}^{(0)} < x_{N_0}^{(0)} = 1$$

of the interval $[0, 1]$, with $h_i^{(0)} = x_i^{(0)} - x_{i-1}^{(0)}$ for $i = 1, \dots, N_0$,

- Compute the corresponding solution $u_{h^{(0)}} \in V_{h^{(0)}}$.
- Given a computed solution $u_{h^{(m)}} \in V_{h^{(m)}}$ for some $m \geq 0$, defined on a subdivision \mathcal{T}_m , stop if

$$K_0 \left(\sum_{i=1}^{N_m} \left(h_i^{(m)} \right)^4 \|R(u_{h^{(m)}})\|_{L_2(x_{i-1}^{(m)}, x_i^{(m)})}^2 \right)^{1/2} \leq TOL.$$

- If not, then determine a new subdivision

$$\mathcal{T}_{m+1} : \quad 0 = x_0^{(m+1)} < x_1^{(m+1)} < \dots < x_{N_{m+1}-1}^{(m+1)} < x_{N_{m+1}}^{(m+1)} = 1$$

and an associated finite element space $V_{h^{(m+1)}}$, such that

$$K_0 \left(\sum_{i=1}^{N_{m+1}} \left(h_i^{(m+1)} \right)^4 \|R(u_{h^{(m)}})\|_{L_2(x_{i-1}^{(m+1)}, x_i^{(m+1)})}^2 \right)^{1/2} = TOL,$$

and continue.

Let us consider the second-order ordinary differential equation

$$\begin{aligned} - (a(x)u')' + b(x)u' + c(x)u &= f(x), \quad x \in (0, 1) \\ u(0) &= 0, \quad u(1) = 0 \end{aligned}$$

Suppose, for example, that

$$a(x) \equiv 1, \quad b(x) \equiv 20, \quad c(x) \equiv 10 \quad \text{and} \quad f(x) \equiv 1$$

In this case, the analytical solution, u , can be expressed in closed form:

$$u(x) = C_1 e^{\lambda_1 x} + C_2 e^{\lambda_2 x} + \frac{1}{10}$$

where λ_1 and λ_2 are the two roots of the characteristic polynomial of the differential equation,


$$-\lambda^2 + 20\lambda + 10 = 0,$$


i.e.,


$$\lambda_1 = 10 + \sqrt{110}, \quad \lambda_2 = 10 - \sqrt{110}$$


and C_1 and C_2 are constants chosen so as to ensure that $u(0) = 0$ and $u(1) = 0$; hence

$$C_1 = \frac{e^{\lambda_2} - 1}{10(e^{\lambda_1} - e^{\lambda_2})}, \quad C_2 = \frac{1 - e^{\lambda_1}}{10(e^{\lambda_1} - e^{\lambda_2})}$$









Conclusion

- .
- .
- .