

# Lecture 5: Numerical quadratures

Pr. Ismail Merabet

Univ. of K-M-Ouargla

November 23, 2024

# Contents

- 1 Introduction
- 2 Newton-Cotes formulae
- 3 Error estimate
- 4 Composite formulae

# Introduction

In this lecture, we derive formulae for **numerical integration** (or numerical quadrature rules) and estimate the associated approximation error.

# Introduction

In this lecture, we derive formulae for **numerical integration** (or numerical quadrature rules) and estimate the associated approximation error.

Let  $f$  be a real-valued function, defined and continuous on a closed real interval  $[a, b]$ , and suppose that we have to evaluate the integral

$$\int_a^b f(x) dx$$

# Introduction

In this lecture, we derive formulae for **numerical integration** (or numerical quadrature rules) and estimate the associated approximation error.

Let  $f$  be a real-valued function, defined and continuous on a closed real interval  $[a, b]$ , and suppose that we have to evaluate the integral

$$\int_a^b f(x) dx$$

Since **polynomials are easy to integrate**, the idea, roughly speaking, is to approximate the function  $f$  by its Lagrange interpolation polynomial  $p_n$  of degree  $n$ , and integrate  $p_n$  instead.

# Introduction

In this lecture, we derive formulae for **numerical integration** (or numerical quadrature rules) and estimate the associated approximation error.

Let  $f$  be a real-valued function, defined and continuous on a closed real interval  $[a, b]$ , and suppose that we have to evaluate the integral

$$\int_a^b f(x) dx$$

Since **polynomials are easy to integrate**, the idea, roughly speaking, is to approximate the function  $f$  by its Lagrange interpolation polynomial  $p_n$  of degree  $n$ , and integrate  $p_n$  instead. Thus,

$$\int_a^b f(x) dx \approx \int_a^b p_n(x) dx \tag{1}$$

# Newton-Cotes formulae

For a positive integer  $n$ , let  $x_i, i = 0, \dots, n$ , denote the interpolation points; for the sake of simplicity, we shall assume that these are equally spaced, namely,

$$x_i = a + ih \quad \text{for } i = 0, \dots, n, \quad \text{where } h = (b - a)/n.$$

# Newton-Cotes formulae

For a positive integer  $n$ , let  $x_i, i = 0, \dots, n$ , denote the interpolation points; for the sake of simplicity, we shall assume that these are equally spaced, namely,

$$x_i = a + ih \quad \text{for } i = 0, \dots, n, \quad \text{where } h = (b - a)/n.$$

The Lagrange interpolation polynomial of degree  $n$  for the function  $f$ , with these interpolation points, is of the form

$$p_n(x) = \sum_{k=0}^n L_k(x) f(x_k)$$



# Newton-Cotes formulae

For a positive integer  $n$ , let  $x_i, i = 0, \dots, n$ , denote the interpolation points; for the sake of simplicity, we shall assume that these are equally spaced, namely,

$$x_i = a + ih \quad \text{for } i = 0, \dots, n, \quad \text{where } h = (b - a)/n.$$

The Lagrange interpolation polynomial of degree  $n$  for the function  $f$ , with these interpolation points, is of the form

$$p_n(x) = \sum_{k=0}^n L_k(x) f(x_k)$$

Inserting this into the right-hand side of (1) yields

$$\int_a^b f(x) dx \approx \sum_{k=0}^n w_k f(x_k) \tag{2}$$

# Newton-Cotes formulae

For a positive integer  $n$ , let  $x_i, i = 0, \dots, n$ , denote the interpolation points; for the sake of simplicity, we shall assume that these are equally spaced, namely,

$$x_i = a + ih \quad \text{for } i = 0, \dots, n, \quad \text{where } h = (b - a)/n.$$

The Lagrange interpolation polynomial of degree  $n$  for the function  $f$ , with these interpolation points, is of the form

$$p_n(x) = \sum_{k=0}^n L_k(x) f(x_k)$$

Inserting this into the right-hand side of (1) yields

$$\int_a^b f(x) dx \approx \sum_{k=0}^n w_k f(x_k) \tag{2}$$

where

$$w_k = \int_a^b L_k(x) dx, \quad k = 0, \dots, n$$

## Definition

Numerical quadrature rules, with **equally spaced quadrature points**, are called **Newton-Cotes formulae**.

In order to illustrate the general idea, we consider two simple examples:

- Trapezium rule.
- Simpson's rule.

# Trapezium rule

In this case we take  $n = 1$ :  $x_0 = a$  and  $x_1 = b$ ,

# Trapezium rule

In this case we take  $n = 1$ :  $x_0 = a$  and  $x_1 = b$ , the Lagrange interpolation polynomial of degree 1 for the function  $f$  is simply

# Trapezium rule

In this case we take  $n = 1$ :  $x_0 = a$  and  $x_1 = b$ , the Lagrange interpolation polynomial of degree 1 for the function  $f$  is simply

$$\begin{aligned}p_1(x) &= L_0(x)f(x_0) + L_1(x)f(x_1) \\&= \frac{x - x_1}{x_0 - x_1}f(x_0) + \frac{x - x_0}{x_1 - x_0}f(x_1) \\&= \frac{1}{b - a}[(b - x)f(a) + (x - a)f(b)]\end{aligned}$$

# Trapezium rule

In this case we take  $n = 1$ :  $x_0 = a$  and  $x_1 = b$ , the Lagrange interpolation polynomial of degree 1 for the function  $f$  is simply

$$\begin{aligned} p_1(x) &= L_0(x)f(x_0) + L_1(x)f(x_1) \\ &= \frac{x - x_1}{x_0 - x_1}f(x_0) + \frac{x - x_0}{x_1 - x_0}f(x_1) \\ &= \frac{1}{b - a}[(b - x)f(a) + (x - a)f(b)] \end{aligned}$$

Integrating  $p_1(x)$  from  $a$  to  $b$  yields

$$\int_a^b f(x) dx \approx \frac{b - a}{2}[f(a) + f(b)].$$

# Trapezium rule

In this case we take  $n = 1$ :  $x_0 = a$  and  $x_1 = b$ , the Lagrange interpolation polynomial of degree 1 for the function  $f$  is simply

$$\begin{aligned} p_1(x) &= L_0(x)f(x_0) + L_1(x)f(x_1) \\ &= \frac{x - x_1}{x_0 - x_1}f(x_0) + \frac{x - x_0}{x_1 - x_0}f(x_1) \\ &= \frac{1}{b - a}[(b - x)f(a) + (x - a)f(b)] \end{aligned}$$

Integrating  $p_1(x)$  from  $a$  to  $b$  yields

$$\int_a^b f(x) dx \approx \frac{b - a}{2}[f(a) + f(b)].$$

This numerical integration formula is called **the trapezium rule**.



# Simpson's rule

A slightly more sophisticated quadrature rule is obtained by taking  $n = 2$ .

# Simpson's rule

A slightly more sophisticated quadrature rule is obtained by taking  $n = 2$ . In this case:

$$x_0 = a, x_1 = (a + b)/2 \text{ and } x_2 = b$$

the Lagrange polynomial of degree 2, with these interpolation points, for the function  $f$  is:

$$\begin{aligned} p_2(x) &= L_0(x)f(x_0) + L_1(x)f(x_1) + L_2(x)f(x_2) \\ &= \frac{(x - x_1)(x - x_2)}{(x_0 - x_1)(x_0 - x_2)}f(x_0) + \frac{(x - x_0)(x - x_2)}{(x_1 - x_0)(x_1 - x_2)}f(x_1) \\ &\quad + \frac{(x - x_0)(x - x_1)}{(x_2 - x_0)(x_2 - x_1)}f(x_2). \end{aligned}$$

# Simpson's rule

A slightly more sophisticated quadrature rule is obtained by taking  $n = 2$ . In this case:

$$x_0 = a, x_1 = (a + b)/2 \text{ and } x_2 = b$$

the Lagrange polynomial of degree 2, with these interpolation points, for the function  $f$  is:

$$\begin{aligned} p_2(x) &= L_0(x)f(x_0) + L_1(x)f(x_1) + L_2(x)f(x_2) \\ &= \frac{(x - x_1)(x - x_2)}{(x_0 - x_1)(x_0 - x_2)}f(x_0) + \frac{(x - x_0)(x - x_2)}{(x_1 - x_0)(x_1 - x_2)}f(x_1) \\ &\quad + \frac{(x - x_0)(x - x_1)}{(x_2 - x_0)(x_2 - x_1)}f(x_2). \end{aligned}$$

Integrating  $p_2(x)$  from  $a$  to  $b$  gives

$$\int_a^b f(x) dx \approx \frac{b - a}{6} \left[ f(a) + 4f\left(\frac{a + b}{2}\right) + f(b) \right].$$

This numerical integration formula is known as **Simpson's rule**.

# Error estimate

Our next task is to estimate the size of the error in the numerical integration formula (2), that is, the error that has been committed by integrating the Lagrange interpolation polynomial of  $f$  instead of the function  $f$  itself.

Our next task is to estimate the size of the error in the numerical integration formula (2), that is, the error that has been committed by integrating the Lagrange interpolation polynomial of  $f$  instead of the function  $f$  itself. The error in (2) is defined by

$$E_n(f) = \int_a^b f(x) dx - \sum_{k=0}^n w_k f(x_k)$$

Our next task is to estimate the size of the error in the numerical integration formula (2), that is, the error that has been committed by integrating the Lagrange interpolation polynomial of  $f$  instead of the function  $f$  itself. The error in (2) is defined by

$$E_n(f) = \int_a^b f(x) dx - \sum_{k=0}^n w_k f(x_k)$$

The next theorem provides a useful bound on  $E_n(f)$  under the additional hypothesis that the function  $f$  is sufficiently smooth.

## Theorem 1

Suppose that  $f$  is a real-valued function defined on the interval  $[a, b]$ , and let  $f^{(n+1)}$  be continuous on  $[a, b]$ . Then,

$$|E_n(f)| \leq \frac{M_{n+1}}{(n+1)!} \int_a^b |\pi_{n+1}(x)| dx \quad (3)$$

where :

- $M_{n+1} = \max_{\zeta \in [a, b]} |f^{(n+1)}(\zeta)|$  and
- $\pi_{n+1}(x) = (x - x_0) \dots (x - x_n)$ .

# Proof:

Recalling the inequality:

$$|f(x) - p_n(x)| \leq \frac{M_{n+1}}{(n+1)!} |\pi_{n+1}(x)| \quad (4)$$

and the definition of the weights  $w_k$  in (2),



# Proof:

Recalling the inequality:

$$|f(x) - p_n(x)| \leq \frac{M_{n+1}}{(n+1)!} |\pi_{n+1}(x)| \quad (4)$$

and the definition of the weights  $w_k$  in (2), we can rewrite  $E_n(f)$  as follows:

$$\begin{aligned} E_n(f) &= \int_a^b f(x) dx - \int_a^b \left( \sum_{k=0}^n L_k(x) f(x_k) \right) dx \\ &= \int_a^b (f(x) - p_n(x)) dx \end{aligned}$$

# Proof:

Recalling the inequality:

$$|f(x) - p_n(x)| \leq \frac{M_{n+1}}{(n+1)!} |\pi_{n+1}(x)| \quad (4)$$

and the definition of the weights  $w_k$  in (2), we can rewrite  $E_n(f)$  as follows:

$$\begin{aligned} E_n(f) &= \int_a^b f(x) dx - \int_a^b \left( \sum_{k=0}^n L_k(x) f(x_k) \right) dx \\ &= \int_a^b (f(x) - p_n(x)) dx \end{aligned}$$

Thus,

$$|E_n(f)| \leq \int_a^b |f(x) - p_n(x)| dx$$

# Proof:

Recalling the inequality:

$$|f(x) - p_n(x)| \leq \frac{M_{n+1}}{(n+1)!} |\pi_{n+1}(x)| \quad (4)$$

and the definition of the weights  $w_k$  in (2), we can rewrite  $E_n(f)$  as follows:

$$\begin{aligned} E_n(f) &= \int_a^b f(x) dx - \int_a^b \left( \sum_{k=0}^n L_k(x) f(x_k) \right) dx \\ &= \int_a^b (f(x) - p_n(x)) dx \end{aligned}$$

Thus,

$$|E_n(f)| \leq \int_a^b |f(x) - p_n(x)| dx$$

The desired error estimate (3) follows by inserting (4) into the right-hand side of this inequality.

Let us apply Theorem 1 to estimate the size of the error that has been committed by applying the trapezium rule to the integral  $\int_a^b f(x) dx$ . In this case (3) reduces to

$$\begin{aligned} |E_1(f)| &\leq \frac{1}{2} M_2 \int_a^b |(x-a)(x-b)| dx \\ &= \frac{1}{2} M_2 \int_a^b (b-x)(x-a) dx \\ &= \frac{1}{12} (b-a)^3 M_2 \end{aligned} \tag{14}$$

Let us apply Theorem 1 to estimate the size of the error that has been committed by applying the trapezium rule to the integral  $\int_a^b f(x) dx$ . In this case (3) reduces to

$$\begin{aligned} |E_1(f)| &\leq \frac{1}{2} M_2 \int_a^b |(x-a)(x-b)| dx \\ &= \frac{1}{2} M_2 \int_a^b (b-x)(x-a) dx \\ &= \frac{1}{12} (b-a)^3 M_2 \end{aligned} \tag{14}$$

An analogous but slightly more tedious calculation shows that, for Simpson's rule,

$$|E_2(f)| \leq \frac{1}{6} M_3 \int_a^b \left| (x-a) \left( x - \frac{a+b}{2} \right) (x-b) \right| dx \tag{5}$$

$$= \frac{1}{192} M_3 (b-a)^4 \tag{6}$$

Unfortunately, (5) gives a very pessimistic estimate of the error in Simpson's rule;

Unfortunately, (5) gives a very pessimistic estimate of the error in Simpson's rule; indeed, it is easy to verify that, whenever  $f$  is a polynomial of degree 3 we have  $E_2(f) = 0$  while the right-hand side of (5) is a positive real number.

Unfortunately, (5) gives a very pessimistic estimate of the error in Simpson's rule; indeed, it is easy to verify that, whenever  $f$  is a polynomial of degree 3 we have  $E_2(f) = 0$  while the right-hand side of (5) is a positive real number. The next lemma will allow us to sharpen this crude bound.



Unfortunately, (5) gives a very pessimistic estimate of the error in Simpson's rule; indeed, it is easy to verify that, whenever  $f$  is a polynomial of degree 3 we have  $E_2(f) = 0$  while the right-hand side of (5) is a positive real number. The next lemma will allow us to sharpen this crude bound.

### Lemma 1

Suppose that  $f$  is a real-valued function defined on the interval  $[a, b]$  and  $f^{(4)}$  is a continuous function on  $[a, b]$ . Then

$$\int_a^b f(x) dx - \frac{b-a}{6} \left[ f(a) + f\left(\frac{a+b}{2}\right) + f(b) \right] = -\frac{(b-a)^5}{2880} f^{(4)}(\xi) \quad (7)$$

for some  $\xi$  in  $(a, b)$ .

# Proof:

Performing the change of variables

$$x = \frac{a+b}{2} + \frac{b-a}{2}t, \quad t \in [-1, 1]$$

and defining the function  $t \mapsto F(t)$  by  $F(t) = f(x)$ , we have that

$$\int_a^b f(x) dx = \frac{b-a}{6} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] \quad (8)$$

$$= \frac{b-a}{2} \left( \int_{-1}^1 F(t) dt - \frac{1}{3}[F(-1) + 4F(0) + F(1)] \right) \quad (9)$$

# Proof:

Performing the change of variables

$$x = \frac{a+b}{2} + \frac{b-a}{2}t, \quad t \in [-1, 1]$$

and defining the function  $t \mapsto F(t)$  by  $F(t) = f(x)$ , we have that

$$\int_a^b f(x) dx = \frac{b-a}{6} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] \quad (8)$$

$$= \frac{b-a}{2} \left( \int_{-1}^1 F(t) dt - \frac{1}{3}[F(-1) + 4F(0) + F(1)] \right) \quad (9)$$

Let us introduce

$$G(t) = \int_{-t}^t F(\tau) d\tau - \frac{t}{3}[F(-t) + 4F(0) + F(t)], \quad t \in [-1, 1]$$

then the right-hand side of (9) is simply  $\frac{1}{2}(b-a)G(1)$ .

# Proof:

Performing the change of variables

$$x = \frac{a+b}{2} + \frac{b-a}{2}t, \quad t \in [-1, 1]$$

and defining the function  $t \mapsto F(t)$  by  $F(t) = f(x)$ , we have that

$$\int_a^b f(x) dx = \frac{b-a}{6} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] \quad (8)$$

$$= \frac{b-a}{2} \left( \int_{-1}^1 F(t) dt - \frac{1}{3}[F(-1) + 4F(0) + F(1)] \right) \quad (9)$$

Let us introduce

$$G(t) = \int_{-t}^t F(\tau) d\tau - \frac{t}{3}[F(-t) + 4F(0) + F(t)], \quad t \in [-1, 1]$$

then the right-hand side of (9) is simply  $\frac{1}{2}(b-a)G(1)$ .

The remainder of the proof is devoted to showing that  $\frac{1}{2}(b-a)G(1)$  is, in turn, equal to the right-hand side of (7) for some  $\xi$  in  $(a, b)$ .

# Proof:

Performing the change of variables

$$x = \frac{a+b}{2} + \frac{b-a}{2}t, \quad t \in [-1, 1]$$

and defining the function  $t \mapsto F(t)$  by  $F(t) = f(x)$ , we have that

$$\int_a^b f(x) dx = \frac{b-a}{6} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] \quad (8)$$

$$= \frac{b-a}{2} \left( \int_{-1}^1 F(t) dt - \frac{1}{3}[F(-1) + 4F(0) + F(1)] \right) \quad (9)$$

Let us introduce

$$G(t) = \int_{-t}^t F(\tau) d\tau - \frac{t}{3}[F(-t) + 4F(0) + F(t)], \quad t \in [-1, 1]$$

then the right-hand side of (9) is simply  $\frac{1}{2}(b-a)G(1)$ .

The remainder of the proof is devoted to showing that  $\frac{1}{2}(b-a)G(1)$  is, in turn, equal to the right-hand side of (7) for some  $\xi$  in  $(a, b)$ . For this purpose, we define

We shall apply Rolle's Theorem four times to the function  $H$ .

We shall apply Rolle's Theorem four times to the function  $H$ . Noting that  $H(0) = H(1) = 0$ , we deduce that there exists  $\zeta_1 \in (0, 1)$  such that  $H'(\zeta_1) = 0$ .

We shall apply Rolle's Theorem four times to the function  $H$ . Noting that  $H(0) = H(1) = 0$ , we deduce that there exists  $\zeta_1 \in (0, 1)$  such that  $H'(\zeta_1) = 0$ . However, also,  $H'(0) = 0$ , so there exists  $\zeta_2 \in (0, \zeta_1)$  such that  $H''(\zeta_2) = 0$ ; but,  $H''(0) = 0$ ,



We shall apply Rolle's Theorem four times to the function  $H$ . Noting that  $H(0) = H(1) = 0$ , we deduce that there exists  $\zeta_1 \in (0, 1)$  such that  $H'(\zeta_1) = 0$ . However, also,  $H'(0) = 0$ , so there exists  $\zeta_2 \in (0, \zeta_1)$  such that  $H''(\zeta_2) = 0$ ; but,  $H''(0) = 0$ , so there is  $\zeta_3 \in (0, \zeta_2)$  such that  $H'''(\zeta_3) = 0$ .

We shall apply Rolle's Theorem four times to the function  $H$ . Noting that  $H(0) = H(1) = 0$ , we deduce that there exists  $\zeta_1 \in (0, 1)$  such that  $H'(\zeta_1) = 0$ . However, also,  $H'(0) = 0$ , so there exists  $\zeta_2 \in (0, \zeta_1)$  such that  $H''(\zeta_2) = 0$ ; but,  $H''(0) = 0$ , so there is  $\zeta_3 \in (0, \zeta_2)$  such that  $H'''(\zeta_3) = 0$ . Finally, noting that  $H'''(0) = 0$ , it follows that there is  $\zeta_4 \in (0, \zeta_3)$  such that  $H^{(4)}(\zeta_4) = 0$ .

We shall apply Rolle's Theorem four times to the function  $H$ . Noting that  $H(0) = H(1) = 0$ , we deduce that there exists  $\zeta_1 \in (0, 1)$  such that  $H'(\zeta_1) = 0$ . However, also,  $H'(0) = 0$ , so there exists  $\zeta_2 \in (0, \zeta_1)$  such that  $H''(\zeta_2) = 0$ ; but,  $H''(0) = 0$ , so there is  $\zeta_3 \in (0, \zeta_2)$  such that  $H'''(\zeta_3) = 0$ . Finally, noting that  $H'''(0) = 0$ , it follows that there is  $\zeta_4 \in (0, \zeta_3)$  such that  $H^{(4)}(\zeta_4) = 0$ . Thus,

$$H^{(4)}(\zeta_4) = G^{(4)}(\zeta_4) - 120\zeta_4 G(1) = 0$$

We shall apply Rolle's Theorem four times to the function  $H$ . Noting that  $H(0) = H(1) = 0$ , we deduce that there exists  $\zeta_1 \in (0, 1)$  such that  $H'(\zeta_1) = 0$ . However, also,  $H'(0) = 0$ , so there exists  $\zeta_2 \in (0, \zeta_1)$  such that  $H''(\zeta_2) = 0$ ; but,  $H''(0) = 0$ , so there is  $\zeta_3 \in (0, \zeta_2)$  such that  $H'''(\zeta_3) = 0$ . Finally, noting that  $H'''(0) = 0$ , it follows that there is  $\zeta_4 \in (0, \zeta_3)$  such that  $H^{(4)}(\zeta_4) = 0$ . Thus,

$$H^{(4)}(\zeta_4) = G^{(4)}(\zeta_4) - 120\zeta_4 G(1) = 0$$

Now

$$G^{(4)}(t) = -\frac{1}{3} (F'''(t) - F'''(-t)) - \frac{t}{3} (F^{(4)}(t) + F^{(4)}(-t))$$

and therefore

$$G(1) = -\frac{1}{360\zeta_4} (F'''(\zeta_4) - F'''(-\zeta_4)) - \frac{1}{360} (F^{(4)}(\zeta_4) + F^{(4)}(-\zeta_4))$$

We shall apply Rolle's Theorem four times to the function  $H$ . Noting that  $H(0) = H(1) = 0$ , we deduce that there exists  $\zeta_1 \in (0, 1)$  such that  $H'(\zeta_1) = 0$ . However, also,  $H'(0) = 0$ , so there exists  $\zeta_2 \in (0, \zeta_1)$  such that  $H''(\zeta_2) = 0$ ; but,  $H''(0) = 0$ , so there is  $\zeta_3 \in (0, \zeta_2)$  such that  $H'''(\zeta_3) = 0$ . Finally, noting that  $H'''(0) = 0$ , it follows that there is  $\zeta_4 \in (0, \zeta_3)$  such that  $H^{(4)}(\zeta_4) = 0$ . Thus,

$$H^{(4)}(\zeta_4) = G^{(4)}(\zeta_4) - 120\zeta_4 G(1) = 0$$

Now

$$G^{(4)}(t) = -\frac{1}{3} (F'''(t) - F'''(-t)) - \frac{t}{3} (F^{(4)}(t) + F^{(4)}(-t))$$

and therefore

$$G(1) = -\frac{1}{360\zeta_4} (F'''(\zeta_4) - F'''(-\zeta_4)) - \frac{1}{360} (F^{(4)}(\zeta_4) + F^{(4)}(-\zeta_4))$$

Applying Lagrange's Mean Value Theorem to the first term on the right, we deduce that

$$G(1) = -\frac{1}{180} F^{(4)}(\zeta_5) - \frac{1}{360} (F^{(4)}(\zeta_4) + F^{(4)}(-\zeta_4))$$

for some  $\zeta_5 \in (-\zeta_4, \zeta_4)$ .

We shall apply Rolle's Theorem four times to the function  $H$ . Noting that  $H(0) = H(1) = 0$ , we deduce that there exists  $\zeta_1 \in (0, 1)$  such that  $H'(\zeta_1) = 0$ . However, also,  $H'(0) = 0$ , so there exists  $\zeta_2 \in (0, \zeta_1)$  such that  $H''(\zeta_2) = 0$ ; but,  $H''(0) = 0$ , so there is  $\zeta_3 \in (0, \zeta_2)$  such that  $H'''(\zeta_3) = 0$ . Finally, noting that  $H'''(0) = 0$ , it follows that there is  $\zeta_4 \in (0, \zeta_3)$  such that  $H^{(4)}(\zeta_4) = 0$ . Thus,

$$H^{(4)}(\zeta_4) = G^{(4)}(\zeta_4) - 120\zeta_4 G(1) = 0$$

Now

$$G^{(4)}(t) = -\frac{1}{3} (F'''(t) - F'''(-t)) - \frac{t}{3} (F^{(4)}(t) + F^{(4)}(-t))$$

and therefore

$$G(1) = -\frac{1}{360\zeta_4} (F'''(\zeta_4) - F'''(-\zeta_4)) - \frac{1}{360} (F^{(4)}(\zeta_4) + F^{(4)}(-\zeta_4))$$

Applying Lagrange's Mean Value Theorem to the first term on the right, we deduce that

$$G(1) = -\frac{1}{180} F^{(4)}(\zeta_5) - \frac{1}{360} (F^{(4)}(\zeta_4) + F^{(4)}(-\zeta_4))$$

for some  $\zeta_5 \in (-\zeta_4, \zeta_4)$ . Equivalently,

Since, by our hypothesis on  $f$ ,  $F^{(4)}$  is a continuous function on  $[-1, 1]$  and  $-1 < -\zeta_4 < \zeta_5 < \zeta_4 < 1$ , it follows that there exists  $\theta$  in  $[-\zeta_4, \zeta_4]$  such that

$$\frac{F^{(4)}(-\zeta_4) + 2F^{(4)}(\zeta_5) + F^{(4)}(\zeta_4)}{4} = F^{(4)}(\theta)$$

consequently,

$$G(1) = -\frac{1}{90}F^{(4)}(\theta) = -\frac{1}{1440}(b-a)^4 f^{(4)}(\xi) \quad (10)$$

where  $\xi = \frac{1}{2}(a+b) + \frac{1}{2}(b-a)\theta$ .

Since, by our hypothesis on  $f$ ,  $F^{(4)}$  is a continuous function on  $[-1, 1]$  and  $-1 < -\zeta_4 < \zeta_5 < \zeta_4 < 1$ , it follows that there exists  $\theta$  in  $[-\zeta_4, \zeta_4]$  such that

$$\frac{F^{(4)}(-\zeta_4) + 2F^{(4)}(\zeta_5) + F^{(4)}(\zeta_4)}{4} = F^{(4)}(\theta)$$

consequently,

$$G(1) = -\frac{1}{90}F^{(4)}(\theta) = -\frac{1}{1440}(b-a)^4 f^{(4)}(\xi) \quad (10)$$

where  $\xi = \frac{1}{2}(a+b) + \frac{1}{2}(b-a)\theta$ . Finally, (7) is obtained by inserting (10) into (9).



Since, by our hypothesis on  $f$ ,  $F^{(4)}$  is a continuous function on  $[-1, 1]$  and  $-1 < -\zeta_4 < \zeta_5 < \zeta_4 < 1$ , it follows that there exists  $\theta$  in  $[-\zeta_4, \zeta_4]$  such that

$$\frac{F^{(4)}(-\zeta_4) + 2F^{(4)}(\zeta_5) + F^{(4)}(\zeta_4)}{4} = F^{(4)}(\theta)$$

consequently,

$$G(1) = -\frac{1}{90}F^{(4)}(\theta) = -\frac{1}{1440}(b-a)^4 f^{(4)}(\xi) \quad (10)$$

where  $\xi = \frac{1}{2}(a+b) + \frac{1}{2}(b-a)\theta$ . Finally, (7) is obtained by inserting (10) into (9).

Now Lemma 1 yields the following bound on the error in Simpson's rule:

$$|E_2(f)| \leq \frac{1}{2880}(b-a)^5 M_4 \quad (11)$$

Since, by our hypothesis on  $f$ ,  $F^{(4)}$  is a continuous function on  $[-1, 1]$  and  $-1 < -\zeta_4 < \zeta_5 < \zeta_4 < 1$ , it follows that there exists  $\theta$  in  $[-\zeta_4, \zeta_4]$  such that

$$\frac{F^{(4)}(-\zeta_4) + 2F^{(4)}(\zeta_5) + F^{(4)}(\zeta_4)}{4} = F^{(4)}(\theta)$$

consequently,

$$G(1) = -\frac{1}{90}F^{(4)}(\theta) = -\frac{1}{1440}(b-a)^4 f^{(4)}(\xi) \quad (10)$$

where  $\xi = \frac{1}{2}(a+b) + \frac{1}{2}(b-a)\theta$ . Finally, (7) is obtained by inserting (10) into (9).

Now Lemma 1 yields the following bound on the error in Simpson's rule:

$$|E_2(f)| \leq \frac{1}{2880}(b-a)^5 M_4 \quad (11)$$

This is a considerable improvement over (5) in the sense that if  $f \in \mathcal{P}_3$  then the right-hand side of (11) is equal to zero and thereby  $E_2(f) = 0$  which now correctly reflects the fact that polynomials of degree three are integrated by Simpson's rule

$$f(b) - f(a) = f'(\zeta)(b - a).$$

without error. (As remarked earlier, this property was not borne out by our initial crude bound (5).)

$$f(b) - f(a) = f'(\zeta)(b - a).$$

without error. (As remarked earlier, this property was not borne out by our initial crude bound (5).)

By considering the right-hand side of the error bound in Theorem 1 we may be led to believe that by increasing  $n$ , that is by approximating the integrand by Lagrange polynomials of increasing degree and integrating these exactly, we shall be reducing the size of the quadrature error  $E_n(f)$ .

$$f(b) - f(a) = f'(\zeta)(b - a).$$

without error. (As remarked earlier, this property was not borne out by our initial crude bound (5).)

By considering the right-hand side of the error bound in Theorem 1 we may be led to believe that by increasing  $n$ , that is by approximating the integrand by Lagrange polynomials of increasing degree and integrating these exactly, we shall be reducing the size of the quadrature error  $E_n(f)$ . It is easy to find examples which show that this is not always the case.

$$f(b) - f(a) = f'(\zeta)(b - a).$$

without error. (As remarked earlier, this property was not borne out by our initial crude bound (5).)

By considering the right-hand side of the error bound in Theorem 1 we may be led to believe that by increasing  $n$ , that is by approximating the integrand by Lagrange polynomials of increasing degree and integrating these exactly, we shall be reducing the size of the quadrature error  $E_n(f)$ . It is easy to find examples which show that this is not always the case. A better approach is to divide the interval  $[a, b]$  into an increasing number of subintervals (of decreasing size) and use a numerical integration formula of fixed order on each of the subintervals.

$$f(b) - f(a) = f'(\zeta)(b - a).$$

without error. (As remarked earlier, this property was not borne out by our initial crude bound (5).)

By considering the right-hand side of the error bound in Theorem 1 we may be led to believe that by increasing  $n$ , that is by approximating the integrand by Lagrange polynomials of increasing degree and integrating these exactly, we shall be reducing the size of the quadrature error  $E_n(f)$ . It is easy to find examples which show that this is not always the case. A better approach is to divide the interval  $[a, b]$  into an increasing number of subintervals (of decreasing size) and use a numerical integration formula of fixed order on each of the subintervals. Quadrature rules based on this approach are called composite formulae, and will be described in detail in the next section.

# Composite trapezium rule

- . We shall consider only some very simple composite quadrature rules: the composite trapezium rule, and the composite Simpson rule.



# Composite trapezium rule

. We shall consider only some very simple composite quadrature rules: the composite trapezium rule, and the composite Simpson rule.

This is obtained by dividing the interval  $[a, b]$  into  $m$  equal subintervals, each of width  $h = (b - a)/m$ , so that

$$\int_a^b f(x) dx = \sum_{i=1}^m \int_{x_{i-1}}^{x_i} f(x) dx$$

where

$$x_i = a + ih = a + \frac{i}{m}(b - a), \quad i = 0, \dots, m$$

# Composite trapezium rule

. We shall consider only some very simple composite quadrature rules: the composite trapezium rule, and the composite Simpson rule.

This is obtained by dividing the interval  $[a, b]$  into  $m$  equal subintervals, each of width  $h = (b - a)/m$ , so that

$$\int_a^b f(x) dx = \sum_{i=1}^m \int_{x_{i-1}}^{x_i} f(x) dx$$

where

$$x_i = a + ih = a + \frac{i}{m}(b - a), \quad i = 0, \dots, m$$

Each of the integrals is then evaluated by the trapezium rule, namely,

$$\int_{x_{i-1}}^{x_i} f(x) dx \approx \frac{1}{2}h[f(x_{i-1}) + f(x_i)]$$

giving the complete approximation

$$\int_a^b f(x) dx \approx h \left[ \frac{1}{2}f(x_0) + f(x_1) + \dots + f(x_{m-1}) + \frac{1}{2}f(x_m) \right] \quad (12)$$

# Error in composite rules

The error in the composite trapezium rule can be estimated by using the error bound (??) for the trapezium rule on each individual subinterval  $[x_{i-1}, x_i], i = 1, \dots, m$ .

# Error in composite rules

The error in the composite trapezium rule can be estimated by using the error bound (??) for the trapezium rule on each individual subinterval  $[x_{i-1}, x_i]$ ,  $i = 1, \dots, m$ . Indeed,

$$\begin{aligned}\mathcal{E}_1(f) &:= \int_a^b f(x) dx - h \left[ \frac{1}{2} f(x_0) + f(x_1) + \dots + f(x_{m-1}) + \frac{1}{2} f(x_m) \right] \\ &= \sum_{i=1}^m \left[ \int_{x_{i-1}}^{x_i} f(x) dx - \frac{1}{2} h [f(x_{i-1}) + f(x_i)] \right]\end{aligned}$$

Applying (??) to each of the terms under the summation sign,

$$|\mathcal{E}_1(f)| \leq \frac{1}{12} h^3 \sum_{i=1}^m \left( \max_{\zeta \in [x_{i-1}, x_i]} |f''(\zeta)| \right) \quad (13)$$

$$\leq \frac{1}{12m^2} (b-a)^3 M_2, \quad (14)$$

where  $M_2 = \max_{\zeta \in [a,b]} |f''(\zeta)|$ .

# Composite Simpson's rule.

Let us suppose that the interval  $[a, b]$  has been divided into  $2m$  subintervals by the points  $x_i = a + ih, i = 0, \dots, 2m$ , where  $h = (b - a)/(2m)$ , and let us apply Simpson's rule on each of the intervals  $[x_{2i-2}, x_{2i}]$ ,  $i = 1, \dots, m$ , giving

$$\begin{aligned}\int_a^b f(x) dx &= \sum_{i=1}^m \int_{x_{2i-2}}^{x_{2i}} f(x) dx \\ &\approx \sum_{i=1}^m \frac{2h}{6} [f(x_{2i-2}) + 4f(x_{2i-1}) + f(x_{2i})]\end{aligned}$$

# Composite Simpson's rule.

Let us suppose that the interval  $[a, b]$  has been divided into  $2m$  subintervals by the points  $x_i = a + ih, i = 0, \dots, 2m$ , where  $h = (b - a)/(2m)$ , and let us apply Simpson's rule on each of the intervals  $[x_{2i-2}, x_{2i}]$ ,  $i = 1, \dots, m$ , giving

$$\begin{aligned}\int_a^b f(x) dx &= \sum_{i=1}^m \int_{x_{2i-2}}^{x_{2i}} f(x) dx \\ &\approx \sum_{i=1}^m \frac{2h}{6} [f(x_{2i-2}) + 4f(x_{2i-1}) + f(x_{2i})]\end{aligned}$$

Equivalently,

$$\begin{aligned}\int_a^b f(x) dx &\approx \frac{h}{3} [f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + \dots \\ &\quad + 2f(x_{2m-2}) + 4f(x_{2m-1}) + f(x_{2m})]\end{aligned}$$

# Composite Simpson's rule.

Let us suppose that the interval  $[a, b]$  has been divided into  $2m$  subintervals by the points  $x_i = a + ih, i = 0, \dots, 2m$ , where  $h = (b - a)/(2m)$ , and let us apply Simpson's rule on each of the intervals  $[x_{2i-2}, x_{2i}]$ ,  $i = 1, \dots, m$ , giving

$$\begin{aligned}\int_a^b f(x) dx &= \sum_{i=1}^m \int_{x_{2i-2}}^{x_{2i}} f(x) dx \\ &\approx \sum_{i=1}^m \frac{2h}{6} [f(x_{2i-2}) + 4f(x_{2i-1}) + f(x_{2i})]\end{aligned}$$

Equivalently,

$$\begin{aligned}\int_a^b f(x) dx &\approx \frac{h}{3} [f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + \dots \\ &\quad + 2f(x_{2m-2}) + 4f(x_{2m-1}) + f(x_{2m})]\end{aligned}$$

The numerical integration formula (??) is called **the composite Simpson rule**.

In order to estimate the error in the composite Simpson rule, we proceed in the same way as for the composite trapezium rule:

$$\begin{aligned}\mathcal{E}_2(f) &:= \int_a^b f(x) dx - \sum_{i=1}^m \frac{h}{3} [f(x_{2i-2}) + 4f(x_{2i-1}) + f(x_{2i})] \\ &= \sum_{i=1}^m \left[ \int_{x_{2i-2}}^{x_{2i}} f(x) dx - \frac{h}{3} [f(x_{2i-2}) + 4f(x_{2i-1}) + f(x_{2i})] \right]\end{aligned}$$

Applying (11) to each individual term under the summation sign and recalling that  $b - a = 2mh$ , we obtain the following error bound:

$$|\mathcal{E}_2(f)| \leq \frac{1}{2880m^4} (b - a)^5 M_4. \quad (15)$$

The composite rules (12) and (??) provide greater accuracy than the basic formulae considered in previous section; this is clearly seen by comparing the error bounds (14) and (15) for the two composite rules with (??) and (11), the error estimates for the basic trapezium and Simpson formula, respectively.



In order to estimate the error in the composite Simpson rule, we proceed in the same way as for the composite trapezium rule:

$$\begin{aligned}\mathcal{E}_2(f) &:= \int_a^b f(x) dx - \sum_{i=1}^m \frac{h}{3} [f(x_{2i-2}) + 4f(x_{2i-1}) + f(x_{2i})] \\ &= \sum_{i=1}^m \left[ \int_{x_{2i-2}}^{x_{2i}} f(x) dx - \frac{h}{3} [f(x_{2i-2}) + 4f(x_{2i-1}) + f(x_{2i})] \right]\end{aligned}$$

Applying (11) to each individual term under the summation sign and recalling that  $b - a = 2mh$ , we obtain the following error bound:

$$|\mathcal{E}_2(f)| \leq \frac{1}{2880m^4} (b - a)^5 M_4. \quad (15)$$

The composite rules (12) and (??) provide greater accuracy than the basic formulae considered in previous section; this is clearly seen by comparing the error bounds (14) and (15) for the two composite rules with (??) and (11), the error estimates for the basic trapezium and Simpson formula, respectively. The inequalities (14) and (15) indicate that, as long as the function  $f$  is sufficiently smooth, the errors in the composite rules can be made arbitrarily small by choosing a sufficiently large number of