

SEIRS Malaria Model with Seasonal Variations

ALOKEME Israël

2025-07-03

Introduction

This simulation models the spread of malaria using a SEIRS model that includes **seasonal variation** in transmission. Parameters are based on estimated data and some literature from **Benin and West Africa**.

Full R Code

```
# Load necessary packages
library(deSolve)
library(tidyverse)
library(reshape2)

# Define the SEIRS model with seasonality
SEIRS_model <- function(t, state, parameters) {
  with(as.list(c(state, parameters)), {

    P <- S + E + I + R
    beta_t <- beta * (1 + A * sin(2 * pi * t / T))
    lam <- beta_t * E / P

    dS <- -lam * S + mu * R
    dE <- lam * S - alpha * E
    dI <- alpha * E - epsilon * I
    dR <- epsilon * I - mu * R

    list(c(dS, dE, dI, dR))
  })
}

# Initial population and states
initP <- 1000000
initE <- 1
initI <- 1
initR <- 0
initS <- initP - initE - initI - initR
istate <- c(S = initS, E = initE, I = initI, R = initR)
```

```

# Parameters based on Benin context
parameters <- c(
  mu = 1/365,      # loss of immunity ~ 1 year
  beta = 0.16,     # transmission coefficient for R0 ~ 5
  A = 0.7,         # seasonal amplitude
  alpha = 1/12,    # incubation ~12 days
  T = 365,         # seasonal cycle (1 year)
  epsilon = 1/30   # infectious period ~30 days
)

# Time vector
t <- seq(0, 365, by = 1)

# Solve the ODEs
out <- ode(y = istate, times = t, func = SEIRS_model, parms = parameters)
out_df <- as.data.frame(out)

# Melt the dataframe and plot
out_long <- melt(out_df, id.vars = "time")

ggplot(out_long, aes(x = time, y = value, color = variable)) +
  geom_line(size = 1) +
  labs(title = "SEIRS Malaria Model with Seasonality", subtitle = "Benin
context",
  y = "Population count",
  x = "Time (days)") +
  theme_minimal()

```

SEIRS Malaria Model with Seasonality

Benin context

