

SECTION 4: ROUNDOFF AND TRUNCATION ERRORS

MAE 4020/5020 – Numerical Methods with MATLAB

2

Definitions of Error

True Error

3

- ***Absolute error*** – the difference between an approximation and the true value

$$E_t = (\text{approx. value}) - (\text{true value}) = \hat{x} - x$$

- ***Relative error*** – the true error as a percentage of the true value

$$\varepsilon_t = \frac{\hat{x} - x}{x} \cdot 100\%$$

- Both definitions ***require knowledge of the true value!***
 - ▣ If we had that, why would we be approximating?

Approximating the Error

4

- Since we don't know the true value, we can only approximate the error
- Often, approximations are made *iteratively*
 - ▣ Approximate the error as the change in the approximate value from one iteration to the next

$$\hat{E} = \hat{x}_{i+1} - \hat{x}_i$$

Relative Approximate Error

5

- We don't know the true value, so we can't calculate the true error – approximate the error
- ***Relative approximate error*** – an approximation of the error relative to the approximation itself

$$\varepsilon_a = \frac{\text{approx. error}}{\text{approximation}} \cdot 100\% = \frac{\hat{E}}{\hat{x}} \cdot 100\%$$

Stopping Criterion

6

- For iterative approximations, continue to iterate until the relative approximate error magnitude is less than a specified ***stopping criterion***

$$|\varepsilon_a| < \varepsilon_s$$

- For accuracy to ***at least n significant figures*** set the stopping criterion to

$$\varepsilon_s = (0.5 \times 10^{2-n})\%$$

7

Roundoff Error

Roundoff Errors


8

- **Roundoff errors** occur due to the way in which computers represent numerical values
- Computer representation of numerical values is limited in terms of:
 - ▣ **Magnitude** – there are upper and lower bounds on the magnitude of numbers that can be represented
 - ▣ **Precision** – not all numbers can be represented exactly
- Certain types of mathematical manipulations are more susceptible to roundoff error than others

Number Systems – Decimal

9

- We are accustomed to the ***decimal*** number system
 - ▣ A ***base-10*** number system
 - ▣ Ten digits: 0, 1, 2, 3, 4, 5, 6, 7, 8, 9
 - ▣ Each digit represents an integer power of 10



Binary Number System

10

- Computers represent numbers in **binary** format
 - ▣ A **base-2** number system
 - ▣ Two digits: 0, 1
 - Easy to store binary values in computer hardware – an on or off switch – a high or low voltage
 - ▣ One digit is a **bit** – eight bits is a **byte**
 - ▣ Each bit represents an integer power of 2

10110101


1	0	1	1	0	1	0	1
→	→	→	→	→	→	→	→
							$1 \times 2^0 = 1$
							$0 \times 2^1 = 0$
							$1 \times 2^2 = 4$
							$0 \times 2^3 = 0$
							$1 \times 2^4 = 16$
							$1 \times 2^5 = 32$
							$0 \times 2^6 = 0$
							$1 \times 2^7 = 128$

$= (181)_{10}$

IEEE Double-Precision Format

11

- By default, MATLAB uses double-precision floating point to store numeric values - `double`
 - ▣ 64-bit binary word



$$\pm \left(1 + \sum_{i=1}^{52} f_i \cdot 2^{-i} \right) \times 2^e$$

IEEE Double-Precision Format

12

□ **Mantissa**

- Only the ***fractional portion*** of the mantissa stored
- Bit to the left of the binary point assumed to be 1
 - ***Normalized numbers***
 - Really a 53-bit value


□ **Exponent**

- 11-bit signed integer value: -1022 ... 1023
- Two special cases:
 - $e = 0x000$ (i.e. ***all zeros***): zero if $f = 0$, ***subnormal #'s*** if $f \neq 0$
 - $e = 0x7FF$ (i.e. ***all ones***): ∞ if $f = 0$, NaN if $f \neq 0$

Normalized numbers

13

- Leading zeros are removed
 - ▣ Most significant digit (must be a 1 in binary) moved to the left of the binary point
 - ▣ 53rd bit of the mantissa (always 1) needn't be stored
- Maximum mantissa



Subnormal Numbers

14

- If $f_{52} = 1$, always, then the smallest number that could be represented is: $2^{-1022} \approx 2.225 \times 10^{-308}$
- If we allow for $f_{52} = 0$, then the most significant bit is somewhere to the right of the binary point
 - ▣ Leading zeros – not normalized ... ***subnormal***
 - ▣ Allows for smaller numbers, filling in the hole around zero
- Subnormal numbers represented by setting the exponent to zero
- Smallest subnormal number:

$$2^{-1022-52} = 2^{-1074} \approx 5 \times 10^{-324}$$

Doubles – Range

15

□ Maximum value:

$$max = \left(1 + \sum_{i=1}^{52} 2^{-i}\right) \times 2^{1023} \approx \mathbf{1.798 \times 10^{308}}$$

□ Minimum normal value:

$$min_{norm} = 2^{-1022} \approx \mathbf{2.225 \times 10^{-308}}$$

□ Minimum subnormal value:

$$min_{sub} = 2^{-1022-52} = 2^{-1074} \approx \mathbf{5 \times 10^{-324}}$$

□ Precision – *machine epsilon*

$$\epsilon = 2^{-52} \approx \mathbf{2.22 \times 10^{-16}}$$

Roundoff Error – Mathematical Operations

16

Certain types of mathematical operations are more susceptible to roundoff errors:

- ***Subtractive cancellation*** – subtracting of two nearly-equal numbers results in a loss of significant digits
- ***Large computations*** – even if the roundoff error from a single operation is small, the cumulative error from many operations may be significant
- ***Adding large and small numbers*** – as in an infinite series
- ***Inner products*** – (i.e. dot product) very common operation – solution of linear systems of equations

17

Truncation Error

Truncation Errors

18

- Errors that result from ***the use of an approximation*** in place of an exact mathematical procedure
 - ▣ E.g. numerical integration, or the approximation derivatives with finite-difference approximations

- To understand how truncation errors arise, and to gain an understanding of their magnitudes, we'll make use of the ***Taylor Series***

Taylor Series

19

- **Taylor's Theorem** – any smooth (i.e., continuously differentiable) function can be approximated as a polynomial
- **Taylor Series**

$$f(x_{i+1}) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_i)}{n!} (x_{i+1} - x_i)^n$$

- This infinite series is an **equality**
 - ▣ An **exact** representation of any smooth function as a polynomial
 - ▣ An **infinite-order polynomial** – impractical


Taylor Series Approximation

20

- Can approximate a function as a polynomial by **truncating** the Taylor series after a finite number of terms

$$f(x_{i+1}) \approx f(x_i) + f'(x_i)h + \frac{f''(x_i)}{2!}h^2 + \dots + \frac{f^{(n)}(x_i)}{n!}h^n$$

where $h = x_{i+1} - x_i$ is the **step size**



Taylor Series Truncation Error

21

- Can account for error by lumping the $n + 1$ and higher-order terms into a single term, R_n

$$f(x_{i+1}) = f(x_i) + f'(x_i)h + \frac{f''(x_i)}{2!}h^2 + \cdots + \frac{f^{(n)}(x_i)}{n!}h^n + R_n$$

- R_n is the error associated with truncating after n terms


$$R_n = \frac{f^{(n+1)}(\xi)}{(n+1)!}h^{n+1}$$

- ξ is some (**unknown**) value of x between x_i and x_{i+1}

Derivative Mean-Value Theorem

22

- If $f(x)$ and $f'(x)$ are continuous on $[x_i, x_{i+1}]$, then there is a point on this interval, ξ , where $f'(\xi)$ is the slope of the line joining $f(x_i)$ and $f(x_{i+1})$



Truncation Error – Dependence on Step Size

23

$$R_n = \frac{f^{(n+1)}(\xi)}{(n+1)!} h^{n+1}$$

- We don't know ξ , so we don't know R_n
 - ▣ We do know it's proportional to h^{n+1} , where h is the **step size**
 - ▣ Error is **on the order of** h^{n+1}

$$R_n = O(h^{n+1})$$

- If $n = 1$ (first-order approx.), **halving** the step size will **quarter** the error

Truncation Errors in Practice

- Discretizing equations
- Finite-difference approximations

Discretization of Equations

25

- As engineers, many of the mathematical expressions we are interested in are differential equations
 - ▣ We know how to evaluate derivatives analytically
 - ▣ Need an approximation for the derivative operation in order to solve numerically
- **Discretization** – conversion of a continuous function, e.g. differentiation, to a discrete approximation for numerical evaluation

Finite Difference Approximations

26

- Recall the definition of a derivative

$$f'(x_i) = \lim_{h \rightarrow 0} \frac{f(x_{i+1}) - f(x_i)}{h}$$

- Remove the limit to approximate this numerically

$$f'(x_i) \approx \frac{f(x_{i+1}) - f(x_i)}{h}$$

- This is the ***forward difference approximation***
 - ▣ Uses value at x_i and forward one step at x_{i+1} to approximate the derivative at x_i

Discretizing Equations – Example

27

- A free-falling object (bungee jumper example from the text) can be modeled as

$$\frac{dv}{dt} = g - \frac{c_d}{m} v^2$$

where v is velocity, m is mass, g is gravitational acceleration, and c_d is a lumped drag coefficient

- This is a non-linear ordinary differential equation (ODE), which can be solved analytically to yield

$$v(t) = \sqrt{\frac{mg}{c_d}} \tanh \left(\sqrt{\frac{gc_d}{m}} \cdot t \right)$$

Discretizing Equations – Example

28

- To solve numerically instead, approximate the derivative operation with a finite difference

$$\frac{v(t_{i+1}) - v(t_i)}{t_{i+1} - t_i} \cong g - \frac{c_d}{m} v(t_i)^2$$

- Solving for $v(t_{i+1})$ and using h to denote the time step yields

$$v(t_{i+1}) \cong v(t_i) + \left[g - \frac{c_d}{m} v(t_i)^2 \right] h$$

- We've transformed the ***differential equation*** to a ***difference equation***
 - ▣ An ***algebraic*** equation
 - ▣ Can be solved iteratively – using a loop

Discretizing Equations – Example

29

$$v(t_{i+1}) \cong v(t_i) + \left[g - \frac{c_d}{m} v(t_i)^2 \right] h$$

- The term in the square brackets is the original diff. eq., i.e. it is $v'(t)$
- The difference equation is a ***first-order Taylor series approximation***

$$v(t_{i+1}) = v(t_i) + v'(t_i) h + R_1$$

- Where we know that the error is on the order of the step size squared

$$R_1 = O(h^2)$$

- ***Taylor series provides a relation between the step size and the accuracy of the numerical solution to the diff. eqn.***

Finite Difference Methods

30

- The preceding example showed
 - ▣ One method – forward difference – for numerically approximating a derivative
 - ▣ Transformation of a differential equation to a difference equation
 - ▣ How Taylor series can provide an understanding of the error associated with an approximation
- Now we'll take a closer look at three ***finite difference methods*** and how Taylor series can help us understand the error associated with each

Forward Difference

31

- Can also derive the forward difference approximation from the Taylor Series

$$f(x_{i+1}) = f(x_i) + f'(x_i)h + R_1$$

- Solving for $f'(x_i)$

$$f'(x_i) = \frac{f(x_{i+1}) - f(x_i)}{h} - \frac{R_1}{h}$$

- We've already seen that

$$R_1 = O(h^2)$$

- So, the error term is

$$\frac{R_1}{h} = O(h)$$


- The forward difference, including error, is

$$f'(x_i) = \frac{f(x_{i+1}) - f(x_i)}{h} + O(h)$$

- ***Error of the forward difference approximation is on the order of the step size***

Forward Difference

32



- Value of the function, $f(x)$, at x_i and forward one step at x_{i+1} used to approximate the derivative at x_i

Backward Difference

33

- **Backward difference** uses value of $f(x)$ at x_i and **one step backward** at x_{i-1} to approximate the derivative at x_i

$$f'(x_i) \approx \frac{f(x_i) - f(x_{i-1})}{h}$$

- This can also be developed by expanding the Taylor series backward

$$f(x_{i-1}) = f(x_i) - f'(x_i)h + R_1$$

- Then solving for $f'(x_i)$

$$f'(x_i) = \frac{f(x_i) - f(x_{i-1})}{h} + \frac{R_1}{h}$$

- Again the error is on the order of the step size

$$\frac{R_1}{h} = O(h)$$


- The backward difference expression, including error, becomes

$$f'(x_i) = \frac{f(x_i) - f(x_{i-1})}{h} + O(h)$$

- **Error of the backward difference approximation is on the order of the step size**

Backward Difference

34



- Now use the value of $f(x)$ at x_i and backward one step at x_{i-1} to approximate the derivative at x_i

- Again, error is $R = O(h)$

Central Difference

35

- **Central difference** uses value of $f(x)$ **one step backward** at x_{i-1} and one step ahead at x_{i+1} to approximate the derivative at x_i

$$f'(x_i) \approx \frac{f(x_{i+1}) - f(x_{i-1}))}{2h}$$

- This can also be developed by subtracting the backward Taylor series from the forward series
- Second-order derivative terms cancel, leaving

$$f(x_{i+1}) = f(x_{i-1}) + 2f'(x_i)h + R_2$$

- Now, the remainder term is

$$R_2 = O(h^3)$$


- The central difference expression, including error, becomes

$$f'(x_i) = \frac{f(x_{i+1}) - f(x_{i-1}))}{2h} + O(h^2)$$

- **Error of the central difference approximation is on the order of the step size squared**
- Central difference method is more accurate than forward or backward
 - ▣ Uses more information

Central Difference

36



- Now use the value of $f(x)$ backward one step at x_{i-1} and forward one step at x_{i+1} to approximate the derivative at x_i

- **Reduced error:**

$$R = O(h^2)$$

37

Total Numerical Error

Total Numerical Error


38

- Total numerical error is the ***sum of roundoff and truncation error***
 - ***Roundoff error*** is largely out of your control, and, with double precision arithmetic, it is not typically an issue
 - ***Truncation error*** can be a significant problem, but can be reduced by decreasing step size
- Reducing step size reduces truncation error, but may also result in subtractive cancellation, thereby increasing roundoff error
- Choose step size to minimize total error
 - Or, more typically, to reduce truncation error to an acceptable level

Total Numerical Error

39

- Reducing step size reduces truncation error, but may also result in subtractive cancellation, thereby increasing roundoff error
- Could choose step size to minimize total error
- But, more typically, reduce step size just enough to reduce truncation error to an acceptable level



Central Difference Error Analysis

40

- First derivative of a function in terms of the central difference approximation is

$$f'(x_i) = \frac{f(x_{i+1}) - f(x_{i-1}))}{2h} - \frac{f^{(3)}(\xi)}{6} h^2$$

- The last term on the right is the truncation error
- There is also roundoff error associated with each value

$$f(x_{i-1}) = \tilde{f}(x_{i-1}) - e_{i-1}$$

$$f(x_{i+1}) = \tilde{f}(x_{i+1}) - e_{i+1}$$

where $\tilde{f}(x_i)$ represents a rounded value, and e_i is the corresponding roundoff error

Central Difference Error Analysis

41

- Substituting the expressions for the rounded values into the expression for the true derivative yields

$$f'(x_i) = \frac{f(x_{i+1}) - f(x_{i-1}))}{2h} - \frac{f^{(3)}(\xi)}{6} h^2 + \frac{e_{i+1} - e_{i-1}}{2h}$$

- Giving a total error of

Roundoff error \rightarrow $err = \frac{e_{i+1} - e_{i-1}}{2h} - \frac{f^{(3)}(\xi)}{6} h^2$ \leftarrow Truncation error

- ***Truncation error increases with step size***
- ***Roundoff error decreases with step size***