X

derivative

Examples

Random

Assuming "derivative" refers to a computation | Use as a general topic or referring to a mathematical definition or a word instead

■ function to differentiate: x^(x^2)

Also include: differentiation variable


Derivative:

$$\frac{d}{dx}\left(x^{x^2}\right) = x^{x^2} \left(x + 2x\log(x)\right)$$

Step-by-step solution

 $\log(x)$ is the natural logarithm


Plots:



Complex-valued plot ▼

Enable interactivity

Complex-valued plot ▼



Enable interactivity

Alternate form:

$$x^{x^2+1}\left(2\log(x)+1\right)$$

Expanded form:

$$x^{x^2+1} + 2x^{x^2+1}\log(x)$$

Root

Approximate form

Step-by-step solution

$$x = \frac{1}{\sqrt{e}}$$

Properties as a real function:

Exact forms

Domain:

 $\{x \in \mathbb{R} : x > 0\}$ (all positive real numbers)

Range:

 $\{y \in \mathbb{R}: y \geq -0.416128\}$

R is the set of real numbers

 $x \left(2 \log(x) + 1 \right) + x^3 \left(2 \log^2(x) + \log(x) \right) + O\left(x^4\right)$

(generalized Puiseux series)

Big-O notation »

Indefinite integral:

 $\int x^{x^2} (x + 2x \log(x)) dx = x^{x^2} + \text{constant}$

Step-by-step solution

Differential geometric curves:

(requires interactivity)

Enable Interactivity

POWERED BY THE WOLFRAM LANGUAGE

Standard computation time exceeded...

Try again with additional computation time »

Related Queries:

= osculating circle of x^(x^2)

= integrate x^(x^2) dx

= polar plot abs(theta^(theta^2))

= d^2/dx^2 x^(x^2)

About Pro Products Mobile Apps Business Solutions For Developers Resources & Tools

Blog Community Participate Contact Connect

© 2015 Wolfram Alpha LLC—A Wolfram Research Company Terms Privacy Policy