

empty

21-128 and 15-151

[3pt] Problem Sheet 1

[7pt]

Solutions to the following six exercises and optional bonus problem are to be submitted to Gradescope by

11 PM on Wednesday, the 2nd of September, 2026.

0.6em

Instructions. This assignment has 30 points, together with an optional 1-point bonus problem. Unless a problem explicitly says otherwise, every answer must be justified by a proof. A counterexample must satisfy all of the hypotheses and must be checked explicitly. You may cite results proved in lecture or recitation, provided that you identify the result you are using.

* Problem1[4points] Let $a, b \in \mathbb{Z}_{>0}$.

[(a)]

1. Suppose that $a \mid b$. Prove that

$$a \mid a^2 + b^2.$$

2. Is the converse true? In other words, if

$$a \mid a^2 + b^2,$$

must $a \mid b$? Prove your answer.

* Problem2[4points] Find, with proof, all real solutions of

$$5 = 2\sqrt{x} + \sqrt{10 - x}.$$

Your solution should identify the domain of the equation, explain whether each squaring step is reversible, and check every candidate in the original equation.

* Problem3[5points] Let $D(m, n)$ be the predicate “ m divides n ,” where all variables range over $\mathbb{Z}_{>0}$. Determine whether each proposition below is true or false.

$\begin{array}{l} \text{(a)} \text{ for all } m, \text{ exists } n; D(m, n), \\ \text{(b)} \text{ exists } n, \text{ for all } m; D(m, n), \\ \text{(c)} \text{ for all } m, n; D(m, n) \end{array}$

For every true proposition, give a valid witness or a rule for choosing a witness. For every false proposition, state and prove its negation.

* Problem4[5points]

[(a)]

1. Prove that

$$(p \rightarrow q) \vee (p \rightarrow r) \iff (p \rightarrow (q \wedge r))$$

are logically equivalent. You may use a truth table or standard logical equivalences, but every step must be clear.

2. Let P and Q be predicates on a set S . Prove that

$$\left(\bigwedge_{x \in S} P(x) \right) \vee \left(\bigwedge_{x \in S} Q(x) \right) \iff \bigwedge_{x \in S} (P(x) \vee Q(x))$$

Then give explicit choices of S , P , and Q showing that the converse implication is false.

* Problem5[7points] Consider the equation

$$x^4y + ay + x = 0,$$

where $a, x, y \in \mathbb{R}$.

[(a)]

1. Show that the proposition

$$(\forall a \in \mathbb{R})(\forall x \in \mathbb{R})(\exists! y \in \mathbb{R})(x^4y + ay + x = 0)$$

is false. Give one pair (a, x) for which no real solution y exists and another pair (a, x) for which more than one real solution y exists.

2. Determine, with proof, the set of all real numbers a for which

$$(\forall x \in \mathbb{R})(\exists! y \in \mathbb{R})(x^4y + ay + x = 0)$$

is true.

* Problem6[5points] Let $r \in \mathbb{Q} \setminus \{0\}$ and let $a, b \in \mathbb{R} \setminus \mathbb{Q}$. Determine which of the following expressions are irrational for *every* permitted choice of r, a, b :

$$a + r, \sqrt[4]{a}, \sqrt[4]{a} + b, \sqrt[4]{ab}.$$

For each expression that must be irrational, give a proof. For each expression that need not be irrational, give an explicit counterexample and verify that all numbers in the example satisfy the hypotheses.

*Bonus Problem [1 bonus point] Katie, Hasita, and Michelle are perfect logicians. Each has a positive integer written on their hat. The numbers are not necessarily distinct, and it is common knowledge that two of the three numbers add to the third. Each logician can see the other two hats but not their own.

The following conversation is public: **Katie:** I do not know my number.

Hasita: I do not know my number.

Michelle: I do not know my number.

Katie: Now I know my number. It is 50.

[(a)]

1. Determine the numbers on Hasita's and Michelle's hats, and justify every inference.
2. At Michelle's first turn—after they have heard Katie and Hasita each say “I do not know”—characterize all ordered pairs (u, v) that they could see on Katie's and Hasita's hats for which they would be able to determine their own number. Your condition should be both necessary and sufficient.