

NEUROPHARMACOLOGY LAB REPORT — CONTROLLED SESSION

COMPOUND DETAILS

Compound	Triaxion-47	Drug Class	SSRI
Dose (mg eq.)	20mg	BBB Coefficient	0.60
Receptor Targets	5-HT1A		
Session Date	2026-06-12T16:21:30.852Z		

Tests run in an isolated Controlled Lab session. Live brain was not modified. Platform biologically constrained by HCP-MMP1.0, Allen Brain Atlas, and Brainnetome Atlas data.

PEAK RECEPTOR OCCUPANCY

5-HT1A
2.8%

Occupancy curves follow rapid absorption / first-order elimination kinetics (high-level approximation). Not equivalent to full pharmacokinetic modeling.

REGIONAL ACTIVATION EFFECTS

Region	Receptor	Activation Delta	Predicted Effect (consistent with literature)
Amygdala	5-HT1A	-16.9%	reduced exploratory drive
PrefrontalCortex	5-HT1A	-16.8%	reduced exploratory drive
Hippocampus	5-HT1A	-16.5%	reduced exploratory drive

Effects represent simulated predictions from population-level receptor binding models. Not equivalent to in vivo pharmacology data.

VALIDATION SUMMARY

Integrity: PASS Stability: PASS Verdict: WEAK EFFECT

ANSInteroceptiveCoupling did not produce a measurable separation from baseline or shuffled control on pre-registered metrics. Claims of biological significance are not supported by this run.

Language note: This report uses "consistent with", "suggests", or "did not support" for all mechanistic claims. The simulation does not "prove" or "demonstrate" pharmacological outcomes.