RESEARCH PAPER · THEORETICAL PHYSICS

Negative-Mass Antimatter and the Baryon Asymmetry Problem

A CPT-Symmetric Cosmological Framework

Author John, CaesarBot & Iuppiter Group
Published April 2026 Status Preprint, not peer reviewed

Category Physics · Cosmology References 50+ References

ABSTRACT

The observed baryon asymmetry of the universe — the overwhelming dominance of matter over antimatter — remains one of the deepest unsolved problems in modern physics. The Standard Model, even supplemented with the Sakharov conditions for baryogenesis, fails to produce sufficient CP violation to account for the observed asymmetry ratio of $\eta B \approx 6.1 \times 10^{-10}$. This report explores an alternative hypothesis: that the apparent absence of antimatter is not the result of a dynamical asymmetry generated in the early universe, but rather a consequence of antimatter possessing negative gravitational mass and, through the Feynman–Stückelberg interpretation, propagating backward in the temporal direction. Under this framework, the universe at the Big Bang is exactly CPT-symmetric, with equal quantities of matter and antimatter created, but with antimatter gravitationally repelled from matter and occupying a CPT-conjugate temporal sector. We develop this hypothesis by synthesizing several lines of established and speculative physics: Bondi's (1957) negative mass framework in general relativity, the Feynman–Stückelberg interpretation of antiparticles, Villata's (2011)

CPT-based argument for antimatter–matter gravitational repulsion, the Boyle–Finn–Turok (2018) CPT-symmetric universe model, and Farnes' (2018) negative-mass cosmology. We examine the cosmological implications for Big Bang nucleosynthesis, CMB observations, dark energy, and large-scale structure formation. Experimental constraints from the ALPHA-g (2023), AEGIS, and GBAR experiments at CERN are critically evaluated. We address the principal counterarguments — including the runaway paradox, energy conservation concerns, and the 2023 ALPHA-g results showing gravitational attraction of antihydrogen — with intellectual honesty about where the hypothesis retains viability and where it faces serious challenges. We conclude that while the simplest version of the negative-mass antimatter hypothesis (antigravity for antihydrogen in our temporal sector) is now empirically disfavored by ALPHA-g, more sophisticated versions involving CPT-symmetric cosmological boundary conditions remain viable and experimentally distinguishable from standard cosmology.

KEYWORDS: baryon asymmetry, antimatter, negative mass, CPT symmetry, antigravity, dark energy, Feynman–Stückelberg interpretation, cosmological constant

A Speculative Theoretical Investigation

SECTION I

1. Introduction

1.1 The Matter–Antimatter Asymmetry Problem

One of the most profound puzzles in contemporary physics is deceptively simple to state: Why is there something rather than nothing? More precisely, why does the observable universe consist almost entirely of matter, with antimatter appearing only in trace quantities produced by high-energy processes such as cosmic ray interactions and radioactive decay?

The Standard Model of particle physics, combined with the hot Big Bang cosmological model, predicts that the early universe should have produced matter and antimatter in nearly equal quantities. At temperatures above the QCD confinement scale ($T \gtrsim 200$ MeV), quarks and antiquarks existed in thermal equilibrium, continuously created and annihilated in pairs. As the universe cooled, these pairs should have mutually annihilated, leaving behind a universe of pure radiation — photons and neutrinos — with essentially no residual baryonic matter.

Yet we observe precisely the opposite: a universe filled with galaxies, stars, planets, and observers, composed overwhelmingly of matter. The baryon-to-photon ratio, as measured by the cosmic microwave background (CMB) anisotropies by the Planck satellite, is (Planck Collaboration, 2020):

$$\eta_B = \frac{n_B - n_{\bar{B}}}{n_\gamma} = (6.143 \pm 0.019) \times 10^{-10}$$

This number, while small, represents an enormous asymmetry: for every billion antimatter particles produced in the Big Bang, there were approximately one billion and one matter particles. The billion pairs annihilated, and the lone excess matter particle survived to form everything we observe.

The absence of significant antimatter in the observable universe is confirmed by multiple independent lines of evidence:

1. Cosmic ray composition: The observed flux of antiprotons in cosmic rays is consistent with secondary production from high-energy collisions and shows no evidence of primary antimatter sources (AMS-02 Collaboration, Aguilar et al., 2016).

1. Gamma-ray background: If regions of antimatter existed adjacent to regions of matter, their boundaries would produce annihilation radiation at 511 keV and at the pion mass scale (~135 MeV). No such signatures have been observed at cosmological scales (Cohen, De Rújula, & Glashow, 1998).

of ηB (Fields, Olive, Yeh, & Young, 2020).

1. CMB anisotropies: The acoustic peak structure of the CMB power spectrum is sensitive to the baryon-to-photon ratio and yields values fully consistent with BBN (Planck Collaboration, 2020).

1.2 The Sakharov Conditions and Their Insufficiency

In a landmark 1967 paper, Andrei Sakharov identified three necessary conditions for the dynamical generation of a baryon asymmetry from an initially symmetric state (Sakharov, 1967):

1. Baryon number (B) violation: Interactions must exist that change the net baryon number.

1. C and CP violation: Charge conjugation symmetry and the combined charge-parity symmetry must be violated, so that processes creating baryons proceed at different rates than those creating antibaryons.

1. Departure from thermal equilibrium: The baryon-number-violating, CP-violating processes must occur out of thermal equilibrium, since CPT symmetry guarantees that in equilibrium, any asymmetry-generating process is exactly compensated by its CPT conjugate.

The Standard Model, remarkably, satisfies all three conditions in principle:

- B violation occurs through non-perturbative electroweak sphaleron processes, which violate B + L while conserving B - L ('t Hooft, 1976; Klinkhamer & Manton, 1984).

- CP violation is present in the CKM quark mixing matrix, as demonstrated experimentally in the kaon and B-meson systems (Christenson, Cronin, Fitch, & Turlay, 1964; BaBar Collaboration, 2001; Belle Collaboration, 2001).

- Departure from equilibrium could occur during the electroweak phase transition if it is first-order.

However, the Standard Model fails quantitatively on at least two fronts:

First, the amount of CP violation in the CKM matrix is insufficient by many orders of magnitude. The Jarlskog invariant $J \approx 3 \times 10^{-5}$, and when combined with the relevant mass scales, produces an asymmetry of order $\eta_{BSM} \sim 10^{-18}$, roughly eight orders of magnitude too small (Gavela, Hernández, Orloff, & Pène, 1994; Huet & Sather, 1995).

Second, lattice gauge theory calculations have demonstrated that the electroweak phase transition in the Standard Model (with the observed Higgs mass of $m_H \approx 125$ GeV) is a smooth crossover rather than a first-order phase transition, invalidating the third Sakharov condition within the Standard Model framework (Kajantie, Laine, Rummukainen, & Shaposhnikov, 1996).

These failures have motivated an enormous body of work on beyond-Standard-Model (BSM) baryogenesis mechanisms, including:

- GUT baryogenesis via the decay of superheavy bosons (Yoshimura, 1978; Weinberg, 1979)
- Electroweak baryogenesis with extended Higgs sectors providing a first-order phase transition and additional CP violation (Morrissey & Ramsey-Musolf, 2012)
- Leptogenesis via the out-of-equilibrium decay of heavy right-handed neutrinos, with subsequent sphaleron conversion of lepton asymmetry to baryon asymmetry (Fukugita & Yanagida, 1986)
- Affleck-Dine baryogenesis using flat directions in supersymmetric theories (Affleck & Dine, 1985)

Despite decades of effort, none of these mechanisms has been experimentally confirmed, and the baryon asymmetry problem remains open.

1.3 An Alternative Perspective: No Asymmetry At All?

This report explores a radically different approach. Rather than seeking a dynamical mechanism to generate a matter–antimatter asymmetry, we ask: What if no asymmetry was ever generated? What if the Big Bang produced exactly equal quantities of matter and antimatter, and the apparent absence of

antimatter is an observational artifact arising from a fundamental property of antimatter that we have not yet correctly identified?

Specifically, we investigate the hypothesis that:

1. Antimatter possesses negative gravitational mass, causing it to be gravitationally repelled by matter.

1. Under the Feynman–Stückelberg interpretation, antiparticles propagate backward in time.

1. At the Big Bang, the CPT symmetry of the laws of physics was exactly realized: equal matter and antimatter were created, but antimatter, having negative mass, was repelled into the opposite temporal direction, forming a CPT-mirror image of our universe.

1. The observed baryon "asymmetry" is therefore not an asymmetry of the universe as a whole, but merely a feature of our local temporal sector.

This hypothesis, while speculative, draws on legitimate threads in theoretical physics and has been partially developed by several authors, notably Villata (2011), Benoit-Lévy and Chardin (2012), Boyle, Finn, and Turok (2018), and Farnes (2018). It makes specific, falsifiable predictions that are being tested by ongoing experiments at CERN.

1.4 Scope and Organization

This report is structured as follows. Section 2 establishes the theoretical framework, reviewing the Feynman–Stückelberg interpretation of antiparticles, CPT symmetry, and the physics of negative mass in general relativity and quantum field theory. Section 3 develops the negative mass–antimatter hypothesis formally, connecting it to CPT-symmetric cosmological models. Section 4 explores the cosmological implications for nucleosynthesis, the CMB, dark energy, and structure formation. Section 5 critically evaluates the experimental evidence, particularly the recent ALPHA-g results from CERN. Section 6 examines the relationship between negative-mass antimatter and the dark sector (dark matter and dark energy). Section 7 presents the mathematical formalism in detail. Section 8 addresses criticisms and

SECTION II

2. Theoretical Framework

2.1 The Feynman–Stückelberg Interpretation

The idea that antiparticles can be interpreted as particles traveling backward in time originated independently with Ernst Stückelberg (1941, 1942) and Richard Feynman (1948, 1949). In Feynman's formulation of quantum electrodynamics, a positron is mathematically equivalent to an electron propagating backward in time:

$$\psi_{\text{positron}}(x, t) \sim \psi^*_{\text{electron}}(x, -t)$$

More precisely, in the Feynman propagator formalism, the negative-energy solutions of the Dirac equation that propagate backward in time are reinterpreted as positive-energy antiparticles propagating forward in time. This is not merely a mathematical trick; it is built into the structure of quantum field theory. The Feynman propagator for a scalar field is:

$$\Delta_F(x - y) = \int \frac{d^4p}{(2\pi)^4} \frac{i}{p^2 - m^2 + i\varepsilon}$$

The $i\varepsilon$ prescription ensures that positive-energy solutions propagate forward in time while negative-energy solutions propagate backward in time. Under charge conjugation, the backward-propagating negative-energy electron becomes a forward-propagating positive-energy positron.

In Feynman diagrams, this interpretation is manifest: an electron line with an arrow pointing backward in time is identified with a positron line pointing forward in time. This is a fundamental feature of relativistic quantum field theory and has been experimentally validated to extraordinary precision through the success of QED calculations.

The key question is whether this mathematical equivalence between "antiparticle going forward in time" and "particle going backward in time" has deeper physical significance, or whether it is merely a convenient computational device. The hypothesis under investigation takes the former view seriously.

2.2 CPT Symmetry

The CPT theorem, proved by Lüders (1954), Pauli (1955), and Jost (1957), states that any Lorentz-invariant local quantum field theory with a Hermitian Hamiltonian is invariant under the combined operation of:

- C (charge conjugation): replacing particles with antiparticles
- P (parity): spatial inversion $x \to -x$
- T (time reversal): $t \to -t$

Formally, for any quantum field theory satisfying the Wightman axioms:

$$\Theta \, L(x) \, \Theta^{-1} = L(-x)$$

where $\Theta = CPT$ is the combined CPT operator. This means that the CPT-transformed version of any physical process has exactly the same probability amplitude as the original process. Equivalently:

$$M(A \to B) = M(B_{\text{(mirror, time-reversed)}} \to A_{\text{(mirror, time-reversed)}})$$

CPT symmetry has been tested to extraordinary precision. The most stringent tests come from the neutral kaon system, where CPT-violating effects are constrained at the level of $|\delta_{(CPT)}| < 10^{-19}$ GeV (CPLEAR Collaboration, Angelopoulos et al., 1998), and from comparisons of the electron and positron magnetic moments, which agree to better than one part in 10^{12} (Hanneke, Fogwell, & Gabrielse, 2008).

The relevance of CPT to our hypothesis is profound. Under CPT:

- A particle with positive mass, positive charge, and forward time evolution maps to an antiparticle with (potentially) negative mass, negative charge, and backward time evolution.

The question is whether CPT merely relates the internal quantum numbers of particles and antiparticles (as is the standard interpretation) or whether it also relates their gravitational properties (as proposed by Villata and others).

2.3 Negative Mass in General Relativity

The concept of negative mass has a long and respectable history in general relativity. Hermann Bondi (1957) provided the first systematic analysis, distinguishing between three types of mass:

1. Inertial mass (m_i): the resistance to acceleration, appearing in $F = m_i a$.

1. Active gravitational mass (m_a): the source of gravitational field.

1. Passive gravitational mass (m_(p)): the response to an external gravitational field.

The equivalence principle, tested to extraordinary precision ($|m_i - m_(p)|/m_i < 10^{-15}$ by the MICROSCOPE satellite; Touboul et al., 2017), asserts that $m_i = m_(p)$. Conservation of momentum requires $m_a = m_(p)$. Thus, in standard physics, all three masses are equal.

Bondi showed that if all three masses are simultaneously negative (maintaining the equivalence principle), the dynamics are self-consistent but exotic. For two point masses $M > 0$ and $m < 0$ interacting gravitationally:

- The positive mass M attracts both positive and negative masses.

- The negative mass m repels both positive and negative masses.

- The negative mass accelerates *toward* the positive mass (because $F = m_i a$ with $m_i < 0$ reverses the direction of acceleration).

- The positive mass accelerates *away from* the negative mass.

This produces the famous runaway pair: the negative mass chases the positive mass, both accelerating indefinitely without violating conservation of energy or momentum (since the kinetic energy of the

Bonnor (1989) extended Bondi's analysis and examined the behavior of negative mass in various general-relativistic contexts, including the Schwarzschild solution and cosmological models. He showed that a negative-mass Schwarzschild solution is repulsive and produces a naked singularity rather than a black hole.

Forward (1990) explored potential technological applications of negative mass, including propulsion systems. While highly speculative, Forward's work helped maintain theoretical interest in the concept.

2.4 Negative Mass in Quantum Field Theory

In quantum field theory, the concept of negative mass is more constrained but not entirely excluded. The mass parameter m in the Lagrangian density:

$$L = \psi(i\gamma^\mu \partial_\mu - m)\psi$$

is conventionally taken to be positive. However, this is a convention related to the choice of vacuum state rather than a fundamental requirement. The physical mass is the pole of the propagator:

$$G(p) = (i(\gamma^\mu p_\mu + m))/(p^2 - m^2 + i\varepsilon)$$

and what is physically measurable is m^2, not the sign of m itself. In the standard interpretation, both signs of m yield the same physics for free fields.

The situation changes when gravity is included. In the semiclassical approximation, the source of gravity is the stress-energy tensor:

$$T\mu\nu = (2)/(\sqrt{-g}) \text{frac}\delta S_{(matter)}\delta g_{(\mu\nu)}$$

For a Dirac field, the energy density (which serves as the source of gravity in the Newtonian limit) is:

$$T^{oo} = \psi\gamma^o (i\gamma^i \partial_i + m)\psi$$

gravitational coupling, antimatter would gravitate differently than matter.

2.5 The Dirac Sea Reinterpretation

Dirac's original (1930) interpretation of the negative-energy solutions of his equation involved a "sea" of filled negative-energy states. A hole in this sea — an absence of a negative-energy, negative-charge electron — manifests as a positive-energy, positive-charge particle: the positron.

This picture, while superseded by the modern QFT formalism, offers an interesting perspective on the mass of antimatter. In the Dirac sea picture, a positron is the absence of a negative-energy electron. If we assign gravitational properties based on the energy of the state, the absence of negative energy is equivalent to the presence of positive energy — and the positron should gravitate normally.

However, this argument can be turned around. If we consider the negative-energy electron as having negative gravitational mass, then its absence (the positron) has the opposite gravitational mass — namely, positive gravitational mass in its rest frame. But this depends crucially on whether the "negative energy" of the Dirac sea has gravitational reality.

In the modern QFT reinterpretation, the vacuum is the state of minimum energy, and all particles (including antiparticles) have positive energy by construction. The gravitational properties of antiparticles are then determined by the positive-energy stress-energy tensor they carry. This standard argument predicts that antimatter falls down, not up.

The tension between the Dirac sea picture (where antiparticles are "holes" in a negative-energy sea) and the modern QFT picture (where antiparticles are positive-energy excitations) remains a source of conceptual ambiguity regarding the gravitational properties of antimatter — an ambiguity that can ultimately only be resolved experimentally.

2.6 Previous Theoretical Work on Antimatter Gravity

Several authors have argued, on various theoretical grounds, that antimatter might experience repulsive gravity.

Morrison (1958) noted that the experimental evidence for normal gravitational behavior of antimatter was, at the time, entirely indirect and called for direct tests.

Schiff (1958, 1959) argued that virtual antimatter loops in quantum corrections to the gravitational interaction of normal matter would lead to inconsistencies if antimatter responded differently to gravity. This argument was long considered decisive, but subsequent authors (notably Nieto & Goldman, 1991) pointed out loopholes in Schiff's reasoning.

Nieto and Goldman (1991) provided a comprehensive review of the theoretical arguments for and against antigravity for antimatter, concluding that the question was more open than commonly assumed.

Chardin and Rax (1992) proposed that CP violation in the kaon system could be explained as a gravitational effect if antimatter experienced repulsive gravity.

Villata (2011) presented an argument based on CPT symmetry and general relativity. He noted that under CPT transformation, the geodesic equation:

$$(d^2 x^\mu)/(d\tau^2) + \Gamma^\mu_{\alpha\beta}(dx^\alpha)/(d\tau)(dx^\beta)/(d\tau) = 0$$

transforms such that the Christoffel symbols, which depend on the metric and hence on the mass of the source, change sign when the source mass undergoes CPT transformation. This led Villata to conclude that matter and antimatter should repel each other gravitationally.

Benoit-Lévy and Chardin (2012) developed a cosmological model (the "Dirac-Milne" universe) based on matter–antimatter gravitational repulsion, showing that it could account for several cosmological observations without requiring dark energy or inflation.

---

3. The Negative Mass–Antimatter Hypothesis

3.1 Statement of the Hypothesis

We now develop the central hypothesis of this report in formal terms. The hypothesis consists of four interconnected claims:

Claim 1 (Negative Gravitational Mass): Antimatter possesses negative gravitational mass. Specifically, the active gravitational mass of an antiparticle has the opposite sign to that of the corresponding particle:

m_(g)p = -m_(g)p

while the inertial mass remains positive in magnitude (ensuring positive kinetic energy in the local frame):

|m_ip| = |m_ip|

This means the equivalence principle is violated for antimatter, with m_(g) / m_i = -1 for antiparticles.

Claim 2 (Gravitational Repulsion): As a consequence of Claim 1, matter and antimatter gravitationally repel each other:

F_(grav)(m, m) = +fracG |m_(g)p| |m_(g)p|r^2 r (repulsive)

while antimatter–antimatter gravitational interaction is attractive (like charges repel in gravity, unlike in electromagnetism, because the product of two negative masses is positive).

Claim 3 (Temporal Segregation): Through the Feynman–Stückelberg interpretation, the negative gravitational mass of antimatter is equivalent to propagation backward in time. At the Big Bang singularity (or more precisely, at the CPT-symmetric boundary), antimatter was repelled into the t < 0 direction while matter evolved into t > 0.

Claim 4 (CPT-Symmetric Cosmology): The universe as a whole is exactly CPT-symmetric. What we observe as the "universe" is only the $t > 0$ matter-dominated half. The $t < 0$ half is an antimatter-dominated, parity-reversed, time-reversed mirror image. The baryon asymmetry is not an asymmetry of physics but an anthropic selection effect: observers made of matter can only exist in the matter-dominated sector.

3.2 Connection to CPT Symmetry and General Relativity

Villata (2011) provided the most rigorous connection between CPT symmetry and antigravity for antimatter. His argument proceeds as follows.

In general relativity, the motion of a test particle in a gravitational field is described by the geodesic equation. Consider a matter test particle moving in the field of a matter source. Under CPT transformation, we transform the test particle into an antiparticle (C), reverse spatial coordinates (P), and reverse time (T).

The geodesic equation in the weak-field limit gives:

$$(d^2 r)/(dt^2) = -\nabla \Phi$$

where $\Phi = -GM/r$ is the Newtonian gravitational potential of the source mass M.

Under T: $t \rightarrow -t$, so $d^2r/dt^2 \rightarrow d^2r/dt^2$ (the double time derivative is T-even).

Under P: $r \rightarrow -r$, so $\nabla \rightarrow -\nabla$, and $d^2r/dt^2 \rightarrow -d^2r/dt^2$.

Under C applied to the source: if C reverses the gravitational "charge" (mass) of the source, then $\Phi \rightarrow -\Phi$.

Combining: the CPT-transformed equation becomes:

$$-(d^2 r)/(dt^2) = -(-\nabla)(-\Phi) = -\nabla\Phi$$

which gives:

This is a repulsive gravitational interaction. The antiparticle is repelled by the matter source.

The critical and controversial step in this argument is the assumption that charge conjugation C reverses the sign of gravitational mass. In standard general relativity, mass-energy is always positive and C does not affect it. Villata's argument requires an extension of CPT to the gravitational sector in a specific way that is not mandated by the standard formulation of the CPT theorem (which is proven only for flat-spacetime QFT).

3.3 The Boyle–Finn–Turok CPT-Symmetric Universe

In 2018, Boyle, Finn, and Turok proposed a remarkable cosmological model that realizes the spirit of our Claim 4 without requiring negative mass per se (Boyle, Finn, & Turok, 2018). They noted that the Standard Model of particle physics, combined with general relativity, is CPT symmetric. They then asked: What if the universe itself respects this symmetry?

Their proposal is that the universe does not begin at the Big Bang but rather extends through it, with the Big Bang serving as a fixed point of a CPT transformation. Before the Big Bang (in the $t < 0$ direction), the universe is the CPT mirror of the $t > 0$ universe:

- Matter is replaced by antimatter (C)
- Left and right are reversed (P)
- Time runs in the opposite direction (T)

In this picture, the universe is a CPT-symmetric pair, analogous to a particle–antiparticle pair created from the vacuum. The total CPT charge of the universe is zero.

Critically, Boyle, Finn, and Turok showed that imposing CPT symmetry as a cosmological boundary condition has concrete observable consequences:

condition implies the existence of a stable, massive fermion that behaves exactly like dark matter. They identified this with one of the right-handed neutrinos, predicting its mass to be in a specific range.

1. No inflation needed: The CPT boundary condition naturally explains the low entropy of the early universe without requiring an inflationary epoch.

1. Prediction for neutrino masses: The model predicts that the lightest neutrino is exactly massless.

The Boyle–Finn–Turok model is not identical to our negative-mass antimatter hypothesis but is closely related in spirit. Both frameworks invoke CPT symmetry as a fundamental principle of cosmology. The key difference is that Boyle–Finn–Turok achieve CPT symmetry through boundary conditions on standard fields, while our hypothesis achieves it through negative gravitational mass causing physical separation of matter and antimatter sectors.

A synthesis is possible: the CPT-symmetric boundary condition of Boyle–Finn–Turok could be realized through the mechanism of gravitational repulsion between matter and antimatter. If antimatter has negative gravitational mass, it is naturally repelled toward the $t < 0$ sector, providing a dynamical reason for the CPT symmetry of the cosmological boundary conditions.

3.4 Formal Development: Gravitational CPT and Temporal Segregation

We now present a more formal development of the temporal segregation mechanism. Consider the Einstein field equations:

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = (8\pi G)/(c^4)\, T_{\mu\nu}$$

In a universe containing both matter (with stress-energy $T_{\mu\nu}{}^{(+)}$, positive mass density $\rho > 0$) and antimatter (with stress-energy $T_{\mu\nu}{}^{(-)}$, negative mass density $-\rho < 0$, per our hypothesis):

$$G_{\mu\nu} = (8\pi G)/(c^4)(T_{\mu\nu}{}^{(+)} + T_{\mu\nu}{}^{(-)})$$

If at the initial moment $t = 0$ (the Big Bang), matter and antimatter are created in equal quantities, the total stress-energy is:

$$T_{(\mu\nu)\text{total}} = T_{(\mu\nu)}(^{+}) + T_{(\mu\nu)}(^{-}) = 0$$

The net gravitational effect is zero — consistent with creation from "nothing" (as in the Tryon (1973) and Vilenkin (1982) proposals for the universe as a vacuum fluctuation).

The gravitational repulsion between matter and antimatter sectors drives their separation. In the language of the Friedmann equation for a two-component fluid:

$$H^2 = (8\pi G)/(3)(\rho_{+} + \rho_{-}) = (8\pi G)/(3)(\rho - \rho) = 0$$

This is problematic for a conventional spatial separation (the expansion rate vanishes). However, if we interpret the separation as temporal — with matter evolving toward $t > 0$ and antimatter toward $t < 0$ — then each sector independently satisfies:

$$H^2 = (8\pi G)/(3)\rho$$

with positive ρ in each sector's own temporal frame.

This temporal segregation resolves the baryon asymmetry problem by construction: an observer in the $t > 0$ sector sees only matter and concludes (incorrectly) that there is a matter–antimatter asymmetry. The full CPT-symmetric universe has exact baryon symmetry.

3.5 Relationship Between Negative Mass and Backward Time Propagation

A deep connection exists between negative mass and backward time propagation, rooted in the structure of special relativity. The energy-momentum relation:

$$E^2 = p^2c^2 + m^2c^4$$

admits both positive and negative energy solutions: $E = \pm\sqrt{p^2c^2 + m^2c^4}$.

In quantum mechanics, the time evolution of a state with energy E is:

$$\psi(t) = \psi(0)\, e^{-iEt/\hbar}$$

A negative-energy solution $E < 0$ evolving forward in time is mathematically equivalent to a positive-energy solution evolving backward in time:

$$e^{-i(-|E|)t/\hbar} = e^{+i|E|t/\hbar} = e^{-i|E|(-t)/\hbar}$$

If we further identify negative energy with negative mass through $E = mc^2$, we obtain the equivalence:

Negative mass, forward in time iff Positive mass, backward in time

This is precisely the Feynman–Stückelberg interpretation applied to the gravitational sector. An antiparticle with negative gravitational mass propagating forward in time is equivalent to a particle with positive gravitational mass propagating backward in time. Our hypothesis unifies the Feynman–Stückelberg interpretation of antiparticles with the concept of negative gravitational mass.

SECTION IV

4. Cosmological Implications

4.1 Big Bang Nucleosynthesis

Big Bang nucleosynthesis (BBN) is one of the most precisely tested pillars of standard cosmology. The predicted abundances of light elements (^{4}He, D, ^{3}He, ^{7}Li) depend sensitively on the baryon-to-photon ratio ηB, the number of relativistic species N_(eff), and the neutron lifetime τ_(n).

In the negative-mass antimatter framework, the key question is whether the $t > 0$ matter sector undergoes BBN identically to the standard model. If all antimatter has been expelled to the $t < 0$ sector by

the time of nucleosynthesis (T ~ 1 MeV, t ~ 1 s), then BBN in the matter sector proceeds exactly as in standard cosmology, with no modifications.

The critical test is the timescale of matter–antimatter separation relative to the timescale of BBN. If gravitational repulsion is the only mechanism for separation, we need to estimate whether separation is complete by t ~ 1 s.

In the Dirac-Milne cosmology of Benoit-Lévy and Chardin (2012), where matter and antimatter coexist in the same temporal sector but are gravitationally segregated spatially, BBN requires modifications because antimatter domains still exist and annihilation continues at domain boundaries. They showed that BBN can still produce acceptable light-element abundances in this scenario, but with different physics.

In our temporal-segregation scenario, separation is more complete (the sectors are in different temporal directions), and standard BBN is recovered. This is a significant advantage of the temporal-segregation version of the hypothesis.

4.2 CMB Observations

The cosmic microwave background provides the most detailed snapshot of the early universe at the time of recombination ($z \approx 1100$, $T \approx 3000$ K). The CMB power spectrum is sensitive to:

- The baryon density $\Omega_b h^2$ (through odd/even peak ratios)
- The matter density $\Omega_{(m)} h^2$ (through the matter-radiation equality epoch)
- The spatial curvature $\Omega_{(k)}$ (through the angular diameter distance to last scattering)
- The amplitude and spectral index of primordial perturbations ($A_{(s)}$, $n_{(s)}$)

In the Boyle–Finn–Turok CPT-symmetric model, the CMB predictions are modified primarily through:

1. The absence of inflation, which changes the mechanism for generating primordial perturbations

1. The prediction that the lightest neutrino is massless

Boyle and Turok (2022) showed that the CPT-symmetric model can produce a scale-invariant spectrum of primordial perturbations through the analytic continuation of the universe through the Big Bang, achieving a fit to CMB data comparable to inflation.

If negative-mass antimatter provides the dynamical mechanism for CPT symmetry, additional observational signatures might include:

- Modified primordial gravitational wave spectrum (since gravitational waves probe the full matter content, including any residual negative-mass component)
- Specific signatures in CMB polarization (the B-mode spectrum)
- Modified Sachs-Wolfe effect at large angular scales

4.3 Dark Energy and the Cosmological Constant

Perhaps the most intriguing cosmological implication of the negative-mass antimatter hypothesis concerns dark energy. The observed accelerated expansion of the universe, discovered by Riess et al. (1998) and Perlmutter et al. (1999), is parameterized by a cosmological constant Λ or equivalently a dark energy density:

$$\rho_\Lambda = (\Lambda c^2)/(8\pi G) \approx 5.96 \times 10^{-27} \text{ kg/m}^3$$

corresponding to $\Omega_\Lambda \approx 0.685$.

The cosmological constant problem — why ρ_Λ is 120 orders of magnitude smaller than naive quantum field theory estimates — is arguably the worst prediction in all of physics.

Negative mass offers a potential resolution. If antimatter has negative gravitational mass and is distributed cosmologically (even if temporally segregated), its gravitational effect could manifest as an

would be repulsive and could drive accelerated expansion.

Farnes (2018) developed this idea into a concrete cosmological model. He proposed a continuously created negative-mass fluid that drives cosmic acceleration. In Farnes' model, the modified Friedmann equation is:

$$H^2 = (8\pi G)/(3)(\rho_+ + \rho_-)$$

where $\rho_- < 0$ is continuously created to maintain a constant negative-mass density (analogous to the steady-state creation field of Hoyle and Narlikar). This produces an effective cosmological constant:

$$\Lambda_{(eff)} = -8\pi G \rho_-$$

Farnes showed that this model can fit supernova data, BAO measurements, and CMB observations comparably to ΛCDM. While the continuous creation mechanism is ad hoc, it demonstrates that negative mass can mimic dark energy phenomenologically.

4.4 Large-Scale Structure

The formation of large-scale structure in the universe depends on the growth of gravitational instabilities from primordial density perturbations. In the standard ΛCDM model, cold dark matter provides the gravitational scaffolding for structure formation, with baryons falling into dark matter potential wells after decoupling from photons at recombination.

In the negative-mass antimatter framework, structure formation is modified in several ways:

1. Modified growth rate: If the negative-mass component exerts repulsive gravity, it acts as a negative pressure component (similar to dark energy) that suppresses the growth of structure at late times. This must be consistent with observations of the matter power spectrum P(k) and the growth rate $f\sigma_8$.

d. Void structure: Negative-mass antimatter, being gravitationally repelled by matter, would accumulate in cosmic voids. If any residual negative mass exists in the $\Gamma < 0$ sector, voids would be more empty than in ΛCDM (they would contain negative-mass material that actively pushes matter toward walls and filaments). This could potentially explain the observed "void phenomenon" — the fact that voids appear emptier than ΛCDM predicts (Peebles, 2001).

1. Galaxy rotation curves: In Farnes' model, negative-mass material forms halos around positive-mass galaxies (it is repelled outward but also self-repelling, creating a pressure-supported halo). This negative-mass halo exerts an inward-directed pressure that mimics the effect of a dark matter halo on galaxy rotation curves. However, the detailed profile of such halos and their match to observed rotation curves has not been demonstrated at the precision of NFW or Einasto profiles.

SECTION V

5. Experimental Evidence and Constraints

5.1 The ALPHA-g Experiment at CERN

The most significant experimental development for our hypothesis came in September 2023, when the ALPHA-g collaboration at CERN published the first direct measurement of the gravitational behavior of antimatter (Anderson et al., 2023, Nature 621, 716–722).

The ALPHA-g apparatus traps antihydrogen atoms (a bound state of an antiproton and a positron, electrically neutral) in a vertical magnetic trap. By slowly reducing the trap depth, antihydrogen atoms escape either upward (against Earth's gravity, if antimatter falls down) or downward (with Earth's gravity, if antimatter falls up). The direction of escape is detected by the annihilation signal.

The result: antihydrogen falls down. The ALPHA-g experiment measured:

$(g)/(g) = 0.75 \pm 0.13\ (\text{stat+syst}) \pm 0.16\ (\text{simulation})$

where a is the gravitational acceleration of antihydrogen and $g = 9.81\ m/s^2$. The result is consistent with $g/g = +1$ (normal gravity) and rules out $g/g = -1$ (antigravity) at a significance of approximately 100σ.

This result represents a profound challenge to the simplest version of our hypothesis (Claim 1 in its naive form). If antihydrogen has negative gravitational mass, it should fall upward, and ALPHA-g conclusively shows it falls downward.

5.2 Interpretation of ALPHA-g Results

However, the situation is more nuanced than it might initially appear. Several important caveats must be noted:

1. Precision: The ALPHA-g result is consistent with normal gravity but has ~20% uncertainties. It does not yet test the weak equivalence principle at high precision. There is room for a small anomalous gravitational component.

2. What is measured: ALPHA-g measures the gravitational behavior of antihydrogen (one antiproton + one positron) in Earth's gravitational field. This is the gravitational interaction of antimatter with the overwhelmingly matter-dominated local environment. If our hypothesis involves a more subtle modification — for example, if the gravitational mass sign depends on the cosmological sector rather than the particle identity — then ALPHA-g might not be testing the relevant regime.

3. Renormalization of gravitational mass: Some theoretical frameworks (e.g., Hajdukovic, 2011) suggest that the gravitational mass of bound antimatter states could differ from that of free antimatter due to virtual pair effects. The gravitational mass measured by ALPHA-g might not be the "bare" gravitational mass of antiparticles.

4. The temporal-segregation version: The strongest version of our hypothesis — that antimatter was repelled into the $t < 0$ sector at the Big Bang — is not directly tested by ALPHA-g. ALPHA-g tests how antihydrogen (produced and confined in the $t > 0$ sector) behaves in a local gravitational field. If the temporal segregation is a cosmological boundary effect rather than a local force, antihydrogen created in

the laboratory might behave normally in local gravity while the cosmological mechanism operates through different physics.

5.3 AEGIS and GBAR Experiments

Two other antimatter gravity experiments at CERN are pursuing complementary approaches:

AEGIS (Antimatter Experiment: Gravity, Interferometry, Spectroscopy) aims to measure the gravitational acceleration of antihydrogen using a Moiré deflectometer, targeting a precision of 1% on g/g (Kellerbauer et al., 2008).

GBAR (Gravitational Behaviour of Antihydrogen at Rest) takes a different approach, producing ultracold antihydrogen ions (H^+), cooling them sympathetically, then photodetaching the extra positron and observing the free-fall trajectory of the resulting neutral H (Pérez et al., 2015). GBAR aims for precision at the percent level and eventually much better.

These experiments will progressively tighten the constraints on anomalous gravitational behavior of antimatter. If g/g is found to be exactly 1 at high precision, the naive negative-mass hypothesis is conclusively excluded. If a small deviation is detected (for example, g/g = 1.001 or 0.999), this would be a discovery of enormous significance, potentially supporting a modified version of the hypothesis.

5.4 Indirect Constraints

Several indirect constraints on the gravitational properties of antimatter exist:

1. Positronium free-fall: Positronium (e^+e^- bound state) is an equal mixture of matter and antimatter. If antimatter has negative gravitational mass, positronium should have zero gravitational mass and not fall at all. While direct positronium free-fall has not been measured, the gravitational redshift of positronium annihilation photons could provide constraints.

2. Kaon oscillations and CP violation: Chardin and Rax (1992) argued that the observed CP violation in the neutral kaon system could be explained by a gravitational energy difference between the K^o (containing a s quark) and K^o (containing an s quark) in Earth's gravitational field. However, this

explanation has been challenged by the observation that CP violation in the B-meson system has a different pattern than the gravitational explanation would predict.

3. Equivalence principle tests with nuclear binding energy: Atoms contain significant contributions to their mass from virtual quark–antiquark pairs (sea quarks). The fraction of an atom's mass due to virtual antimatter is of order 10%. The precision of equivalence principle tests (10^{-15} by MICROSCOPE) therefore constrains the difference between matter and antimatter gravitational mass at the ~10^{-14} level (Adelberger, Heckel, & Nelson, 2003).

This is perhaps the strongest indirect argument against our hypothesis: if virtual antimatter within matter had negative gravitational mass, the gravitational mass of different elements (which have different virtual antimatter content) would differ, violating the equivalence principle at a level detectable by existing experiments. However, this argument assumes that virtual pairs contribute to gravitational mass in the same way as real particles — an assumption that is plausible but not proven.

5.5 Neutrino-Antineutrino Oscillations

If neutrinos and antineutrinos have different gravitational masses, their oscillation behavior in gravitational fields would differ. The MINOS experiment and subsequent long-baseline neutrino experiments have measured neutrino and antineutrino oscillation parameters. The observed CP-violating phase δ_(CP) in neutrino oscillations (T2K Collaboration, Abe et al., 2020) could potentially receive contributions from gravitational effects if our hypothesis is correct. However, the current precision of neutrino oscillation measurements is insufficient to distinguish gravitational CP violation from intrinsic CP violation.

SECTION VI

6. Relationship to Dark Matter and Dark Energy

6.1 The Dark Sector Problem

The standard ΛCDM cosmological model requires two components with no known particle physics counterpart:

- Dark matter ($\Omega_{DM} \approx 0.265$): gravitationally attractive, non-baryonic, cold, collisionless
- Dark energy ($\Omega_\Lambda \approx 0.685$): gravitationally repulsive, uniform, constant density

Together, these "dark" components constitute ~95% of the energy density of the universe, while the ordinary matter we understand constitutes only ~5%. This situation has been described as a crisis for fundamental physics.

6.2 Farnes' Negative-Mass Cosmology

Farnes (2018) proposed a unified framework in which a single component — a negative-mass fluid — simultaneously explains both dark matter and dark energy. The key features of the model are:

1. Continuous creation: Negative-mass particles are continuously created, maintaining a constant negative-mass density $\rho_- = \text{const}$ even as the universe expands. This is analogous to the C-field of Hoyle and Narlikar (1964) in steady-state cosmology.

1. Effective cosmological constant: The constant negative-mass density produces an effective cosmological constant:

$$\Lambda_{(eff)} = -8\pi G\rho_- > 0$$

This drives the observed accelerated expansion.

1. Dark matter mimicry: Negative-mass particles are repelled by positive-mass galaxies and accumulate in a halo surrounding each galaxy. From the galaxy's perspective, this halo exerts an effective inward pressure (the negative-mass material pushes on the galaxy from outside). Farnes showed through N-body simulations that this can flatten galaxy rotation curves qualitatively.

measurements of the amplitude of matter fluctuations).

6.3 Connecting Farnes' Model to the Antimatter Hypothesis

If the negative-mass fluid in Farnes' model is identified with antimatter, we obtain a framework where:

- Antimatter = negative gravitational mass
- Most antimatter is in the $t < 0$ CPT-mirror sector (explaining baryon asymmetry)
- Residual negative-mass antimatter in the $t > 0$ sector provides dark energy and contributes to dark matter phenomenology
- The continuous creation mechanism could be understood as quantum pair production from the vacuum, where the negative-mass component is the antimatter produced

However, several challenges arise:

1. ALPHA-g: If antimatter is the negative-mass fluid, why does antihydrogen fall down?

1. Annihilation: Why doesn't the negative-mass antimatter in the halo annihilate with the positive-mass matter in the galaxy? (Farnes' model uses a generic negative-mass fluid, not specifically antimatter, partly to avoid this issue.)

1. BBN constraints: Significant quantities of free antimatter at $T \sim 1$ MeV would modify nucleosynthesis through annihilation processes.

6.4 The Boyle–Finn–Turok Dark Matter Candidate

The CPT-symmetric universe model of Boyle, Finn, and Turok offers a different connection to dark matter. Their model predicts that the CPT boundary condition at $t = 0$ selects a specific vacuum state for fermionic fields. For a theory with three right-handed neutrinos (needed for the seesaw mechanism), the

CPT condition requires that one of these neutrinos be stable, with a mass determined by cosmological constraints.

Boyle and Turok (2022) showed that this CPT-mandated right-handed neutrino has the correct relic abundance to serve as dark matter if its mass is approximately:

$m_{(N)} \sim 4.8 \times 10^8$ GeV

This is a remarkably specific prediction from a minimal framework. The dark matter candidate is "ordinary" in the sense that it has positive mass and does not require exotic physics, yet it emerges naturally from the CPT symmetry requirement.

If we combine the Boyle–Finn–Turok framework with our negative-mass antimatter hypothesis, we obtain a picture where:

- Dark matter is a CPT-mandated right-handed neutrino
- Dark energy is explained by the residual gravitational effect of the $t < 0$ antimatter sector
- The baryon asymmetry is resolved by temporal segregation

6.5 Alternative: Negative-Mass Antimatter as Dark Energy Only

A more conservative version of the hypothesis abandons the attempt to explain dark matter through negative mass and focuses solely on dark energy. In this version:

- Dark matter is a separate, conventional (positive-mass) particle (e.g., a WIMP or axion)
- Antimatter has negative gravitational mass and has been temporally segregated
- The boundary between the $t > 0$ and $t < 0$ sectors imposes effective boundary conditions on the gravitational field that produce accelerated expansion

This version is less ambitious but faces fewer immediate challenges. The accelerated expansion of the universe would be a consequence of the universe's CPT-symmetric structure rather than requiring a separate dark energy component.

SECTION VII

7. Mathematical Formalism

7.1 Modified Einstein Field Equations with Negative Mass

We begin by writing the Einstein field equations for a universe containing both positive-mass matter (density ρ_+, pressure p_+) and negative-mass antimatter (density $\rho_- < 0$, pressure p_-):

$$G_{\mu\nu} = (8\pi G)/(c^4)(T_{\mu\nu}^{(+)} + T_{\mu\nu}^{(-)})$$

For a perfect fluid with negative mass:

$$T_{\mu\nu}^{(-)} = (\rho_- + (p_-)/(c^2))u_\mu u_\nu + p_- g_{\mu\nu}$$

where $\rho_- < 0$ and we take $p_- = w_- \rho_- c^2$ with equation of state parameter w_-.

7.2 Modified Friedmann Equations

For a homogeneous, isotropic universe (FLRW metric) containing both positive- and negative-mass components:

$$ds^2 = -c^2 dt^2 + a(t)^2[(dr^2)/(1-kr^2) + r^2 d\Omega^2]$$

The Friedmann equations become:

$$H^2 \equiv ((a)/(a))^2 = (8\pi G)/(3)(\rho_+ + \rho_-) - (kc^2)/(a^2)$$

Case 1: Equal and opposite matter and antimatter ($\rho_+ + \rho_- = 0$).

If the universe contains equal quantities of positive and negative mass:

$$H^2 = -(kc^2)/(a^2)$$

For k = -1 (open universe), $H^2 = c^2/a^2$, giving a(t) = ct. This is the coasting or Milne cosmology, which has a = 0 — neither accelerating nor decelerating. This is the basis of the Dirac-Milne cosmology (Benoit-Lévy & Chardin, 2012).

Case 2: Temporal segregation with residual negative mass.

If most antimatter is in the t < 0 sector but a fraction f remains in t > 0:

$$\rho_{(total)} = \rho_+ - f\rho_+ = (1-f)\rho_+$$

The Friedmann equation for the matter-dominated era becomes:

$$H^2 = (8\pi G)/(3)(1-f)\rho_+ - (kc^2)/(a^2)$$

For the acceleration equation:

$$(a)/(a) = -(4\pi G)/(3)(1-f)\rho_+ + (4\pi G)/(3) \cdot 3 f \rho_+ w_-$$

If the residual negative mass has $w_- < -1/3$ (or if the negative-mass contribution acts effectively as a cosmological constant), this can produce accelerated expansion.

Case 3: Farnes' continuously created negative mass.

Farnes (2018) imposes the condition ρ_- = const, so that as the universe expands, new negative-mass material is continuously created to maintain constant density. The continuity equation:

where Γ is the creation rate, required to satisfy $\rho_- = 0$:

$$\Gamma = -3H(\rho_- + p_-/c^2)$$

For $\rho_- < 0$ and dust-like negative matter ($p_- = 0$):

$$\Gamma = -3H\rho_- > 0$$

The Friedmann equation becomes:

$$H^2 = (8\pi G)/(3)\rho_+ + (8\pi G)/(3)\rho_- = (8\pi G)/(3)\rho_+ - (|\rho_-|8\pi G)/(3)$$

The constant negative-mass density acts exactly as a positive cosmological constant: $\Lambda_{(eff)} = 8\pi G |\rho_-|$.

7.3 Negative Mass in the Schwarzschild Solution

The Schwarzschild solution for a point mass M is:

$$ds^2 = -(1 - (2GM)/(c^2 r))c^2 dt^2 + (1 - (2GM)/(c^2 r))^{-1}dr^2 + r^2 d\Omega^2$$

For $M < 0$ (negative mass):

$$ds^2 = -(1 + (2G|M|)/(c^2 r))c^2 dt^2 + (1 + (2G|M|)/(c^2 r))^{-1}dr^2 + r^2 d\Omega^2$$

Key features of the negative-mass Schwarzschild solution (Bonnor, 1989):

1. No horizon: The factor $(1 + 2G|M|/c^2r) > 1$ for all $r > 0$, so there is no event horizon. The singularity at $r = 0$ is naked.

1. Repulsive gravity: Test particles are repelled. The geodesic equation in the Newtonian limit gives:

$$(d^2r)/(dt^2) = +(G|M|)/(r^2)$$

d. Negative Komar mass: The Komar mass integral yields $M_{(K)} = M < 0$, consistent with the ADM mass.

1. Gravitational lensing: A negative-mass object acts as a *diverging* gravitational lens, spreading light rays apart rather than focusing them. This would produce a characteristic anti-lensing signature observable in principle (Izaguirre, Jia, & Stojkovic, 2020).

7.4 Geodesic Equation for Negative-Mass Test Particles

For a test particle with inertial mass m_i and gravitational mass $m_{(g)} = -|m_{(g)}|$ (if $m_{(g)} < 0$ while $m_i > 0$, violating the equivalence principle), the equation of motion in a gravitational field is:

$$m_i \, (d^2 x^\mu)/(d\tau^2) = m_{(g)} \, (-\Gamma^\mu_{(\alpha\beta)}(dx^\alpha)/(d\tau)(dx^\beta)/(d\tau)) \cdot (m_i)/(m_{(g)})$$

Wait — let us be more careful. The geodesic equation is:

$$(d^2 x^\mu)/(d\tau^2) + \Gamma^\mu_{(\alpha\beta)}(dx^\alpha)/(d\tau)(dx^\beta)/(d\tau) = 0$$

This is independent of the mass of the test particle (this is the universality of free fall / equivalence principle). If we maintain the equivalence principle but allow both m_i and $m_{(g)}$ to be negative (Bondi's approach), then negative-mass test particles follow the same geodesics as positive-mass particles — they fall toward positive masses just as positive masses do.

However, if we violate the equivalence principle by having $m_{(g)} < 0$ while $m_i > 0$ (or vice versa), the equation of motion becomes:

$$m_i \, a^\mu = m_{(g)} \, \nabla^\mu \Phi$$

$$a^\mu = (m_{(g)})/(m_i) \, \nabla^\mu \Phi$$

For $m_{(g)}/m_i = -1$ (our antimatter hypothesis): the acceleration is in the opposite direction. The antiparticle falls up in a gravitational field.

This violation of the equivalence principle is the most radical aspect of our hypothesis and the one most directly constrained by experiment.

7.5 Quantum Field Theory of Gravitationally Repulsive Antimatter

A proper quantum field theory treatment requires coupling the Dirac field to gravity. In curved spacetime, the Dirac equation is:

$$(i\gamma^\mu \nabla_\mu - (mc)/(\hbar))\psi = 0$$

where $\nabla_\mu = \partial_\mu + (1)/(4)\omega_{\mu ab}\gamma_a\gamma b$ is the spinor covariant derivative and $\omega_{\mu ab}$ is the spin connection.

Under CPT, the Dirac field transforms as:

$$\psi \text{ xrightarrowCPT } \eta_{(CPT)}\ \gamma_5\ \psi^*$$

where $\eta_{(CPT)}$ is a phase. Villata (2011) argued that applying this transformation to the Dirac equation in curved spacetime produces:

$$(i\gamma^\mu \nabla_\mu + (mc)/(\hbar))\psi_{(CPT)} = 0$$

Note the sign change of the mass term. If the gravitational coupling depends on this sign, the CPT-transformed field (antimatter) couples with opposite sign to gravity.

However, in standard QFT, both signs of the mass term yield physically equivalent theories (related by a chiral rotation $\psi \to \gamma_5\ \psi$ for massless or by field redefinition for massive fields). The claim that the sign change has gravitational consequences is therefore controversial and depends on the precise coupling between spin and gravity.

7.6 Energy Conditions and Negative Mass

General relativity constrains the stress-energy tensor through various energy conditions:

1. Strong Energy Condition (SEC): $(T_{\mu\nu} - \frac{1}{2}T g_{\mu\nu})u^\mu u^\nu \geq 0$. This implies $\rho + 3p/c^2 \geq 0$.

1. Null Energy Condition (NEC): $T_{\mu\nu} k^\mu k^\nu \geq 0$ for all null k^μ.

1. Dominant Energy Condition (DEC): $T_{\mu\nu} u^\mu$ is non-spacelike for timelike u^μ.

Negative-mass matter violates the WEC ($\rho < 0$). Notably, the cosmological constant / dark energy already violates the SEC ($\rho + 3p/c^2 = \rho_\Lambda + 3(-\rho_\Lambda) = -2\rho_\Lambda < 0$). The energy conditions are not fundamental laws — they are conjectures about the behavior of classical matter. Quantum effects routinely violate the WEC (e.g., the Casimir effect produces regions of negative energy density).

If negative-mass antimatter exists, the WEC is violated, but this is not necessarily fatal to the theory. The key requirement is that the total energy of any isolated system remains non-negative to ensure stability of the vacuum. In the CPT-symmetric framework, the total energy of the universe (summing both temporal sectors) is exactly zero, consistent with creation from nothing.

SECTION VIII

8. Criticisms and Counterarguments

8.1 The Runaway Paradox

The objection: Bondi (1957) showed that a positive-mass particle and a negative-mass particle, interacting gravitationally, form a "runaway pair" — the negative mass chases the positive mass, both accelerating without limit. This appears to violate energy conservation and common sense.

Response: The runaway pair actually does conserve energy and momentum. The kinetic energy of the negative-mass particle is $E_k = \frac{1}{2}m v^2 < 0$ (since $m < 0$), and the total kinetic energy of the pair remains constant (one positive, one negative, summing to zero). Similarly, momentum is conserved

because the negative-mass particle carries negative momentum. The runaway pair is counterintuitive but not inconsistent.

However, in the temporal-segregation version of our hypothesis, the runaway paradox does not arise because matter and antimatter occupy different temporal sectors and never interact locally after the Big Bang event. The gravitational repulsion operates only at the cosmological boundary, not as a local force between nearby particles.

8.2 The ALPHA-g Result

The objection: The 2023 ALPHA-g experiment directly measured that antihydrogen falls downward in Earth's gravity, ruling out antigravity at high significance.

Response: This is the most serious experimental challenge to the hypothesis. We acknowledge that the simplest version — antimatter has negative gravitational mass and falls up — is ruled out by ALPHA-g.

However, several responses are available:

1. The hypothesis is cosmological, not local. The temporal-segregation mechanism may operate through cosmological boundary conditions rather than through a local modification of gravity. Antihydrogen produced in the laboratory exists within the $t > 0$ sector and interacts normally with local gravity, while the cosmological CPT symmetry operates at the level of initial/boundary conditions.

1. Screening effects. In some theoretical frameworks, the gravitational mass of antimatter could be "screened" by the dominant matter environment. In the matter-dominated $t > 0$ sector, the antimatter might be gravitationally "dressed" by its interaction with the matter vacuum, acquiring effective positive gravitational mass (analogous to charge screening in QED).

1. The gravitational mass distinction may be between sectors, not particles. In the Boyle–Finn–Turok framework, the gravitational repulsion is not between individual particles but between the two temporal sectors of the universe. Individual antiparticles within our sector behave normally.

Warp-like response (3) effectively concedes the simple negative-mass hypothesis and replaces it with a more subtle version based on cosmological boundary conditions. This is a legitimate theoretical move but reduces the explanatory power and predictive specificity of the framework.

8.3 Virtual Antimatter and the Equivalence Principle

The objection: Atoms contain significant virtual antimatter (sea quarks, virtual electron-positron pairs). If virtual antimatter had negative gravitational mass, different atoms would have different ratios of gravitational to inertial mass, violating the equivalence principle. MICROSCOPE and torsion balance experiments constrain equivalence principle violations at the 10^{-15} level, seemingly ruling out any difference.

Response: This is a powerful argument in the context of local physics. Possible responses:

1. Virtual pairs may not contribute to gravitational mass in the same way as real particles. Virtual particles are off-shell ($E^2 \neq p^2c^2 + m^2c^4$) and their gravitational properties may differ. The theoretical question of how virtual particles gravitate is unsettled.

1. Renormalization. In renormalized QFT, the physical mass of a particle includes contributions from virtual pairs, but these contributions are already accounted for in the measured inertial mass. If gravitational mass equals total renormalized mass (as the equivalence principle demands), then virtual pairs do not produce a separate gravitational anomaly.

1. The distinction applies only to real, free antiparticles. Virtual antimatter within matter is part of the quantum vacuum dressing of the matter particle and gravitates normally. Only free, on-shell antiparticles possess negative gravitational mass. This is *ad hoc* but not inconsistent.

8.4 Energy Conservation

The objection: If matter has positive energy and antimatter has negative energy, their annihilation should produce zero energy, not the observed radiation (e.g., electron-positron annihilation produces two 511 keV photons).

Response: This objection confuses gravitational mass/energy with inertial mass/energy. In our framework:

- The *inertial* mass of both particles and antiparticles is positive.
- The *gravitational* mass of antiparticles is negative.
- Annihilation converts inertial mass-energy into radiation: $e^+ + e^- \rightarrow 2\gamma$, releasing $2m_e c^2 = 1.022$ MeV. This is the inertial energy, which is positive for both.
- The *gravitational* energy balance is: the gravitational mass goes from $(+m_e) + (-m_e) = 0$ to two photons with gravitational mass $+E/c^2$ each. This appears to violate gravitational energy conservation.

This is a genuine problem. The resolution might be:

1. Photons as their own antiparticles have zero net gravitational charge in a CPT sense. The gravitational energy of annihilation photons comes from the conversion of kinetic/binding energy, not from gravitational mass.

1. The equivalence of gravitational and inertial mass for photons is well-tested (gravitational lensing, Shapiro delay), so photons definitely have positive gravitational mass. This creates a genuine bookkeeping problem for the hypothesis.

We acknowledge this as one of the most serious theoretical difficulties with the framework.

8.5 Why Mainstream Physics Rejects Negative Mass

The mainstream rejection of negative gravitational mass for antimatter rests on several pillars:

1. The equivalence principle is the foundation of general relativity and has been tested to extraordinary precision. Violating it requires abandoning or modifying GR.

gravitational mass. However, the CPT theorem is proven for flat-spacetime QFT and its extension to curved spacetime (where gravity is present) is not rigorous.

1. Positive energy theorems (Schoen & Yau, 1979; Witten, 1981) prove that the total energy of an asymptotically flat spacetime satisfying the dominant energy condition is non-negative. Negative mass violates the dominant energy condition.

1. Stability concerns: A universe containing both positive and negative mass has no ground state — energy could be extracted indefinitely by separating positive and negative mass-energy.

1. ALPHA-g: As discussed, direct measurement shows antihydrogen falls down.

These objections are serious. We note, however, that:

- The equivalence principle is empirical, not derived from first principles
- The CPT theorem has not been extended to quantum gravity
- Dark energy already violates the strong energy condition
- The stability concern is addressed by the CPT-symmetric framework (total energy is zero)
- ALPHA-g tests local gravity, not cosmological CPT

8.6 Photon Self-Conjugacy Problem

The objection: The photon is its own antiparticle. If our hypothesis assigns negative gravitational mass to antiparticles, what gravitational mass does the photon have? It cannot be both positive and negative.

Response: The photon, being its own antiparticle, has zero "gravitational charge" in the matter/antimatter sense. Its gravitational interaction comes entirely from its energy-momentum, which is always positive. The mass/antimass distinction applies only to particles that are distinct from their

antiparticles (Dirac-type particles). For Majorana-type particles (self-conjugate), the gravitational mass is determined by their energy content, as in standard GR.

This actually provides a consistency check: self-conjugate particles (photons, Z^0, Higgs, graviton, and potentially Majorana neutrinos) must gravitate normally. Only Dirac-type particles (quarks, charged leptons, Dirac neutrinos if applicable) have distinct antiparticles that could carry negative gravitational mass.

SECTION IX

9. Conclusions and Future Directions

9.1 Summary of the Hypothesis

We have explored the hypothesis that the observed baryon asymmetry of the universe can be explained by antimatter possessing negative gravitational mass, leading to temporal segregation of matter and antimatter at the Big Bang into CPT-conjugate sectors of the universe. This hypothesis draws on:

- The Feynman–Stückelberg interpretation (antiparticles as backward-in-time particles)
- Bondi's negative-mass framework in general relativity
- Villata's CPT argument for matter-antimatter gravitational repulsion
- Boyle, Finn, and Turok's CPT-symmetric universe model
- Farnes' negative-mass cosmology

9.2 Where the Hypothesis is Strong

1. Elegance and economy: The hypothesis resolves the baryon asymmetry problem without requiring new physics beyond gravity. No new particles, no new symmetry breaking, no new CP violation —

1. CPT symmetry as a cosmological principle: The idea that the universe as a whole respects CPT symmetry is aesthetically compelling and has been shown by Boyle, Finn, and Turok to have concrete, testable consequences (prediction of dark matter candidate, lightest neutrino mass = 0).

1. Connection to dark energy: The negative-mass framework offers a conceptual path toward understanding the cosmological constant as a consequence of the universe's CPT structure rather than as a separate, unexplained component.

1. Creation from nothing: If the total mass-energy of the universe is zero (positive mass in $t > 0$, negative mass in $t < 0$), the universe can arise as a quantum fluctuation from nothing, connecting to the proposals of Tryon (1973) and Vilenkin (1982).

1. Testability: The hypothesis makes predictions that are being actively tested (antimatter gravity at CERN, neutrino mass hierarchy, CMB signatures).

9.3 Where the Hypothesis is Weak or Speculative

1. ALPHA-g: The most direct test — does antihydrogen fall up or down? — has returned a result inconsistent with the simplest version of the hypothesis. Antihydrogen falls down.

1. Energy conservation in annihilation: The bookkeeping of gravitational energy in matter–antimatter annihilation is problematic.

1. Virtual antimatter and the equivalence principle: Indirect constraints from precision tests of the equivalence principle are severe, though not conclusive due to ambiguities in how virtual particles gravitate.

1. Violation of the equivalence principle: The hypothesis, in its simplest form, requires violating the foundational principle of general relativity.

1. This CPT argument is contested: The extension of CPT to the gravitational sector (necessary for Villata's argument) is not supported by a rigorous proof.

1. Quantitative predictions are lacking: The hypothesis, as developed here, makes qualitative predictions but lacks the precise quantitative predictions (e.g., CMB power spectrum) that would allow a statistical comparison with ΛCDM.

9.4 Future Directions

1. Precision antimatter gravity: ALPHA-g, AEGIS, and GBAR at CERN will progressively improve the precision of g/g measurements. A result of exactly 1.000 would close the local-antigravity version of the hypothesis. Any deviation, however small, would be revolutionary.

1. CPT-symmetric cosmology: The Boyle–Finn–Turok program is developing precise predictions for the CMB and large-scale structure. Comparison with Planck and future CMB experiments (CMB-S4, LiteBIRD) can test the framework.

1. Neutrino mass: The prediction that the lightest neutrino is massless can be tested by next-generation neutrino experiments (JUNO, DUNE, Project 8/KATRIN).

1. Gravitational wave astronomy: The negative-mass framework predicts specific modifications to the gravitational wave background and potentially to the inspiral of compact objects containing antimatter (though natural antimatter compact objects are not expected to exist in the $t > 0$ sector).

1. Anti-lensing searches: A negative-mass concentration would act as a diverging gravitational lens. Systematic searches for anti-lensing signatures in large surveys (Euclid, Vera Rubin Observatory) could constrain or detect negative-mass concentrations.

1. Theoretical development: A rigorous quantum gravity treatment of the gravitational properties of antimatter is needed. This requires progress on the quantization of gravity itself — arguably the biggest open problem in theoretical physics.

d. Positronium gravity: A measurement of the free-fall acceleration of positronium (a matter–antimatter bound state) would provide a unique test. If antimatter has negative gravitational mass, positronium should have zero gravitational mass and float.

9.5 Final Assessment

The negative-mass antimatter hypothesis, in its simplest and most dramatic form, is now empirically disfavored by the ALPHA-g experiment. However, the deeper insight — that CPT symmetry may be a fundamental cosmological principle, and that the baryon asymmetry may reflect the structure of the universe rather than a dynamical process — remains viable and scientifically productive. The Boyle–Finn–Turok CPT-symmetric universe, which captures the essential idea without requiring locally measurable antigravity, is an active and promising research program.

The history of physics teaches that our deepest puzzles often require reconsidering our most basic assumptions. The baryon asymmetry problem may ultimately be resolved by new sources of CP violation within conventional baryogenesis, or it may point toward a radically different understanding of the relationship between matter, antimatter, gravity, and time. The experiments and theoretical developments of the coming decade will be decisive.

Report completed February 2026. The author acknowledges that this work is speculative and is intended as a rigorous exploration of a non-standard hypothesis, not as advocacy for its correctness.

© 2026 IUPPITER GROUP · ALL RIGHTS RESERVED

The views expressed are those of the author and represent independent academic analysis.