Foam: A Python package for Forward Asteroseismic Modelling of gravity modes

Mathias Michielsen [1]

1 Institute of Astronomy, KU Leuven, Celestijnenlaan 200D, B-3001 Leuven, Belgium

Summary

Foam is a python package to perform forward asteroseismic modelling of gravity modes. It automates and streamlines a considerable fraction of the modelling process, and can be configured to use various different modelling methodologies. This includes different ways to match the theoretically predicted oscillations to observations, the option to use different sets of observables to compare to their theoretically predicted values, the use of different merit functions to determine the goodness of fit, and the option to consider nested subgrids in a statistically meaningfull way. See Michielsen et al. (2021) and Michielsen et al. (2023) (accepted) for an application of these methodologies to model observed gravity modes.

Statement of need

Stars spend approximately 90% of their evolution on their so called *main sequence*, during which they fuse fusing hydrogen into helium in their cores. In stars with masses above about 1.2 times the mass of the sun, the stellar core in which these fusion processes take place becomes convective. Macroscopic element transport in and near the convective cores of these stars has a large influence on their life, since it transports additional hydrogen from outside of the nuclear fusion region into this region. In this way it both prolongs the main-sequence lifetime of stars and enlarges the mass of the helium core at the end of the main sequence, which significantly influences all later stages of their evolution. However, these transport processes provide the largest uncertainties in stellar structure and evolution models for stars with convective cores, due to our poor understanding of macroscopic element transport and limited number of useful observations to test the theories. (See e.g. Anders & Pedersen (2023) for a review on this topic.)

Through asteroseismology, the study of stellar pulsations, we gain the means to unravel the interior structure of stars (Conny Aerts et al., 2010; C. Aerts, 2021). Gravity modes in particular have a high sensitivity to the properties of the near-core region. We can exploit the probing power of gravity modes, observed in e.g. Slowly Pulsating B-type stars (Waelkens, 1991), to investigate the physics in the interior of these stars, particularly the transition region between the convective core and radiative envelope.

Foam was developed to streamline the forward modelling process of gravity modes, including multiple options and considerations in the statistical analysis of the grids of stellar equillibrium models.

Acknowledgements

39 No. 11F7120N.

References

41 Aerts, C. (2021). Probing the interior physics of stars through asteroseismology. *Reviews of*
42   *Modern Physics*, *93*(1), 015001. https://doi.org/10.1103/RevModPhys.93.015001

43 Aerts, Conny, Christensen-Dalsgaard, J., & Kurtz, D. W. (2010). *Asteroseismology*. Springer,
44   Astronomy; Astrophysics Library.

45 Anders, E. H., & Pedersen, M. G. (2023). Convective Boundary Mixing in Main-Sequence
46   Stars: Theory and Empirical Constraints. *Galaxies*, *11*(2), 56. https://doi.org/10.3390/
47   galaxies11020056

48 Michielsen, M., Aerts, C., & Bowman, D. M. (2021). Probing the temperature gradient in
49   the core boundary layer of stars with gravito-inertial modes. The case of KIC 7760680.
50   *Astronomy and Astrophysics*, *650*, A175. https://doi.org/10.1051/0004-6361/202039926

51 Waelkens, C. (1991). Slowly pulsating B stars. *Astronomy and Astrophysics*, *246*, 453.