




Foam: A Python package for forward asteroseismic modelling of gravity modes

Mathias Michielsen ¹

¹ Institute of Astronomy, KU Leuven, Celestijnenlaan 200D, B-3001 Leuven, Belgium

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright
and release the work under a
Creative Commons Attribution 4.0
International License ([CC BY 4.0](#))

Summary

Asteroseismology, the study of stellar pulsations, offers insights into the internal structures and evolution of stars. Analysing the variations in a star's brightness allows the determination of fundamental properties such as mass, radius, age, and chemical composition. Asteroseismology heavily relies on computational tools, but a significant number of them are closed-source, thus inaccessible to the broader astronomic community. This manuscript presents Foam, a python package designed to perform forward asteroseismic modelling of stars exhibiting gravity modes. It automates and streamlines a considerable fraction of the modelling process, comparing grids of theoretical stellar models and their oscillation frequencies to observed frequency sets in stars.

Foam offers the flexibility to employ diverse modelling approaches, allowing users to choose different methodologies for matching theoretically predicted oscillations to observations. It provides options to utilise various sets of observables for comparison with their theoretical counterparts, employ different merit functions for assessing goodness of fit, and to incorporate nested subgrids in a statistically rigorous manner. For applications of these methodologies in modelling observed gravity modes, refer to Michielsen et al. (2021) and Michielsen et al. (2023).

Introduction

Stars spend approximately 90% of their evolution on their so called *main sequence*, during which they fuse hydrogen into helium in their cores. In stars with masses above about 1.2 times the mass of the sun, the stellar core in which these fusion processes take place becomes convective. Macroscopic element transport in and near the convective cores of these stars has a large influence on their life, since it transports additional hydrogen from outside of the nuclear fusion region into this region. In this way it both prolongs the main-sequence lifetime of stars and enlarges the mass of the helium core at the end of the main sequence, which significantly influences all later stages of their evolution. However, these transport processes provide the largest uncertainties in stellar structure and evolution models for stars with convective cores, due to our poor understanding of macroscopic element transport and limited number of useful observations to test the theories. (See e.g. [Anders & Pedersen, 2023](#) for a review on this topic.)

Through asteroseismology, we gain the means to unravel the interior structure of stars ([Aerts et al., 2010](#); [Aerts, 2021](#)). Gravity (g-) modes in particular have a high sensitivity to the properties of the near-core region. These modes have buoyancy as their restoring force, have dominantly horizontal displacements, and oscillate with a period of several hours to a few days. Additionally, they can only propagate in the non-convective regions in the star, which makes their propagation cavity very sensitive to the size of the convective core. We can exploit the probing power of g-modes, observed in e.g. Slowly Pulsating B-type stars ([Waelkens, 1991](#)), to

41 investigate the physics in the interior of these stars, particularly the transition region between
42 the convective core and radiative envelope.

43 Statement of need

44 Some tools have been developed and made publicly available to model and determine stel-
45 lar parameters of solar-like oscillators, such as AIMS (Rendle et al., 2019), BASTA (Aguirre
46 Børsen-Koch et al., 2022), and pySYD (Chontos et al., 2022). However, there are several key
47 differences between the modelling of the pressure (p-) modes observed in solar-like oscillators,
48 and the modelling of the g-modes observed in more massive stars. First and foremost, the
49 well-known asteroseismic scaling relations used for solar-like oscillators cannot be extrapolated
50 to main-sequence stars with a convective core. Secondly, the effect of rotation on p-modes is
51 often included in a perturbative way, whereas the g-mode frequencies are strongly dependent
52 on rotation and require the inclusion of the Coriolis acceleration in a non-perturbative way.
53 Additionally the mass regime of stars with convective cores is subject to strong correlations
54 between several model parameters, which sometimes follow non-linear relationships. In this con-
55 text, the Mahalanobis distance (MD) (see Aerts et al., 2018 for its application to asteroseismic
56 modelling) provides a more appropriate merit function than the often used χ^2 , since it tackles
57 both these non-linear correlations and includes uncertainties for the theoretical predictions.
58 The use of a different, more appropriate merit function significantly impacts modelling results.
59 This is demonstrated by Michielsen et al. (2021) in their comparison between the results
60 obtained by employing the MD versus χ^2 , applied in the modelling of an observed star.

61 Foam was developed to be complimentary to the available modelling tools for solar-like oscillators.
62 It provides a framework for the forward modelling of g-modes in main-sequence stars with
63 convective cores, and tackles the differences in the modelling approach as compared to the
64 case of solar-like oscillators. Foam therefore extends the efforts to provide publicly available,
65 open-source tools for asteroseismic modelling to the g-mode domain, given that the currently
66 available tools predominantly concern the solar-like oscillators.

67 Software package overview

68 Foam is designed as a customisable pipeline. It will match theoretical models to observations,
69 computing the goodness of fit of each model based on the selected merit function. Afterwards
70 it will determine the best model alongside the uncertainty region of this solution based on
71 statistical criteria. On the observational side, it will take a list of frequencies as an input,
72 optionally complemented by additional information such as a set of surface properties (effective
73 temperature, surface gravity, luminosity, element surface abundances...). On the theoretical
74 side Foam will use a grid of theoretical stellar models, calculated by the user to suit their
75 specific needs. Although the current implementation is made for a grid of stellar equilibrium
76 models computed by MESA (Jermyn et al., 2023; Paxton et al., 2011, 2013, 2015, 2018, 2019),
77 whose pulsation frequencies are computed with GYRE (Townsend et al., 2018; Townsend &
78 Teitler, 2013), the majority of the code is not inherently dependent on MESA. By making
79 certain adjustments to the modelling pipeline, Foam could potentially employ grids generated
80 by different stellar evolution codes. Some suggestions how to approach this are given in
81 the description of the theoretical model grid in the online documentation. However, the
82 implementation of such functionality currently remains out of the scope of the project.

83 The script to run the pipeline can be altered in order to change the modelling approach you want
84 to take. The various configuration options, the installation procedure, and a walkthrough of how
85 to create your own modelling setup, are described in more detail in the online documentation.
86 Although it relies on grids of stellar equilibrium models computed by MESA as the source of
87 the theoretical model grid, MESA's installation itself is not required for Foam to function. The
88 installation of GYRE is however required, specifically since Foam relies on the tar_fit.mX.kX.h5

89 files included in the GYRE installation. This allows us to rescale the g-modes for various
 90 stellar rotation rates, following the traditional approximation of rotation Townsend (2020) and
 91 assuming rigid rotation. This facilitates computing the oscillation frequencies for the grid of
 92 stellar equilibrium models only once, and subsequently rescaling them to find the optimised
 93 rotation rate (see Michielsen et al., 2023). This approach avoids repeating the oscillation
 94 computations for a variety of rotation values, which would introduce extra dimensionality in
 95 the modelling problem in the form of adding the rotation rate as an additional free parameter.

96 Foam's modelling procedure can be broken down into following sequential [steps of the pipeline](#):
 97 - Extract all required parameters and quantities from the files in the theoretical MESA and
 98 GYRE grids. - Construct the theoretical pulsation patterns for each stellar model. Thereafter
 99 select theoretical pulsation patterns matching the observational pattern whilst optimising their
 100 rotation rates. Finally combine this information with the models' surface properties. - Calculate
 101 the likelihood of all the theoretical patterns according to the specified merit functions and
 102 observables. This list of observables consist of the pulsations, but can optionally be extended
 103 with spectroscopic or astrometric information. - Exclude all the models that fall outside an
 104 n-sigma error box on the spectroscopic and astrometric constraints as acceptable solutions. -
 105 Calculate the Akaike information criterion (AIC) (Claeskens & Hjort, 2008) corrected for small
 106 sample size. This statistical criterion rewards goodness of fit, but penalises model complexity in
 107 the form of additional free parameters. The AIC thus allows a statistical comparison between
 108 models of different (nested) grids where the number of free parameters is not the same. -
 109 Calculate the 2 sigma uncertainty region of the maximum likelihood solution using Bayes'
 110 theorem. - Make corner plots for all combinations of the different modelling choices (See
 111 [Figure 1](#) for an example). - Construct a table with the best model of the grid for each
 112 combination of different modelling choices.

113 Next to the tables with the best model parameters and their AIC values, the cornerplots
 114 provide a quick way to assess the output of the pipeline and visualise the modelling results.
 115 [Figure 1](#) shows an example of such a cornerplot for the modelling of KIC 4930889 performed
 116 by Michielsen et al. (2023). It gives a clear indication of which models are included (coloured)
 117 or excluded (greyscale) from the uncertainty region, and indicates what the best models of the
 118 grid are (yellow, see the colour bar).




Figure 1: Cornerplot with the parameters in the grid and the rotation. The 50% best models are shown, colour-coded according to the log of their merit function value. Models in colour fall within the 2 sigma error ellipse, while those in greyscale fall outside of it. Figures on the diagonal show binned parameter distributions of the models in the error ellipse, and the panel at the top right shows an Hertzsprung-Russell (HR) diagram with 1 and 3 sigma observational error boxes. Figure taken from Michielsen et al. (2023).

Acknowledgements

The research leading to the development of this package has received funding from the Research Foundation Flanders (FWO) by means of a PhD scholarship to MM under project No. 11F7120N. MM is grateful to T. Van Reeth for his help concerning the scaling of g-modes with rotation, and to A. Kemp for his suggestions regarding the online documentation.

References

- Aerts, C. (2021). Probing the interior physics of stars through asteroseismology. *Reviews of Modern Physics*, 93(1), 015001. <https://doi.org/10.1103/RevModPhys.93.015001>
- Aerts, C., Christensen-Dalsgaard, J., & Kurtz, D. W. (2010). *Asteroseismology*. Springer, Astronomy; Astrophysics Library.
- Aerts, C., Molenberghs, G., Michielsen, M., Pedersen, M. G., Björklund, R., Johnston, C., Mombarg, J. S. G., Bowman, D. M., Buysschaert, B., Pápics, P. I., Sekaran, S., Sundqvist, J. O., Tkachenko, A., Truyaert, K., Van Reeth, T., & Vermeyen, E. (2018). Forward Asteroseismic Modeling of Stars with a Convective Core from Gravity-mode Oscillations: Parameter Estimation and Stellar Model Selection. *The Astrophysical Journal Supplement Series*, 237, 15. <https://doi.org/10.3847/1538-4365/aaccfb>

- 135 Aguirre Børsen-Koch, V., Rørsted, J. L., Justesen, A. B., Stokholm, A., Verma, K., Winther,
136 M. L., Knudstrup, E., Nielsen, K. B., Sahlholdt, C., Larsen, J. R., Cassisi, S., Serenelli, A.
137 M., Casagrande, L., Christensen-Dalsgaard, J., Davies, G. R., Ferguson, J. W., Lund, M.
138 N., Weiss, A., & White, T. R. (2022). The BAYesian STellar algorithm (BASTA): a fitting
139 tool for stellar studies, asteroseismology, exoplanets, and Galactic archaeology. *509*(3),
140 4344–4364. <https://doi.org/10.1093/mnras/stab2911>
- 141 Anders, E. H., & Pedersen, M. G. (2023). Convective Boundary Mixing in Main-Sequence
142 Stars: Theory and Empirical Constraints. *Galaxies*, *11*(2), 56. <https://doi.org/10.3390/galaxies11020056>
143
- 144 Chontos, A., Huber, D., Sayeed, M., & Yamsiri, P. (2022). pySYD: Automated measurements
145 of global asteroseismic parameters. *The Journal of Open Source Software*, *7*(79), 3331.
146 <https://doi.org/10.21105/joss.03331>
- 147 Claeskens, G., & Hjort, N. L. (2008). *Model Selection and Model Averaging*, *Cambridge Series*
148 *in Statistical and Probabilistic Mathematics*.
- 149 Eckart, G. (1960). Hydrodynamics of oceans and atmospheres. *Hydrodynamics of Oceans and*
150 *Atmospheres*, Pergamon Press, Oxford. <https://doi.org/10.1002/qj.49708938224>
- 151 Jermyn, A. S., Bauer, E. B., Schwab, J., Farmer, R., Ball, W. H., Bellinger, E. P., Dotter, A.,
152 Joyce, M., Marchant, P., Mombarg, J. S. G., Wolf, W. M., Sunny Wong, T. L., Cinquegrana,
153 G. C., Farrell, E., Smolec, R., Thoul, A., Cantiello, M., Herwig, F., Toloza, O., ... Timmes,
154 F. X. (2023). Modules for Experiments in Stellar Astrophysics (MESA): Time-dependent
155 Convection, Energy Conservation, Automatic Differentiation, and Infrastructure. *265*(1),
156 15. <https://doi.org/10.3847/1538-4365/acae8d>
- 157 Michielsen, M., Aerts, C., & Bowman, D. M. (2021). Probing the temperature gradient in
158 the core boundary layer of stars with gravito-inertial modes. The case of KIC 7760680.
159 *Astronomy and Astrophysics*, *650*, A175. <https://doi.org/10.1051/0004-6361/202039926>
- 160 Michielsen, M., Van Reeth, T., Tkachenko, A., & Aerts, C. (2023). Probing the physics in the
161 core boundary layers of the double-lined B-type binary KIC 4930889 from its gravito-inertial
162 modes. *679*, A6. <https://doi.org/10.1051/0004-6361/202244192>
- 163 Paxton, B., Bildsten, L., Dotter, A., Herwig, F., Lesaffre, P., & Timmes, F. (2011). Modules
164 for Experiments in Stellar Astrophysics (MESA). *192*(1), 3. <https://doi.org/10.1088/0067-0049/192/1/3>
165
- 166 Paxton, B., Cantiello, M., Arras, P., Bildsten, L., Brown, E. F., Dotter, A., Mankovich, C.,
167 Montgomery, M. H., Stello, D., Timmes, F. X., & Townsend, R. (2013). Modules for
168 Experiments in Stellar Astrophysics (MESA): Planets, Oscillations, Rotation, and Massive
169 Stars. *208*(1), 4. <https://doi.org/10.1088/0067-0049/208/1/4>
- 170 Paxton, B., Marchant, P., Schwab, J., Bauer, E. B., Bildsten, L., Cantiello, M., Dessart, L.,
171 Farmer, R., Hu, H., Langer, N., Townsend, R. H. D., Townsley, D. M., & Timmes, F. X.
172 (2015). Modules for Experiments in Stellar Astrophysics (MESA): Binaries, Pulsations, and
173 Explosions. *220*(1), 15. <https://doi.org/10.1088/0067-0049/220/1/15>
- 174 Paxton, B., Schwab, J., Bauer, E. B., Bildsten, L., Blinnikov, S., Duffell, P., Farmer, R.,
175 Goldberg, J. A., Marchant, P., Sorokina, E., Thoul, A., Townsend, R. H. D., & Timmes, F.
176 X. (2018). Modules for Experiments in Stellar Astrophysics (MESA): Convective Boundaries,
177 Element Diffusion, and Massive Star Explosions. *234*(2), 34. <https://doi.org/10.3847/1538-4365/aaa5a8>
178
- 179 Paxton, B., Smolec, R., Schwab, J., Gautschi, A., Bildsten, L., Cantiello, M., Dotter, A., Farmer,
180 R., Goldberg, J. A., Jermyn, A. S., Kanbur, S. M., Marchant, P., Thoul, A., Townsend, R.
181 H. D., Wolf, W. M., Zhang, M., & Timmes, F. X. (2019). Modules for Experiments in
182 Stellar Astrophysics (MESA): Pulsating Variable Stars, Rotation, Convective Boundaries,
183 and Energy Conservation. *243*(1), 10. <https://doi.org/10.3847/1538-4365/ab2241>

- 184 Rendle, B. M., Buldgen, G., Miglio, A., Reese, D., Noels, A., Davies, G. R., Campante, T. L.,
185 Chaplin, W. J., Lund, M. N., Kuszewicz, J. S., Scott, L. J. A., Scuflaire, R., Ball, W. H.,
186 Smetana, J., & Nsamba, B. (2019). AIMS - a new tool for stellar parameter determinations
187 using asteroseismic constraints. *484*(1), 771–786. <https://doi.org/10.1093/mnras/stz031>
- 188 Townsend, R. H. D. (2020). Improved asymptotic expressions for the eigenvalues of Laplace's
189 tidal equations. *497*(3), 2670–2679. <https://doi.org/10.1093/mnras/staa2159>
- 190 Townsend, R. H. D., Goldstein, J., & Zweibel, E. G. (2018). Angular momentum transport by
191 heat-driven g-modes in slowly pulsating B stars. *475*, 879–893. <https://doi.org/10.1093/mnras/stx3142>
- 192
- 193 Townsend, R. H. D., & Teitler, S. A. (2013). GYRE: an open-source stellar oscillation code
194 based on a new Magnus Multiple Shooting scheme. *435*, 3406–3418. <https://doi.org/10.1093/mnras/stt1533>
- 195
- 196 Waelkens, C. (1991). Slowly pulsating B stars. *Astronomy and Astrophysics*, *246*, 453.

DRAFT