

Numerik I – Homework 5

Deadline: 29.3.2019, 13:00

Restriction The use of built-in functions for Fourier transform is not allowed.

Exercise 1 (6 P.) (Programming task)

Write a function that implements the Fourier transform given by Definition 5.2 in the lecture notes. The function should take the real-valued vector $(f_k)_{k=0}^{N-1}$ as the input argument and return as the output

- (a) the Fourier coefficients \hat{f}_k ;
- (b) the coefficients \hat{A}_m and \hat{B}_m of the trigonometric polynomial $T(t)$ for real-valued f_k (Equation 5.5 in the lecture notes);
- (c) a plot of the trigonometric polynomial $T(t)$.

Exercise 2 (6 P.) (Programming task)

On the course homepage there is a data file $f.dat$ which stores the evaluations of a continuous function $f : [0, 2x_0] \rightarrow \mathbb{R}$

$$f_k = f\left(\frac{2x_0 k}{N}\right) \quad \text{where } k = 0, 1, \dots, N-1$$

for $N = 2^{16}$. The function has the property $f(0) = f(2x_0)$, $x_0 = 101.15520937701599$.

For the refinement level $l = 2, 4, 6, 8, 10, 12, 14, 16$,

- (a) Use your program in Exercise 1 to calculate the trigonometric polynomial $T(t)$ by using the following data sets respectively as the input argument

$$f\left(\frac{2x_0 k}{2^l}\right) \quad \text{where } k = 0, 1, \dots, 2^l - 1;$$

and show the trigonometric polynomials in the plot.

- (b) Calculate the approximation error for each data set

$$\max_{0 \leq k \leq N-1} \left| T_l\left(\frac{2x_0 k}{N}\right) - f_k \right|$$

and record them in a table.