

Numerik I – Assignment 2

Deadline: 08.03.2018, 12:00

Please note that the use of Matlab internal functions like `polyfit` or `polyval` is not permitted to solve these exercises.

Exercise 1

- a) Write a Matlab function that computes the Newton differences as in formula (2.7) of the lecture notes. Input: nodes $(x_i)_{i=0}^n$ as vector, function values $(f_i)_{i=0}^n$ as vector. Output: Newton differences $(b_i)_{i=0}^n$ as vector. (3 points)
- b) Write a Matlab function that evaluates the interpolation polynomial defined via the nodes $(x_i)_{i=0}^n$ and Newton differences $(b_i)_{i=0}^n$ at a given point x (Satz 2.5 in the lecture notes).
Verify the functionality of your programs by interpolating the function $f(x) = x(x-1)$ at $x_0 = 0$, $x_1 = 1$ and $x_2 = 2$. Evaluate the interpolation at $x = 0.5$. What do you expect as result? (2 points)

Please use the code that you developed above to solve the following exercises.

Exercise 2

Let the function

$$f(x) := \frac{x}{1 + \cos(x)^2},$$

and intervals $I_1 := [-2, 2]$ and $I_2 := [-4, 4]$ be given.

- a) Construct the Newton polynomials of degree $n = 5, 10, 15$ on $n + 1$ equispaced nodes in I_1 (the nodes should include the endpoints of the interval). Plot the polynomials and f in the same graph. (2 points)
- b) Construct the Newton polynomials of degree $n = 5, 10, 15$ on $n + 1$ equispaced nodes in I_2 . Plot the polynomials and f in the same graph. (1 point)
- c) Explain the behaviour you observed in b). Use error estimate (2.13) from the lecture notes in your reasoning. (2 points)

- d) Construct the Newton polynomials P_n of degree $n = 5, 10, 15, 20, 25$ on $n+1$ Chebyshev nodes in I_2 . Plot the approximations and f in the same graph. Estimate the maximum norm

$$\|f - P_n\|_{\max}, \quad n = 5, 10, 15, 20, 25$$

by taking the maximum of the absolute value of the difference of f and P_n on 1000 equispaced nodes on I_2 . Display the errors in a table. What do you observe and why? (3 points)

Exercise 3

Let $g(x) := \sin(2\pi x)$.

- a) Evaluate the function g at 21 equispaced nodes in $[-1, 1]$ (the nodes should contain the endpoints -1 and 1). Compute the Newton polynomial P of degree 20. Plot P and f in the same graph. (1 point)
- b) Load the file ‘measurements.dat’ (use the Matlab command `load`). It contains 21 numbers that can be interpreted as function values of an unknown function \bar{g} at the 21 equispaced nodes in $[-1, 1]$ from a). Approximate \bar{g} by computing the interpolating polynomial and plot this approximation and g in the same graph. Knowing a posteriori that the set of ‘measured’ data came from the function $\bar{g}(x) = \sin(2\pi x) + (-1)^{i+1}10^{-4}$, comment on the result of your plot. (2 points)