

# Machine learning from scratch

## Lecture 2: Convex optimization

Alexis Zubiolo

`alexis.zubiolo@gmail.com`

Data Science Team Lead @ Adcash

February 2, 2017

## Context reminder

living area (m <sup>2</sup> )	# bedrooms	intercept	price (1000's BGN)
50	1	1	30
76	2	1	48
26	1	1	12
102	3	1	90

$$h(\mathbf{x}) = \sum_{j=0}^d \theta_j x_j = \boldsymbol{\theta}^T \mathbf{x}$$

## Context reminder

living area (m <sup>2</sup> )	# bedrooms	intercept	price (1000's BGN)
50	1	1	30
76	2	1	48
26	1	1	12
102	3	1	90

$$h(\mathbf{x}) = \sum_{j=0}^d \theta_j x_j = \boldsymbol{\theta}^T \mathbf{x}$$

Suppose we chose the following loss function:

$$\ell(y, \hat{y}) = \frac{1}{2} (y - \hat{y})^2$$

This leads to the following least squares *cost function*:

$$J(\boldsymbol{\theta}) = \frac{1}{2} \sum_{i=1}^n \left( h(\mathbf{x}^{(i)}) - y^{(i)} \right)^2$$

This problem the **ordinary least squares** (OLS) regression model.

Thank you! Questions?

`alexis.zubiollo@gmail.com`

`https://github.com/azubiollo/itstep`