

# Proof of Regularity of 1-, 2-, and 3-Sums of Matroids

Ivan Sergeev

March– 2025

## 1 Preliminaries

### 1.1 Total Unimodularity and Partial Unimodularity

**Definition 1.** We say that a matrix  $A$  is totally unimodular, or TU for short, if for every  $k \in \mathbb{Z}_{\geq 1}$ , every  $k \times k$  submatrix  $T$  of  $A$  has  $\det T \in \{0, \pm 1\}$ .

**Definition 2.** Given  $k \in \mathbb{Z}_{\geq 1}$ , we say that a matrix  $A$  is  $k$ -partially unimodular, or  $k$ -PU for short, if every  $k \times k$  submatrix  $T$  of  $A$  has  $\det T \in \{0, \pm 1\}$ .

**Lemma 3.** A matrix  $A$  is TU if and only if  $A$  is  $k$ -PU for every  $k \in \mathbb{Z}_{\geq 1}$ .

*Proof.* This follows from Definitions 1 and 2. □

### 1.2 Pivoting

#### 1.2.1 Definitions

long tableau pivot, short tableau pivot

#### 1.2.2 Properties

**Lemma 4.** Let  $B = \begin{bmatrix} B_{11} & 0 \\ B_{21} & B_{22} \end{bmatrix} \in \mathbb{Q}^{\{X_1 \cup X_2\} \times \{Y_1 \times Y_2\}}$ . Let  $B' = \begin{bmatrix} B'_{11} & B'_{12} \\ B'_{21} & B'_{22} \end{bmatrix}$  be the result of performing a short tableau pivot on  $(x, y) \in X_1 \times Y_1$  in  $B$ . Then  $B'_{12} = 0$ ,  $B'_{22} = B_{22}$ , and  $\begin{bmatrix} B'_{11} \\ B'_{21} \end{bmatrix}$  is the matrix resulting from performing a short tableau pivot on  $(x, y)$  in  $\begin{bmatrix} B_{11} \\ B_{21} \end{bmatrix}$ .

*Proof.* This follows by a direct calculation. Indeed, because of the 0 block in  $B$ ,  $B_{12}$  and  $B_{22}$  remain unchanged, and since  $\begin{bmatrix} B_{11} \\ B_{21} \end{bmatrix}$  is a submatrix of  $B$  containing the pivot element, performing a short tableau pivot in it is equivalent to performing a short tableau pivot in  $B$  and then taking the corresponding submatrix. □

**Lemma 5.** Let  $k \in \mathbb{Z}_{\geq 1}$ , let  $A \in \mathbb{Q}^{k \times k}$ , and let  $A'$  be the result of performing a short tableau pivot in  $A$  on  $A(x, y) \neq 0$  where  $x, y \in \{1, \dots, k\}$ . Then  $A'$  contains a submatrix  $A''$  of size  $(k-1) \times (k-1)$  with  $|\det A''| = |\det A|/|A(x, y)|$ .

*Proof.* Let  $X = \{1, \dots, k\} \setminus \{x\}$  and  $Y = \{1, \dots, k\} \setminus \{y\}$ , and let  $A'' = A'(X, Y)$ . Since  $A''$  does not contain the pivot row or the pivot column,  $\forall (i, j) \in X \times Y$  we have  $A''(i, j) = A(i, j) - \frac{A(i, y) \cdot A(x, j)}{A(x, y)}$ . For  $\forall j \in Y$ , let  $B_j$  be the matrix obtained from  $A$  by removing row  $x$  and column  $j$ , and let  $B'_j$  be the matrix obtained from  $A''$  by replacing column  $j$  with  $A(X, y)$  (i.e., the pivot column without the pivot element). The cofactor expansion along row  $x$  in  $A$  yields

$$\det A = \sum_{j=1}^k (-1)^{y+j} \cdot A(x, j) \cdot \det B_j.$$

By reordering columns of every  $B_j$  to match their order in  $B_j''$ , we get

$$\det A = (-1)^{x+y} \cdot \left( A(x, y) \cdot \det A' - \sum_{j \in Y} A(x, j) \cdot \det B_j'' \right).$$

By linearity of the determinant applied to  $\det A''$ , we have

$$\det A'' = \det A' - \sum_{j \in Y} \frac{A(x, j)}{A(x, y)} \cdot \det B_j''$$

Therefore,  $|\det A''| = |\det A|/|A(x, y)|$ . □

**Lemma 6.** Let  $k \in \mathbb{Z}_{\geq 1}$ , let  $A \in \mathbb{Q}^{k \times k}$ , and let  $A'$  be the result of performing a short tableau pivot in  $A$  on  $A(x, y) \in \{\pm 1\}$  where  $x, y \in \{1, \dots, k\}$ . Then  $A'$  contains a submatrix  $A''$  of size  $(k-1) \times (k-1)$  with  $|\det A''| = |\det A|$ .

*Proof.* Apply Lemma 5 to  $A$  and use that  $A(x, y) \in \{\pm 1\}$ . □

### 1.3 Vector Matroids

#### 1.3.1 Full Matrix Representation

definition, properties

#### 1.3.2 Standard Matrix Representation

definition, properties

#### 1.3.3 Conversion From Full to Standard Matrix Representation

match lean implementation

**Lemma 7.** Let  $M$  be a matroid represented by a matrix  $A \in \mathbb{Q}^{X \times Y}$  and let  $B$  be a base of  $M$ . Then there exists a matrix  $S \in \mathbb{Q}^{B \times (Y \setminus B)}$  that is a standard representation matrix of  $M$ .

*Proof.* Let  $C = \{A(\bullet, b) \mid b \in B\}$ . Since  $B$  is a base of  $M$ , we can show that  $C$  is a basis in the column space  $\text{span}\{A(\bullet, y) \mid y \in Y\}$ . For every  $y \in Y \setminus B$ , let  $S(\bullet, y)$  be the coordinates of  $A(\bullet, y)$  in basis  $C$ . We can show that  $[I \mid S]$  represents the same matroid as  $A$ , so  $S$  is a standard representation matrix of  $M$ . □

**Lemma 8.** Let  $M$  be a matroid represented by a TU matrix  $A \in \mathbb{Q}^{X \times Y}$  and let  $B$  be a base of  $M$ . Then there exists a matrix  $S \in \mathbb{Q}^{B \times (Y \setminus B)}$  such that  $S$  is TU and  $S$  is a standard representation matrix of  $M$ .

*Proof sketch.* Apply the procedure described in the proof of Lemma 7 to  $A$ . This procedure can be represented as a sequence of elementary row operations, all of which preserve TUness. Dropping the identity matrix at the end also preserves TUness.

write up new proof using general pivoting

□

see details in implementation

#### 1.3.4 Support Matrices

**Definition 9.** Let  $F$  be a field. The support of matrix  $A \in F^{X \times Y}$  is  $A^\# \in \{0, 1\}^{X \times Y}$  given by

$$\forall i \in X, \forall j \in Y, A^\#(i, j) = \begin{cases} 0, & \text{if } A(i, j) = 0, \\ 1, & \text{if } A(i, j) \neq 0. \end{cases}$$

**Definition 10.** Let  $M$  be a matroid, let  $B$  be a base of  $M$ , and let  $e \in E \setminus B$  be an element. The fundamental circuit  $C(e, B)$  of  $e$  with respect to  $B$  is the unique circuit contained in  $B \cup \{e\}$ .

**Lemma 11.** Let  $M$  be a matroid and let  $S \in F^{X \times Y}$  be a standard representation matrix of  $M$  over a field  $F$ . Then  $\forall y \in Y$ , the fundamental circuit of  $y$  w.r.t.  $X$  is  $C(y, X) = \{y\} \cup \{x \in X \mid S(x, y) \neq 0\}$ .

*Proof.* Let  $y \in Y$ . Our goal is to show that  $C'(y, X) = \{y\} \cup \{x \in X \mid D(x, y) \neq 0\}$  is a fundamental circuit of  $y$  with respect to  $X$ .

- $C'(y, X) \subseteq X \cup \{y\}$  by construction.
- $C'(y, X)$  is dependent, since columns of  $[I \mid S]$  indexed by elements of  $C(y, X)$  are linearly dependent.
- If  $C \subsetneq C'(y, X)$ , then  $C$  is independent. To show this, let  $V$  be the set of columns of  $[I \mid S]$  indexed by elements of  $C$  and consider two cases.
  1. Suppose that  $y \notin C$ . Then vectors in  $V$  are linearly independent (as columns of  $I$ ). Thus,  $C$  is independent.
  2. Suppose  $\exists x \in X \setminus C$  such that  $S(x, y) \neq 0$ . Then any nontrivial linear combination of vectors in  $V$  has a non-zero entry in row  $x$ . Thus, these vectors are linearly independent, so  $C$  is independent.

□

**Lemma 12.** Let  $M$  be a matroid and let  $S \in F^{X \times Y}$  be a standard representation matrix of  $M$  over a field  $F$ . Then  $\forall y \in Y$ , column  $S^\#(\bullet, y)$  is the characteristic vector of  $C(y, X) \setminus \{y\}$ .

*Proof.* This directly follows from Lemma 11. □

**Lemma 13.** Let  $A$  be a TU matrix.

1. If a matroid is represented by  $A$ , then it is also represented by  $A^\#$ .
2. If a matroid is represented by  $A^\#$ , then it is also represented by  $A$ .

*Proof.* See Lean implementation.

add details

□

## 1.4 Regular Matroids

**Definition 14.** A matroid  $M$  is regular if there exists  $A \in \mathbb{Q}^{X \times Y}$  such that  $M = M[A]$  and  $A$  is TU.

**Definition 15.** We say that  $A' \in \mathbb{Q}^{X \times Y}$  is a TU signing of  $A \in \mathbb{Z}_2^{X \times Y}$  if  $A'$  is TU and

$$\forall i \in X, \forall j \in Y, |A'(i, j)| = A(i, j).$$

**Lemma 16.** Let  $M$  be a matroid given by a standard representation matrix  $B \in \mathbb{Z}_2^{X \times Y}$ . Then  $M$  is regular if and only if  $B$  has a TU signing.

*Proof.* Suppose that  $M$  is regular. By Definition 14, there exists  $A \in \mathbb{Q}^{X \times Y}$  such that  $M = M[A]$  and  $A$  is TU. Recall that  $X$  (the row set of  $B$ ) is a base of  $M$ . By Lemma 8,  $A$  can be converted into a standard representation matrix  $B' \in \mathbb{Q}^{X \times Y}$  of  $M$  such that  $B'$  is also TU. Since  $B'$  and  $B$  are both standard representations of  $M$ , by Lemma 12 the support matrices  $(B')^\#$  and  $B^\#$  are the same. Moreover,  $B^\# = B$ , since  $B$  has entries in  $\mathbb{Z}_2$ . Thus,  $B'$  is TU and  $(B')^\# = B$ , so  $B'$  is a TU signing of  $B$ .

Suppose that  $B$  has a TU signing  $B' \in \mathbb{Q}^{X \times Y}$ . Then  $A = [I \mid B']$  is TU, as it is obtained from  $B'$  by adjoining the identity matrix. Moreover, by Lemma 13,  $A$  represents the same matroid as  $A^\# = [I \mid B]$ , which is  $M$ . Thus,  $A$  is a TU matrix representing  $M$ , so  $M$  is regular. □

add lemma

## 2 Regularity of 1-Sum

Write up based on Lean implementation

### 3 Regularity of 2-Sum

**Definition 17.** Let  $R$  be a semiring (we will use  $R = \mathbb{Z}_2$  and  $R = \mathbb{Q}$ ). Let  $B_\ell \in R^{(X_\ell \cup \{x\}) \times Y_\ell}$  and  $B_r \in R^{X_r \times (Y_r \cup \{y\})}$  be matrices of the form

$$B_\ell = \begin{bmatrix} A_\ell \\ r \end{bmatrix}, \quad B_r = \begin{bmatrix} c & A_r \end{bmatrix}.$$

The 2-sum  $B = B_\ell \oplus_{2,x,y} B_r$  of  $B_\ell$  and  $B_r$  is defined as

$$B = \begin{bmatrix} A_\ell & 0 \\ D & A_r \end{bmatrix} \quad \text{where} \quad D = c \otimes r.$$

Here  $A_\ell \in R^{X_\ell \times Y_\ell}$ ,  $A_r \in R^{X_r \times Y_r}$ ,  $r \in R^{Y_\ell}$ ,  $c \in R^{X_r}$ ,  $D \in R^{X_\ell \times Y_r}$ , and the indexing is consistent everywhere.

**Definition 18.** A matroid  $M$  is a 2-sum of matroids  $M_\ell$  and  $M_r$  if there exist standard  $\mathbb{Z}_2$  representation matrices  $B$ ,  $B_\ell$ , and  $B_r$  (for  $M$ ,  $M_\ell$ , and  $M_r$ , respectively) of the form given in Definition 17.

**Lemma 19.** Let  $B_\ell$  and  $B_r$  from Definition 17 be TU matrices (over  $\mathbb{Q}$ ). Then  $C = \begin{bmatrix} D & A_r \end{bmatrix}$  is TU.

*Proof.* Since  $B_\ell$  is TU, all its entries are in  $\{0, \pm 1\}$ . In particular,  $r$  is a  $\{0, \pm 1\}$  vector. Therefore, every column of  $D$  is a copy of  $y$ ,  $-y$ , or the zero column. Thus,  $C$  can be obtained from  $B_r$  by adjoining zero columns, duplicating the  $y$  column, and multiplying some columns by  $-1$ . Since all these operations preserve TUness and since  $B_r$  is TU,  $C$  is also TU.  $\square$

**Lemma 20.** Let  $B_\ell$  and  $B_r$  be matrices from Definition 17. Let  $B'_\ell$  and  $B'$  be the matrices obtained by performing a short tableau pivot on  $(x_\ell, y_\ell) \in X_\ell \times Y_\ell$  in  $B_\ell$  and  $B$ , respectively. Then  $B' = B'_\ell \oplus_{2,x,y} B_r$ .

*Proof.* Let

$$B'_\ell = \begin{bmatrix} A'_\ell \\ r' \end{bmatrix}, \quad B' = \begin{bmatrix} B'_{11} & B'_{12} \\ B'_{21} & B'_{22} \end{bmatrix}$$

where the blocks have the same dimensions as in  $B_\ell$  and  $B$ , respectively. By Lemma 4,  $B'_{11} = A'_\ell$ ,  $B'_{12} = 0$ , and  $B'_{22} = A_r$ . Equality  $B'_{21} = c \otimes r'$  can be verified via a direct calculation. Thus,  $B' = B'_\ell \oplus_{2,x,y} B_r$ .  $\square$

**Lemma 21.** Let  $B_\ell$  and  $B_r$  from Definition 17 be TU matrices (over  $\mathbb{Q}$ ). Then  $B_\ell \oplus_{2,x,y} B_r$  is TU.

*Proof.* By Lemma 3, it suffices to show that  $B_\ell \oplus_{2,x,y} B_r$  is  $k$ -PU for every  $k \in \mathbb{Z}_{\geq 1}$ . We prove this claim by induction on  $k$ . The base case with  $k = 1$  holds, since all entries of  $B_\ell \oplus_{2,x,y} B_r$  are in  $\{0, \pm 1\}$  by construction.

Suppose that for some  $k \in \mathbb{Z}_{\geq 1}$  we know that for any TU matrices  $B'_\ell$  and  $B'_r$  (from Definition 17) their 2-sum  $B'_\ell \oplus_{2,x,y} B'_r$  is  $k$ -PU. Now, given TU matrices  $B_\ell$  and  $B_r$  (from Definition 17), our goal is to show that  $B = B_\ell \oplus_{2,x,y} B_r$  is  $(k+1)$ -PU, i.e., that every  $(k+1) \times (k+1)$  submatrix  $T$  of  $B$  has  $\det T \in \{0, \pm 1\}$ .

First, suppose that  $T$  has no rows in  $X_\ell$ . Then  $T$  is a submatrix of  $\begin{bmatrix} D & A_r \end{bmatrix}$ , which is TU by Lemma 19, so  $\det T \in \{0, \pm 1\}$ . Thus, we may assume that  $T$  contains a row  $x_\ell \in X_\ell$ .

Next, note that without loss of generality we may assume that there exists  $y_\ell \in Y_\ell$  such that  $T(x_\ell, y_\ell) \neq 0$ . Indeed, if  $T(x_\ell, y) = 0$  for all  $y$ , then  $\det T = 0$  and we are done, and  $T(x_\ell, y) = 0$  holds whenever  $y \in Y_r$ .

Since  $B$  is 1-PU, all entries of  $T$  are in  $\{0, \pm 1\}$ , and hence  $T(x_\ell, y_\ell) \in \{\pm 1\}$ . Thus, by Lemma 6, performing a short tableau pivot in  $T$  on  $(x_\ell, y_\ell)$  yields a matrix that contains a  $k \times k$  submatrix  $T''$  such that  $|\det T| = |\det T''|$ . Since  $T$  is a submatrix of  $B$ , matrix  $T''$  is a submatrix of the matrix  $B'$  resulting from performing a short tableau pivot in  $B$  on the same entry  $(x_\ell, y_\ell)$ . By Lemma 20, we have  $B' = B'_\ell \oplus_{2,x,y} B_r$  where  $B'_\ell$  is the result of performing a short tableau pivot in  $B_\ell$  on  $(x_\ell, y_\ell)$ . Since TUness is preserved by pivoting and  $B_\ell$  is TU,  $B'_\ell$  is also TU. Thus, by the inductive hypothesis applied to  $T''$  and  $B'_\ell \oplus_{2,x,y} B_r$ , we have  $\det T'' \in \{0, \pm 1\}$ . Since  $|\det T| = |\det T''|$ , we conclude that  $\det T \in \{0, \pm 1\}$ .  $\square$

**Lemma 22.** Let  $M$  be a 2-sum of regular matroids  $M_\ell$  and  $M_r$ . Then  $M$  is also regular.

*Proof.* Let  $B$ ,  $B_\ell$ , and  $B_r$  be standard  $\mathbb{Z}_2$  representation matrices from Definition 18. Since  $M_\ell$  and  $M_r$  are regular, by Lemma 16,  $B_\ell$  and  $B_r$  have TU signings  $B'_\ell$  and  $B'_r$ , respectively. Then  $B' = B'_\ell \oplus_{2,x,y} B'_r$  is a TU signing of  $B$ . Indeed,  $B'$  is TU by Lemma 21, and a direct calculation verifies that  $B'$  is a signing of  $B$ . Thus,  $M$  is regular by Lemma 16.  $\square$

## 4 Regularity of 3-Sum

### 4.1 Definition

**Definition 23.** Let  $B_l \in \mathbb{Z}_2^{(X_l \cup \{x_0, x_1\}) \times (Y_l \cup \{y_2\})}$ ,  $B_r \in \mathbb{Z}_2^{(X_r \cup \{x_2\}) \times (Y_r \cup \{y_0, y_1\})}$  be matrices of the form

$$B_l = \begin{array}{|c|c|c|} \hline & A_l & 0 \\ \hline & 1 & 1 \\ \hline D_l & D_0 & \begin{array}{|c|} \hline 1 \\ \hline 1 \end{array} \\ \hline \end{array} \quad \text{and} \quad B_r = \begin{array}{|c|c|c|c|} \hline 1 & 1 & 0 & 0 \\ \hline D_0 & \begin{array}{|c|} \hline 1 \\ \hline 1 \end{array} & & A_r \\ \hline D_r & & & \\ \hline \end{array} \quad \text{where} \quad D_0 = \begin{array}{|c|c|} \hline 1 & 0 \\ \hline 0 & 1 \end{array} \quad \text{or} \quad D_0 = \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 0 & 1 \end{array}.$$

The 3-sum  $B = B_l \oplus_3 B_r \in \mathbb{Z}_2^{(X_l \cup X_r) \times (Y_l \cup Y_r)}$  of  $B_l$  and  $B_r$  is defined as

$$B = \begin{array}{|c|c|c|} \hline & A_l & 0 \\ \hline & 1 & 1 \\ \hline D_l & D_0 & \begin{array}{|c|} \hline 1 \\ \hline 1 \end{array} \\ \hline D_{lr} & D_r & A_r \\ \hline \end{array} \quad \text{where} \quad D_{lr} = D_r \cdot (D_0)^{-1} \cdot D_l.$$

Here  $x_2 \in X_l$ ,  $x_0, x_1 \in X_r$ ,  $y_0, y_1 \in Y_l$ ,  $y_2 \in Y_r$ ,  $A_l \in \mathbb{Z}_2^{X_l \times Y_l}$ ,  $A_r \in \mathbb{Z}_2^{X_r \times Y_r}$ ,  $D_l \in \mathbb{Z}_2^{\{x_0, x_1\} \times (Y_l \setminus \{y_0, y_1\})}$ ,  $D_r \in \mathbb{Z}_2^{(X_r \setminus \{x_0, x_1\}) \times \{y_0, y_1\}}$ ,  $D_{lr} \in \mathbb{Z}_2^{(X_r \setminus \{x_0, x_1\}) \times (Y_l \setminus \{y_0, y_1\})}$ ,  $D_0 \in \mathbb{Z}_2^{\{x_0, x_1\} \times \{y_0, y_1\}}$ . The indexing is consistent everywhere.

**Remark 24.** In Definition 23,  $D_0$  is non-singular by construction, so  $D_{lr}$  and  $B$  are well-defined. Moreover, a non-singular  $\mathbb{Z}_2^{2 \times 2}$  matrix is either  $\begin{array}{|c|c|} \hline 1 & 0 \\ \hline 0 & 1 \end{array}$  or  $\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 0 & 1 \end{array}$  up to re-indexing. Thus, Definition 23 can be equivalently restated with  $D_0$  required to be non-singular and  $B_l$ ,  $B_r$ , and  $B$  re-indexed appropriately.

**Definition 25.** A matroid  $M$  is a 3-sum of matroids  $M_\ell$  and  $M_r$  if there exist standard  $\mathbb{Z}_2$  representation matrices  $B$ ,  $B_\ell$ , and  $B_r$  (for  $M$ ,  $M_\ell$ , and  $M_r$ , respectively) of the form given in Definition 23.

### 4.2 Canonical Signing

**Definition 26.** We call  $D'_0 \in \mathbb{Q}^{\{x_0, x_1\} \times \{y_0, y_1\}}$  the canonical signing of  $D_0 \in \mathbb{Z}_2^{\{x_0, x_1\} \times \{y_0, y_1\}}$  if

$$D_0 = \begin{array}{|c|c|} \hline 1 & 0 \\ \hline 0 & 1 \end{array} \quad \text{and} \quad D'_0 = \begin{array}{|c|c|} \hline 1 & 0 \\ \hline 0 & -1 \end{array}, \quad \text{or} \quad D_0 = \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 0 & 1 \end{array} \quad \text{and} \quad D'_0 = \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 0 & 1 \end{array}.$$

Similarly, we call  $S' \in \mathbb{Q}^{\{x_0, x_1, x_2\} \times \{y_0, y_1, y_2\}}$  the canonical signing of  $S \in \mathbb{Z}_2^{\{x_0, x_1, x_2\} \times \{y_0, y_1, y_2\}}$  if

$$S = \begin{array}{|c|c|c|} \hline 1 & 1 & 0 \\ \hline D_0 & \begin{array}{|c|} \hline 1 \\ \hline 1 \end{array} & \\ \hline \end{array} \quad \text{and} \quad S' = \begin{array}{|c|c|c|} \hline 1 & 1 & 0 \\ \hline D'_0 & \begin{array}{|c|} \hline 1 \\ \hline 1 \end{array} & \\ \hline \end{array}.$$

To simplify notation, going forward we use  $D_0$ ,  $D'_0$ ,  $S$ , and  $S'$  to refer to the matrices of the form above.

**Lemma 27.** The canonical signing  $S'$  of  $S$  (from Definition 26) is TU.

*Proof.* Verified via a direct calculation. □

**Lemma 28.** Let  $Q$  be a TU signing of  $S$  (from Definition 26). Let  $u \in \{0, \pm 1\}^{\{x_0, x_1, x_2\}}$ ,  $v \in \{0, \pm 1\}^{\{y_0, y_1, y_2\}}$ , and  $Q'$  be defined as follows:

$$u(i) = \begin{cases} Q(x_2, y_0) \cdot Q(x_0, y_0), & i = x_0, \\ Q(x_2, y_0) \cdot Q(x_0, y_0) \cdot Q(x_0, y_2) \cdot Q(x_1, y_2), & i = x_1, \\ 1, & i = x_2, \end{cases}$$

$$v(j) = \begin{cases} Q(x_2, y_0), & j = y_0, \\ Q(x_2, y_1), & j = y_1, \\ Q(x_2, y_0) \cdot Q(x_0, y_0) \cdot Q(x_0, y_2), & j = y_2, \end{cases}$$

$$Q'(i, j) = Q(i, j) \cdot u(i) \cdot v(j) \quad \forall i \in \{x_0, x_1, x_2\}, \forall j \in \{y_0, y_1, y_2\}.$$

Then  $Q' = S'$  (from Definition 26).

*Proof.* Since  $Q$  is a TU signing of  $S$  and  $Q'$  is obtained from  $Q$  by multiplying rows and columns by  $\pm 1$  factors,  $Q'$  is also a TU signing of  $S$ . By construction, we have

$$\begin{aligned} Q'(x_2, y_0) &= Q(x_2, y_0) \cdot 1 \cdot Q(x_2, y_0) = 1, \\ Q'(x_2, y_1) &= Q(x_2, y_1) \cdot 1 \cdot Q(x_2, y_1) = 1, \\ Q'(x_2, y_2) &= 0, \\ Q'(x_0, y_0) &= Q(x_0, y_0) \cdot (Q(x_2, y_0) \cdot Q(x_0, y_0)) \cdot Q(x_2, y_0) = 1, \\ Q'(x_0, y_1) &= Q(x_0, y_1) \cdot (Q(x_2, y_0) \cdot Q(x_0, y_0)) \cdot Q(x_2, y_1), \\ Q'(x_0, y_2) &= Q(x_0, y_2) \cdot (Q(x_2, y_0) \cdot Q(x_0, y_0)) \cdot (Q(x_2, y_0) \cdot Q(x_0, y_0) \cdot Q(x_0, y_2)) = 1, \\ Q'(x_1, y_0) &= 0, \\ Q'(x_1, y_1) &= Q(x_1, y_1) \cdot (Q(x_2, y_0) \cdot Q(x_0, y_0) \cdot Q(x_0, y_2) \cdot Q(x_1, y_2)) \cdot (Q(x_2, y_1)), \\ Q'(x_1, y_2) &= Q(x_1, y_2) \cdot (Q(x_2, y_0) \cdot Q(x_0, y_0) \cdot Q(x_0, y_2) \cdot Q(x_1, y_2)) \cdot (Q(x_2, y_0) \cdot Q(x_0, y_0) \cdot Q(x_0, y_2)) = 1. \end{aligned}$$

Thus, it remains to show that  $Q'(x_0, y_1) = S'(x_0, y_1)$  and  $Q'(x_1, y_1) = S'(x_1, y_1)$ .

Consider the entry  $Q'(x_0, y_1)$ . If  $D_0(x_0, y_1) = 0$ , then  $Q'(x_0, y_1) = 0 = S'(x_0, y_1)$ . Otherwise, we have  $D_0(x_0, y_1) = 1$ , and so  $Q'(x_0, y_1) \in \{\pm 1\}$ , as  $Q'$  is a signing of  $S$ . If  $Q'(x_0, y_1) = -1$ , then

$$\det Q'(\{x_0, x_2\}, \{y_0, y_1\}) = \det \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} = 2 \notin \{0, \pm 1\},$$

which contradicts TUness of  $Q'$ . Thus,  $Q'(x_0, y_1) = 1 = S'(x_0, y_1)$ .

Consider the entry  $Q'(x_1, y_1)$ . Since  $Q'$  is a signing of  $S$ , we have  $Q'(x_1, y_1) \in \{\pm 1\}$ . Consider two cases.

1. Suppose that  $D_0 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ . If  $Q'(x_1, y_1) = 1$ , then  $\det Q = \det \begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix} = -2 \notin \{0, \pm 1\}$ , which contradicts TUness of  $Q'$ . Thus,  $Q'(x_1, y_1) = -1 = S'(x_1, y_1)$ .
2. Suppose that  $D_0 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ . If  $Q'(x_1, y_1) = -1$ , then  $\det Q(\{x_0, x_1\}, \{y_1, y_2\}) = \det \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} = 2 \notin \{0, \pm 1\}$ , which contradicts TUness of  $Q'$ . Thus,  $Q'(x_1, y_1) = 1 = S'(x_1, y_1)$ .

□

**Definition 29.** Let  $X$  and  $Y$  be sets with  $\{x_0, x_1, x_2\} \subseteq X$  and  $\{y_0, y_1, y_2\} \subseteq Y$ . Let  $Q \in \mathbb{Q}^{X \times Y}$  be a TU

matrix. Define  $u \in \{0, \pm 1\}^X$ ,  $v \in \{0, \pm 1\}^Y$ , and  $Q'$  as follows:

$$u(i) = \begin{cases} Q(x_2, y_0) \cdot Q(x_0, y_0), & i = x_0, \\ Q(x_2, y_0) \cdot Q(x_0, y_0) \cdot Q(x_0, y_2) \cdot Q(x_1, y_2), & i = x_1, \\ 1, & i = x_2, \\ 1, & i \in X \setminus \{x_0, x_1, x_2\}, \end{cases}$$

$$v(j) = \begin{cases} Q(x_2, y_0), & j = y_0, \\ Q(x_2, y_1), & j = y_1, \\ Q(x_2, y_0) \cdot Q(x_0, y_0) \cdot Q(x_0, y_2), & j = y_2, \\ 1, & j \in Y \setminus \{y_0, y_1, y_2\}, \end{cases}$$

$$Q'(i, j) = Q(i, j) \cdot u(i) \cdot v(j) \quad \forall i \in X, \forall j \in Y.$$

We call  $Q'$  the canonical re-signing of  $Q$ .

**Lemma 30.** Let  $X$  and  $Y$  be sets with  $\{x_0, x_1, x_2\} \subseteq X$  and  $\{y_0, y_1, y_2\} \subseteq Y$ . Let  $Q \in \mathbb{Q}^{X \times Y}$  be a TU signing of  $Q_0 \in \mathbb{Z}_2^{X \times Y}$  such that  $Q_0(\{x_0, x_1, x_2\}, \{y_0, y_1, y_2\}) = S$  (from Definition 26). Then the canonical re-signing  $Q'$  of  $Q$  is a TU signing of  $Q_0$  and  $Q'(\{x_0, x_1, x_2\}, \{y_0, y_1, y_2\}) = S'$  (from Definition 26).

*Proof.* Since  $Q$  is a TU signing of  $Q_0$  and  $Q'$  is obtained from  $Q$  by multiplying some rows and columns by  $\pm 1$  factors,  $Q'$  is also a TU signing of  $Q_0$ . Equality  $Q'(\{x_0, x_1, x_2\}, \{y_0, y_1, y_2\}) = S'$  follows from Lemma 28.  $\square$

**Definition 31.** Suppose that  $B_l$  and  $B_r$  from Definition 23 have TU signings  $B'_l$  and  $B'_r$ , respectively. Let  $B''_l$  and  $B''_r$  be the canonical re-signings (from Definition 29) of  $B'_l$  and  $B'_r$ , respectively. Let  $A''_l, A''_r, D''_l, D''_r$ , and  $D''_0$  be blocks of  $B''_l$  and  $B''_r$  analogous to blocks  $A_l, A_r, D_l, D_r$ , and  $D_0$  of  $B_l$  and  $B_r$ . The canonical signing  $B''$  of  $B$  is defined as

$$B'' = \begin{array}{|c|c|c|c|} \hline & A''_l & & 0 \\ \hline & \begin{array}{|c|c|c|} \hline 1 & 1 & 0 \\ \hline \end{array} & & \\ \hline D''_l & D''_0 & \begin{array}{|c|} \hline 1 \\ \hline 1 \\ \hline \end{array} & \\ \hline D''_{lr} & D''_r & & A''_r \\ \hline \end{array} \quad \text{where} \quad D''_{lr} = D''_r \cdot (D''_0)^{-1} \cdot D''_l.$$

**Remark 32.** In Definition 31,  $D''_0$  is non-singular by construction, so  $D''_{lr}$  and hence  $B''$  are well-defined.

### 4.3 Properties of Canonical Signing

**Lemma 33.**  $B''$  from Definition 31 is a signing of  $B$ .

*Proof.* By Lemma 30,  $B''_l$  and  $B''_r$  are TU signings of  $B_l$  and  $B_r$ , respectively. As a result, blocks  $A''_l, A''_r, D''_l, D''_r$ , and  $D''_0$  in  $B''$  are signings of the corresponding blocks in  $B$ . Thus, it remains to show that  $D''_{lr}$  is a signing of  $D_{lr}$ . This can be verified via a direct calculation.  $\square$

need details?

**Lemma 34.** Suppose that  $B_r$  from Definition 23 has a TU signing  $B'_r$ . Let  $B''_r$  be the canonical re-signing (from Definition 29) of  $B'_r$ . Let  $c''_0 = B''_r(X_r, y_0)$ ,  $c''_1 = B''_r(X_r, y_1)$ , and  $c''_2 = c''_0 - c''_1$ . Then the following statements hold.

1. For every  $i \in X_r$ ,  $\begin{bmatrix} c''_0(i) & c''_1(i) \end{bmatrix} \in \{0, \pm 1\}^{\{y_0, y_1\}} \setminus \left\{ \begin{bmatrix} 1 & -1 \end{bmatrix}, \begin{bmatrix} -1 & 1 \end{bmatrix} \right\}$ .
2. For every  $i \in X_r$ ,  $c''_2(i) \in \{0, \pm 1\}$ .
3.  $\begin{bmatrix} c''_0 & c''_2 & A''_r \end{bmatrix}$  is TU.
4.  $\begin{bmatrix} c''_1 & c''_2 & A''_r \end{bmatrix}$  is TU.



5.  $\begin{bmatrix} c_0'' & c_1'' & c_2'' & A_r'' \end{bmatrix}$  is TU.

*Proof.* Throughout the proof we use that  $B_r''$  is TU, which holds by Lemma 30.

1. Since  $B_r''$  is TU, all its entries are in  $\{0, \pm 1\}$ , and in particular  $\begin{bmatrix} c_0''(i) & c_1''(i) \end{bmatrix} \in \{0, \pm 1\}^{\{y_0, y_1\}}$ . If  $\begin{bmatrix} c_0''(i) & c_1''(i) \end{bmatrix} = \begin{bmatrix} 1 & -1 \end{bmatrix}$ , then

$$\det B_r''(\{x_2, i\}, \{y_0, y_1\}) = \det \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = -2 \notin \{0, \pm 1\},$$

which contradicts TUness of  $B_r''$ . Similarly, if  $\begin{bmatrix} c_0''(i) & c_1''(i) \end{bmatrix} = \begin{bmatrix} -1 & 1 \end{bmatrix}$ , then

$$\det B_r''(\{x_2, i\}, \{y_0, y_1\}) = \det \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} = 2 \notin \{0, \pm 1\},$$

which contradicts TUness of  $B_r''$ . Thus, the desired statement holds.

2. Follows from item 1 and a direct calculation.  
 3. Performing a short tableau pivot in  $B_r''$  on  $(x_2, y_0)$  yields:

$$B_r'' = \begin{array}{|c|c|c|} \hline \textcircled{1} & 1 & 0 \\ \hline c_0'' & c_1'' & A_r'' \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|} \hline 1 & 1 & 0 \\ \hline -c_0'' & c_1'' - c_0'' & A_r'' \\ \hline \end{array}$$

The resulting matrix can be transformed into  $\begin{bmatrix} c_0'' & c_2'' & A_r'' \end{bmatrix}$  by removing row  $x_2$  and multiplying columns  $y_0$  and  $y_1$  by  $-1$ . Since  $B_r''$  is TU and since TUness is preserved under pivoting, taking submatrices, multiplying columns by  $\pm 1$  factors, we conclude that  $\begin{bmatrix} c_0'' & c_2'' & A_r'' \end{bmatrix}$  is TU.

4. Similar to item 4, performing a short tableau pivot in  $B_r''$  on  $(x_2, y_1)$  yields:

$$B_r'' = \begin{array}{|c|c|c|} \hline 1 & \textcircled{1} & 0 \\ \hline c_0'' & c_1'' & A_r'' \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|} \hline 1 & 1 & 0 \\ \hline c_0'' - c_1'' & -c_1'' & A_r'' \\ \hline \end{array}$$

The resulting matrix can be transformed into  $\begin{bmatrix} c_1'' & c_2'' & A_r'' \end{bmatrix}$  by removing row  $x_2$ , multiplying column  $y_1$  by  $-1$ , and swapping the order of columns  $y_0$  and  $y_1$ . Since  $B_r''$  is TU and since TUness is preserved under pivoting, taking submatrices, multiplying columns by  $\pm 1$  factors, and re-ordering columns, we conclude that  $\begin{bmatrix} c_1'' & c_2'' & A_r'' \end{bmatrix}$  is TU.

5. Let  $V$  be a square submatrix of  $\begin{bmatrix} c_0'' & c_1'' & c_2'' & A_r'' \end{bmatrix}$ . Our goal is to show that  $\det V \in \{0, \pm 1\}$ .

Suppose that column  $c_2''$  is not in  $V$ . Then  $V$  is a submatrix of  $B_r''$ , which is TU. Thus,  $\det V \in \{0, \pm 1\}$ . Going forward we assume that column  $z$  is in  $V$ .

Suppose that columns  $c_0''$  and  $c_1''$  are both in  $V$ . Then  $V$  contains columns  $c_0''$ ,  $c_1''$ , and  $c_2'' = c_0'' - c_1''$ , which are linearly. Thus,  $\det V = 0$ . Going forward we assume that at least one of the columns  $c_0''$  and  $c_1''$  is not in  $V$ .

Suppose that column  $c_1''$  is not in  $V$ . Then  $V$  is a submatrix of  $\begin{bmatrix} c_0'' & c_2'' & A_r'' \end{bmatrix}$ , which is TU by item 3. Thus,  $\det V \in \{0, \pm 1\}$ . Similarly, if column  $c_0''$  is not in  $V$ , then  $V$  is a submatrix of  $\begin{bmatrix} c_1'' & c_2'' & A_r'' \end{bmatrix}$ , which is TU by item 4. Thus,  $\det V \in \{0, \pm 1\}$ .

□

**Lemma 35.** Suppose that  $B_l$  from Definition 23 has a TU signing  $B'_l$ . Let  $B''_l$  be the canonical re-signing (from Definition 29) of  $B'_l$ . Let  $d''_0 = B''_l(x_0, Y_l)$ ,  $d''_1 = B''_l(x_1, Y_l)$ , and  $d''_2 = d''_0 - d''_1$ . Then the following statements hold.

1. For every  $j \in Y_l$ ,  $\begin{bmatrix} d''_0(j) \\ d''_1(j) \end{bmatrix} \in \{0, \pm 1\}^{\{x_0, x_1\}} \setminus \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$ .
2. For every  $j \in Y_l$ ,  $d''_2(j) \in \{0, \pm 1\}$ .
3.  $\begin{bmatrix} A''_l \\ d''_0 \\ d''_2 \end{bmatrix}$  is TU.
4.  $\begin{bmatrix} A''_l \\ d''_1 \\ d''_2 \end{bmatrix}$  is TU.
5.  $\begin{bmatrix} A''_l \\ d''_0 \\ d''_1 \\ d''_2 \end{bmatrix}$  is TU.

*Proof.* Apply Lemma 34 to  $B_l^\top$ , or repeat the same arguments up to transposition.  $\square$

**Lemma 36.** Let  $B''$  be from Definition 31. Let  $c''_0 = B''(X_r, y_0)$ ,  $c''_1 = B''(X_r, y_1)$ , and  $c''_2 = c''_0 - c''_1$ . Similarly, let  $d''_0 = B''(x_0, Y_l)$ ,  $d''_1 = B''(x_1, Y_l)$ , and  $d''_2 = d''_0 - d''_1$ . Then the following statements hold.

1. For every  $i \in X_r$ ,  $c''_2(i) \in \{0, \pm 1\}$ .
2. If  $D''_0 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ , then  $D'' = c''_0 \otimes d''_0 - c''_1 \otimes d''_1$ . If  $D''_0 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ , then  $D'' = c''_0 \otimes d''_0 - c''_0 \otimes d''_1 + c''_1 \otimes d''_1$ .
3. For every  $j \in Y_l$ ,  $D''(X_r, j) \in \{0, \pm c''_0, \pm c''_1, \pm c''_2\}$ .
4. For every  $i \in X_r$ ,  $D''(i, Y_l) \in \{0, \pm d''_0, \pm d''_1, \pm d''_2\}$ .
5.  $\begin{bmatrix} D'' & A''_r \end{bmatrix}$  is TU.
6.  $\begin{bmatrix} A''_l \\ D'' \end{bmatrix}$  is TU.

*Proof.* 1. Holds by Lemma 34.2.

2. Note that

$$\begin{bmatrix} D''_l \\ D''_{lr} \end{bmatrix} = \begin{bmatrix} D''_0 \\ D''_r \end{bmatrix} \cdot (D''_0)^{-1} \cdot D''_l, \quad \begin{bmatrix} D''_0 \\ D''_r \end{bmatrix} = \begin{bmatrix} D''_0 \\ D''_r \end{bmatrix} \cdot (D''_0)^{-1} \cdot D''_0, \quad \begin{bmatrix} D''_0 \\ D''_r \end{bmatrix} = \begin{bmatrix} c''_0 & c''_1 \end{bmatrix}, \quad \begin{bmatrix} D''_l & D''_0 \end{bmatrix} = \begin{bmatrix} d''_0 \\ d''_1 \end{bmatrix}.$$

Thus,

$$D'' = \begin{bmatrix} D''_l & D''_0 \\ D''_{lr} & D''_r \end{bmatrix} = \begin{bmatrix} D''_0 \\ D''_r \end{bmatrix} \cdot (D''_0)^{-1} \cdot \begin{bmatrix} D''_l & D''_0 \end{bmatrix} = \begin{bmatrix} c''_0 & c''_1 \end{bmatrix} \cdot (D''_0)^{-1} \cdot \begin{bmatrix} d''_0 \\ d''_1 \end{bmatrix}.$$

Considering the two cases for  $D''_0$  and performing the calculations yields the desired results.

3. Let  $j \in Y_l$ . By Lemma 35.1,  $\begin{bmatrix} d''_0(j) \\ d''_1(j) \end{bmatrix} \in \{0, \pm 1\}^{\{x_0, x_1\}} \setminus \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$ . Consider two cases.

- (a) If  $D''_0 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ , then by item 2 we have  $D''(X_r, j) = d''_0(j) \cdot c''_0 + (-d''_1(j)) \cdot c''_1$ . By considering all possible cases for  $d''_0(j)$  and  $d''_1(j)$ , we conclude that  $D''(X_r, j) \in \{0, \pm c''_0, \pm c''_1, \pm(c''_0 - c''_1)\}$ .

- (b) If  $D''_0 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ , then by item 2 we have  $D''(X_r, j) = (d''_0(j) - d''_1(j)) \cdot c''_0 + d''_1(j) \cdot c''_1$ . By considering all possible cases for  $d''_0(j)$  and  $d''_1(j)$ , we conclude that  $D''(X_r, j) \in \{0, \pm c''_0, \pm c''_1, \pm(c''_0 - c''_1)\}$ .
4. Let  $i \in X_r$ . By Lemma 34.1,  $\begin{bmatrix} c''_0(i) & c''_1(i) \end{bmatrix} \in \{0, \pm 1\}^{\{y_0, y_1\}} \setminus \left\{ \begin{bmatrix} 1 & -1 \end{bmatrix}, \begin{bmatrix} -1 & 1 \end{bmatrix} \right\}$ . Consider two cases.
- (a) If  $D''_0 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ , then by item 2 we have  $D''(i, Y_l) = c''_0(i) \cdot d''_0 + (-c''_1(i)) \cdot d''_1$ . By considering all possible cases for  $c''_0(i)$  and  $c''_1(i)$ , we conclude that  $D''(i, Y_l) \in \{0, \pm d''_0, \pm d''_1, \pm d''_2\}$ .
- (b) If  $D''_0 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ , then by item 2 we have  $D''(i, Y_l) = c''_0(i) \cdot d''_0 + (c''_1(i) - c''_0(i)) \cdot d''_1$ . By considering all possible cases for  $c''_0(i)$  and  $c''_1(i)$ , we conclude that  $D''(i, Y_l) \in \{0, \pm d''_0, \pm d''_1, \pm d''_2\}$ .
5. By Lemma 34.5,  $\begin{bmatrix} c''_0 & c''_1 & c''_2 & A''_r \end{bmatrix}$  is TU. Since TUness is preserved under adjoining zero columns, copies of existing columns, and multiplying columns by  $\pm 1$  factors,  $\begin{bmatrix} 0 & \pm c''_0 & \pm c''_1 & \pm c''_2 & A''_r \end{bmatrix}$  is also TU. By item 3,  $\begin{bmatrix} D'' & A''_r \end{bmatrix}$  is a submatrix of the latter matrix, hence it is also TU.

6. By Lemma 35.5,  $\begin{bmatrix} A''_l \\ d''_0 \\ d''_1 \\ d''_2 \end{bmatrix}$  is TU. Since TUness is preserved under adjoining zero rows, copies of existing

rows, and multiplying rows by  $\pm 1$  factors,  $\begin{bmatrix} A''_l \\ 0 \\ \pm d''_0 \\ \pm d''_1 \\ \pm d''_2 \end{bmatrix}$  is also TU. By item 4,  $\begin{bmatrix} A''_l \\ D'' \end{bmatrix}$  is a submatrix of the latter matrix, hence it is also TU. □

## 4.4 Proof of Regularity

**Definition 37.** Let  $X_l, Y_l, X_r, Y_r$  be sets and let  $c_0, c_1 \in \mathbb{Q}^{X_r}$  be column vectors such that for every  $i \in X_r$  we have  $c_0(i), c_1(i), c_0(i) - c_1(i) \in \{0, \pm 1\}$ . Define  $\mathcal{C}(X_l, Y_l, X_r, Y_r; c_0, c_1)$  to be the family of matrices of the form  $\begin{bmatrix} A_l & 0 \\ D & A_r \end{bmatrix}$  where  $A_l \in \mathbb{Q}^{X_l \times Y_l}$ ,  $A_r \in \mathbb{Q}^{X_r \times Y_r}$ , and  $D \in \mathbb{Q}^{X_r \times Y_l}$  are such that: (a) for every  $j \in Y_r$ ,

$D(X_r, j) \in \{0, \pm c_0, \pm c_1, \pm(c_0 - c_1)\}$ , (b)  $\begin{bmatrix} c_0 & c_1 & c_0 - c_1 & A_r \end{bmatrix}$  is TU, (c)  $\begin{bmatrix} A_l \\ D \end{bmatrix}$  is TU.

**Lemma 38.** Let  $B''$  be from Definition 31. Then  $B'' \in \mathcal{C}(X_l, Y_l, X_r, Y_r; c''_0, c''_1)$  where  $c''_0 = B''(X_r, y_0)$  and  $c''_1 = B''(X_r, y_1)$ .

*Proof.* Recall that  $c''_0 - c''_1 \in \{0, \pm 1\}^{X_r}$  by Lemma 36.1, so  $\mathcal{C}(X_l, Y_l, X_r, Y_r; c''_0, c''_1)$  is well-defined. To see that  $B'' \in \mathcal{C}(X_l, Y_l, X_r, Y_r; c''_0, c''_1)$ , note that all properties from Definition 37 are satisfied: property (a) holds by Lemma 36.3, property (b) holds by Lemma 34.5, and property (c) holds by Lemma 36.6. □

**Lemma 39.** Let  $C \in \mathcal{C}(X_l, Y_l, X_r, Y_r; c_0, c_1)$  from Definition 37. Let  $x \in X_l$  and  $y \in Y_l$  be such that  $A_l(x, y) \neq 0$ , and let  $C'$  be the result of performing a short tableau pivot in  $C$  on  $(x, y)$ . Then  $C' \in \mathcal{C}(X_l, Y_l, X_r, Y_r; c_0, c_1)$ .

*Proof.* Our goal is to show that  $C'$  satisfies all properties from Definition 37. Let  $C' = \begin{bmatrix} C'_{11} & C'_{12} \\ C'_{21} & C'_{22} \end{bmatrix}$ , and let

$\begin{bmatrix} A'_l \\ D' \end{bmatrix}$  be the result of performing a short tableau pivot on  $(x, y)$  in  $\begin{bmatrix} A_l \\ D \end{bmatrix}$ . Observe the following.

- By Lemma 4,  $C'_{11} = A'_l$ ,  $C'_{12} = 0$ ,  $C'_{21} = D'$ , and  $C'_{22} = A_r$ .

- Since  $\begin{bmatrix} A_l \\ D \end{bmatrix}$  is TU by property (c) for  $C$ , all entries of  $A_l$  are in  $\{0, \pm 1\}$ .
- $A_l(x, y) \in \{\pm 1\}$ , as  $A_l(x, y) \in \{0, \pm 1\}$  by the above observation and  $A_l(x, y) \neq 0$  by the assumption.
- Since  $\begin{bmatrix} A_l \\ D \end{bmatrix}$  is TU by property (c) for  $C$  and since pivoting preserves TUness,  $\begin{bmatrix} A'_l \\ D' \end{bmatrix}$  is also TU.

These observations immediately imply properties (b) and (c) for  $C'$ . Indeed, property (b) holds for  $C'$ , since  $C'_{22} = A_r$  and  $\begin{bmatrix} c_0 & c_1 & c_0 - c_1 & A_r \end{bmatrix}$  is TU by property (b) for  $C$ . On the other hand, property (c) follows from  $C'_{11} = A'_l$ ,  $C'_{21} = D'$ , and  $\begin{bmatrix} A'_l \\ D' \end{bmatrix}$  being TU. Thus, it only remains to show that  $C'$  satisfies property (a). Let  $j \in Y_r$ . Our goal is to prove that  $D'(X_r, j) \in \{0, \pm c_0, \pm c_1, \pm(c_0 - c_1)\}$ .

Suppose  $j = y$ . By the pivot formula,  $D'(X_r, y) = -\frac{D(X_r, y)}{A_l(x, y)}$ . Since  $D(X_r, y) \in \{0, \pm c_0, \pm c_1, \pm(c_0 - c_1)\}$  by property (a) for  $C$  and since  $A_l(x, y) \in \{\pm 1\}$ , we get  $D'(X_r, y) \in \{0, \pm c_0, \pm c_1, \pm(c_0 - c_1)\}$ .

Now suppose  $j \in Y_l \setminus \{y\}$ . By the pivot formula,  $D'(X_r, j) = D(X_r, j) - \frac{A_l(x, j)}{A_l(x, y)} \cdot D(X_r, y)$ . Here  $D(X_r, j), D(X_r, y) \in \{0, \pm c_0, \pm c_1, \pm(c_0 - c_1)\}$  by property (a) for  $C$ , and  $A_l(x, j) \in \{0, \pm 1\}$  and  $A_l(x, y) \in \{\pm 1\}$  by the prior observations. Perform an exhaustive case distinction on  $D(X_r, j), D(X_r, y), A_l(x, j)$ , and  $A_l(x, y)$ . In every case, we can show that either  $\begin{bmatrix} A_l(x, y) & A_l(x, j) \\ D(X_r, y) & D(X_r, j) \end{bmatrix}$  contains a submatrix with determinant not in  $\{0, \pm 1\}$ , which contradicts TUness of  $\begin{bmatrix} A_l \\ D \end{bmatrix}$ , or that  $D'(X_r, j) \in \{0, \pm c_0, \pm c_1, \pm(c_0 - c_1)\}$ , as desired.  $\square$  need details?

**Lemma 40.** Let  $C \in \mathcal{C}(X_l, Y_l, X_r, Y_r; c_0, c_1)$  from Definition 37. Then  $C$  is TU.

*Proof.* By Lemma 3, it suffices to show that  $C$  is  $k$ -PU for every  $k \in \mathbb{Z}_{\geq 1}$ . We prove this claim by induction on  $k$ . The base case with  $k = 1$  holds, since properties (b) and (c) in Definition 37 imply that  $A_l, A_r$ , and  $D$  are TU, so all their entries of  $C = \begin{bmatrix} A_l & 0 \\ D & A_r \end{bmatrix}$  are in  $\{0, \pm 1\}$ , as desired.

Suppose that for some  $k \in \mathbb{Z}_{\geq 1}$  we know that every  $C' \in \mathcal{C}(X_l, Y_l, X_r, Y_r; c_0, c_1)$  is  $k$ -PU. Our goal is to show that  $C$  is  $k$ -PU, i.e., that every  $(k+1) \times (k+1)$  submatrix  $S$  of  $C$  has  $\det V \in \{0, \pm 1\}$ .

First, suppose that  $V$  has no rows in  $X_\ell$ . Then  $V$  is a submatrix of  $\begin{bmatrix} D & A_r \end{bmatrix}$ , which is TU by property (b) in Definition 37, so  $\det V \in \{0, \pm 1\}$ . Thus, we may assume that  $S$  contains a row  $x_\ell \in X_\ell$ .

Next, note that without loss of generality we may assume that there exists  $y_\ell \in Y_\ell$  such that  $V(x_\ell, y_\ell) \neq 0$ . Indeed, if  $V(x_\ell, y) = 0$  for all  $y$ , then  $\det V = 0$  and we are done, and  $V(x_\ell, y) = 0$  holds whenever  $y \in Y_r$ .

Since  $C$  is 1-PU, all entries of  $V$  are in  $\{0, \pm 1\}$ , and hence  $V(x_\ell, y_\ell) \in \{\pm 1\}$ . Thus, by Lemma 6, performing a short tableau pivot in  $V$  on  $(x_\ell, y_\ell)$  yields a matrix that contains a  $k \times k$  submatrix  $S''$  such that  $|\det V| = |\det V''|$ . Since  $V$  is a submatrix of  $C$ , matrix  $V''$  is a submatrix of the matrix  $C'$  resulting from performing a short tableau pivot in  $C$  on the same entry  $(x_\ell, y_\ell)$ . By Lemma 39, we have  $C' \in \mathcal{C}(X_l, Y_l, X_r, Y_r; c_0, c_1)$ . Thus, by the inductive hypothesis applied to  $V''$  and  $C'$ , we have  $\det V'' \in \{0, \pm 1\}$ . Since  $|\det V| = |\det V''|$ , we conclude that  $\det V \in \{0, \pm 1\}$ .  $\square$

**Lemma 41.**  $B''$  from Definition 31 is TU.

*Proof.* Combine the results of Lemmas 38 and 40.  $\square$

**Lemma 42.** Let  $M$  be a 3-sum of regular matroids  $M_\ell$  and  $M_r$ . Then  $M$  is also regular.

*Proof.* Let  $B, B_\ell$ , and  $B_r$  be standard  $\mathbb{Z}_2$  representation matrices from Definition 25. Since  $M_\ell$  and  $M_r$  are regular, by Lemma 16,  $B_\ell$  and  $B_r$  have TU signings. Then the canonical signing  $B''$  from Definition 31 is a TU signing of  $B$ . Indeed,  $B''$  is a signing of  $B$  by Lemma 33, and  $B''$  is TU by Lemma 41. Thus,  $M$  is regular by Lemma 16.  $\square$