Report on the main results of the surveillance under article 11 for annex II, IV and V species (Annex B)

0.1 Member State	**IT**
0.2.1 Species code	**1090**
0.2.2 Species name	**Scyllarides latus**
0.2.3 Alternative species scientific name	N/A
0.2.4 Common name	Magnosa

1. National Level

1.1 Maps

1.1.1 Distribution Map	Yes
1.1.1a Sensitive species	No
1.1.2 Method used - map	Estimate based on expert opinion with no or minimal sampling (1)
1.1.3 Year or period	2001-2012
1.1.4 Additional map	No
1.1.5 Range map	Yes

2. Biogeographical Or Marine Level

2.1 Biogeographical Region

Marine Mediterranean (MMED)

2.2 Published sources

The present species assessment (fields 0.1-2.9) has been compiled by Anna Alonzi, Piero Genovesi, Francesca Ronchi (ISPRA). Information and data have been extracted from MSFD Supporting documents on the Initial Assessmant on Benthic Species, including methodology, data used and results (ISPRA, 2013). Expert judgements have been provided by Leonardo Tunesi (ISPRA).

BIANCHINI, M. L., CHESSA, L., GRECO, S., RAGONESE, S. and SCARPELLI, G., 1996. Morphometric aspects of slipper lobster, Scyllarides latus, in Abstracts of the 6th Colloquium Crustacea Decapoda Mediterranea, Florence 12-15 September, 1996, pp. 11-12.

BIANCHINI M. L., CHESSA L., GRECO S., RAGONESE S., 1996 (cf. a) - An Italian enhancement programfor slipper lobster, Scyllarides latus. IIWorld Fisheries Congress (Brisbane,Australia, 28/07-02/08/1996): 91.

BIANCHINI M. L., RAISA P.F., 1997 - Valutazione della fattibilità e potenzialità del ripopolamento attivo per la magnosa, Scyllarides latus (Crostacei Decapodi). Final report to MiRAAF (Pesca Marittima), 1, 1997.

BIANCHINI M. L., RAGONESE, S., GRECO, S., CHESSA, L. & BIAGI, F., 1998 - Il progetto “Valutazione della fattibilità e potenzialità del ripopolamento attivo per la magnosa, Scyllarides latus (Crostacei Decapodi) ” Rapporto: sintesi e risultati Biologia Marina Mediterranea, 5 (3): 1277-1283.

BIANCHINI M. L., SPANIER. E., RAGONESE S., 1998 - Enzymatic variability of Mediterranean slipper lobster Scyllarides latus, from Sicilian waters. Annals – Annals for Istrian and Mediterranean Studies, Ser Hist Nat., 13: 43-50.

BIANCHINI M. L., BONO G., RAGONESE S., 2001 - Long-term recaptures and growth of slipper lobsters, Scyllarides latus, in the Strait of Sicily (Mediterranean Sea). Crustaceana 74: 673-680.

BIANCHINI M. L., RAGONESE, S., 2003 - In ovo embryonic development of the Mediterranean slipper lobster, scyllarides latus, the lobster newsletter, 16, 10-12.

CHESSA, L. A., PAIS, A. and SERRA, S., 1996, Behavioral observations on slipper lobster Scyllarides latus (Latreille 1803) (Decapoda, Scyllaridae) reared in laboratory, in Abstracts of the 6th Colloquium Crustacea Decapoda Mediterranea, Florence 12-15 September, 1996, pp. 25 26

Report on the main results of the surveillance under article 11 for annex II, IV and V species (Annex B)

CONSORZIOMEDITERRANEO, 2001 - Studio della fattibilità e potenzialità del ripopolamento attivo per la magnosa, Scyllarides latus. Rapporto Finale. Centro Servizi, Lega Pesca; Servizi Orizzontali. Ministero Politiche Agricole & CE.
FISCHER W., GAUCHO L., SCHNEIDER M., 1987 – Fiches FAO d'identification des especies pour les besoins de la peche . Rev. 1. Mediterranee et Mer Noir. Zone de peche 37. Invertebres. Rome, FAO 1: 1-760.
Frau F., Cinti M.F., Paliaga B., Guala I. 2011. Protected species according to the Spa/Bio Protocol (Barcelona Convention) present in the Marine Protected Area Capo Carbonara. 42° Congresso della Società Italiana di Biologia Marina, Olbia, 23-28 maggio 2011: 262-263.
FROESE R., PAULY D. (ed), 2001 – Fishbase Word Wild Web electronic publication. http://www.fishbase.org
HOLTHUIS L.B., 1991 – Marine lobster of the word. FAO Species Catalogue 13: 292 pp.
Regione Autonoma della Sardegna - Assessorato Difesa Ambiente , 2012 - "Servizio di monitoraggio dello stato di conservazione degli habitat e delle specie di importanza comunitaria presenti nei siti della Rete Natura 2000 in Sardegna – Linea 4. Redazione del Rapporto sullo stato di conservazione degli habitat e delle specie "
RELINI G., BERTRAND J., ZAMBONI A. (eds), 1999 – sintesi delle conoscenze sulle risorse da pesca dei fondi del Mediterraneo central (Italia e Corsica). Biol. Mar. Med., 6 (suppl. 1): 868 pp.
ROMEO T., G. FLORIO, F. LENTINI, L. CASTRIOTA, M. FALAUTANO, P. CONSOLI, P.PELUSI, & S. GRECO., 2004 -Morphometric aspects of Scyllarides latus. Mediterranean Marine Science 5/2: 65-71.
SANNA M. T., OLIANAS A., CASTAGNOLA M., SOLLAI L., MANCONI B., SALVADORI S., GIARDINA B., PELLEGRINI M., 2004 - Oxygen-binding modulation of hemocyanin from the slipper lobster Scyllarides latus. Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology 139 (2): 261-268.

2.3 Range

2.3.1 Surface area - Range (km²)	474900	
2.3.2 Method - Range surface area	Estimate based on expert opinion with no or minimal sampling (1)	
2.3.3 Short-term trend period	2001-2012	
2.3.4 Short-term trend direction	unknown (x)	
2.3.5 Short-term trend magnitude	min	max
2.3.6 Long-term trend period		
2.3.7 Long-term trend direction	N/A	
2.3.8 Long-term trend magnitude	min	max
2.3.9 Favourable reference range	area (km²)	
	operator	approximately equal to (≈)
	unkown	No
	method	Expert judgement
2.3.10 Reason for change	Improved knowledge/more accurate dataUse of different method	

2.4 Population

2.4.1 Population size (individuals or agreed exception)	Unit	N/A	
	min		max
2.4.2 Population size (other than individuals)	Unit	N/A	
	min		max

Report on the main results of the surveillance under article 11 for annex II, IV and V species (Annex B)

2.4.3 Additional information	Definition of locality		
	Conversion method		
	Problems		
2.4.4 Year or period			
2.4.5 Method – population size	Absent data (0)		
2.4.6 Short-term trend period	2001-2012		
2.4.7 Short term trend direction	unknown (x)		
2.4.8 Short-term trend magnitude	min	max	confidence interval
2.4.9 Short-term trend method	Absent data (0)		
2.4.10 Long-term trend period			
2.4.11 Long term trend direction	N/A		
2.4.12 Long-term trend magnitude	min	max	confidence interval
2.4.13 Long-term trend method	N/A		
2.4.14 Favourable reference population	number		
	operator	much more than (>>)	
	unknown	No	
	method	Expert judgement	
2.4.15 Reason for change	Improved knowledge/more accurate data		

2.5 Habitat for the Species

2.5.1 Surface area - Habitat (km²)	
2.5.2 Year or period	
2.5.3 Method used - habitat	Absent data (0)
2.5.4 a) Quality of habitat	Good
2.5.4 b) Quality of habitat - method	expert judgement
2.5.5 Short term trend period	2001-2012
2.5.6 Short term trend direction	unknown (x)
2.5.7 Long-term trend period	
2.5.8 Long term trend direction	N/A
2.5.9 Area of suitable habitat (km²)	
2.5.10 Reason for change	Improved knowledge/more accurate data

2.6 Main Pressures

Pressure	ranking	pollution qualifier(s)
scubadiving, snorkelling (G01.07)	high importance (H)	N/A
Fishing and harvesting aquatic resources (F02)	medium importance (M)	N/A
potting (F02.01.01)	medium importance (M)	N/A

2.6.1 Method used – pressures	mainly based on expert judgement and other data (2)

2.7 Main Threats

Threat	ranking	pollution qualifier(s)
Fishing and harvesting aquatic resources (F02)	medium importance (M)	N/A
scubadiving, snorkelling (G01.07)	high importance (H)	N/A
potting (F02.01.01)	medium importance (M)	N/A

2.7.1 Method used – threats	expert opinion (1)

2.8 Complementary Information

Report on the main results of the surveillance under article 11 for annex II, IV and V species (Annex B)

2.8.1 Justification of % thresholds for trends	
2.8.2 Other relevant Information	
2.8.3 Trans-boundary assessment	

2.9 Conclusions (assessment of conservation status at end of reporting period)

2.9.1 Range	assessment	Favourable (FV)
	qualifiers	N/A
2.9.2. Population	assessment	Bad (U2)
	qualifiers	unknown (x)
2.9.3. Habitat	assessment	Favourable (FV)
	qualifiers	N/A
2.9.4. Future prospects	assessment	Inadequate (U1)
	qualifiers	N/A
2.9.5 Overall assessment of Conservation Status	Bad (U2)	
2.9.5 Overall trend in Conservation Status	stable (=)	

3. Natura 2000 coverage and conservation measures - Annex II species

3.1 Population

3.1.1 Population Size	Unit	N/A	
	min		max
3.1.2 Method used	N/A		
3.1.3 Trend of population size within	N/A		

3.2 Conversation Measures

Notes

Species name: Scyllarides latus (1090) Region code: MMED

Field label	Note	User
2.3.1 Surface area - Range (km^2)	It is important to point out that only the surface area of the habitat that can actually host the species should be considered.	ISPRA_AUNA

_F