0.1 Member State	IT
0.2.1 Species code	2621
0.2.2 Species name	Balaenoptera physalus
0.2.3 Alternative species scientific name	N/A
0.2.4 Common name	Balenottera comune

#### 1. National Level

#### **1.1 Maps**

1.1.1 Distribution Map	Yes
1.1.1a Sensitive species	No
1.1.2 Method used - map	Estimate based on partial data with some extrapolation and/or modelling (2)
1.1.3 Year or period	2010-2011
1.1.4 Additional map	No
1.1.5 Range map	Yes

#### 2. Biogeographical Or Marine Level

2.1 Biogeographical Region

#### 2.2 Published sources

#### Marine Mediterranean (MMED)

The present species assessment (fields 0.1-2.9) has been compiled by Anna Alonzi, Piero Genovesi, Francesca Ronchi (ISPRA). Information and data have been extracted from the MSFD Supporting documents on the Initial Assessment on Cetaceans, including methodology, data used and results (ISPRA,2013). Contributing authors: Caterina Fortuna, Mario Acquarone, Aldo Annunziatellis, Antonella Arcangeli, Arianna Azzellino, Nicola Baccetti, Michela Bellingeri, Silvia Bonizzoni, Junio Fabrizio Borsani, Ilaria Caliani, Simonepietro Canese, Roberta Canneri, Nadia Cerioli, Andrea De Lucia, Salvatore Dimatteo, Carmelo Fanizza, Elio Filidei jr., Maria Cristina Fossi, Fulvio Garibaldi, Stefania Gaspari, Otello Giovanardi, Michela Giusti, Guido Gnone, Paolo Guidetti, Drasko Holcer, Giancarlo Lauriano, Letizia Marsili, Antonio Mazzola, Giulia Mo, Aurelie Moulins, Barbara Mussi, Giuseppe

Notarbartolo di Sciara, Lidia Orsi Relini, Daniela Silvia Pace, Simone Panigada, Gianni Pavan, Michela Podestà, Marina Pulcini, Sasa Raicevich, Ettore Randi, Teresa Romeo, Massimiliano Rosso, Antonello Sala, Paola Tepsich, Walter Zimmer e Nicola Zizzo. Expert judgements have been provided by Caterina Fortuna (ISPRA).

A.A.V.V. 2007. Workshop d'inquadramento del fenomeno delle interazioni tra cetacei e attività di pesca in Italia. Roma, 10-11 settembre 2007, Ministero delle politiche agricole, alimentari e forestali, Roma. Progetto «Programma Bycatch Italia-ACCOBAMS» (codice 6G23), finanziato dalla Direzione Generale per la Pesca marittima e Acquacoltura, MiPAAF.

Arcangeli, A., Marini, L., Crosti, R. 2012. Changes in cetacean presence, relative abundance and distribution over 20 years along a trans-regional fixed line transect in the Central Tyrrhenian Sea. Marine Ecology, article first published online: 16 OCT 2012, DOI: 10.1111/maec.12006.

Arrigoni M., Manfredi P., Panigada S., Bramanti L., Santangelo G. 2011 Lifehistory tables of the Mediterranean fin whale from stranding data. Marine Ecology 32 (Suppl. 1):1–9.

Azzellino, A., Airoldi, S., Gaspari, S., Nani, B., 2008. Habitat use of cetaceans along the continental slope and adjacent waters in the western Ligurian Sea.

30/04/2014 10.57.58 Page 1 of 6

Deep Sea Res. 55, 296-323.

Azzellino, A., Lanfredi, C., D'Amico, A., Pavan, G., Podestà, M., Haun, J., 2011. Risk mapping for sensitive species to underwater anthropogenic sound emissions: model development and validation in two Mediterranean areas. Mar. Pollut. Bull. 63, 56-70.

Azzellino, A., Panigada, S., Lanfredi, C., Zanardelli, M., Airoldi, S., Notarbartolo di Sciara, G. 2012. Predictive habitat models for managing marine areas: Spatial and temporal distribution of marine mammals within the Pelagos Sanctuary (Northwestern Mediterranean sea). Ocean & Coastal Management 67 (2012) 63-74.

Buckland ST, Anderson DR, Burnham KP, Laake JL, Borchers DL, Thomas L 2001. Introduction to distance sampling: stimating abundance of biological populations. Oxford University Press, Oxford.

Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., Borchers, D.L., Thomas, L. 2004. Advanced Distance Sampling. Oxford University Press.

Cañadas, A., Fortuna, C., Pulcini, M., Lauriano, G., Bearzi, G., Cotte, C., Raga, J.A., Panigada, S., Politi, E., Rendell, L., B-Nagy, A.1, Pastor, X., Frantzis, A., Mussi, B. 2011. Accobams collaborative effort to map high-use areas by beaked whales in the Mediterranean. Scientific Committee of the International Whaling Commission. SC/63/SM10

Cañadas, A., R. Sagarminaga, R. de Stephanis, E. Urquiola and P.S. Hammond. 2005. Habitat selection models as a conservation tool: proposal of marine protected areas for cetaceans in Southern Spain. Aquatic Conservation: Marine and Freshwater Ecosystems 15:495-521.

Cañadas, A., Sagarminaga, R. and García-Tiscar, S. 2002. Cetacean distribution related with depth and slope in the Mediterranean waters off southern Spain. Deep Sea Research I 49: 2053-2073.

Canese, S., Cardinali, A., Fortuna, C.M., Giusti, M., Lauriano, G., Salvati, E. and Greco, S. 2006. The first identified winter feeding ground of fin whales (Balaenoptera physalus) in the Mediterranean Sea. J. Mar. Biol. Ass. U.K. (2006), 86: 903-907

Castellote, M., Clark, C.W., Lammers, M.O. 2012a. Acoustic and behavioural changes by fin whales (Balaenoptera physalus) in response to shipping and airgun noise. Biological Conservation 147(1):115-122.

Castellote, M., Clark, C.W., Lammers, M.O. 2012b. Fin whale (Balaenoptera physalus) population identity in the western Mediterranean Sea. Marine Mammal Science 28: 325-344.

Caughley G. 1966. Mortality patterns in mammals. Ecology 47:906–918. Cerioni, S., Forni, L., Lo Tenero, A., Nannarelli, S. & Pulcini, M. 1996. A cetacean survey in the Taranto Gulf: work in progress. European Research on Cetaceans 9:102–103.

D'Ortenzio, F. and Ribera d'Alcala, M. 2008. On the trophic regimes of the Mediterranean Sea. Biosciences Discussions 5: 2959-2983.

Emelen J.M. 1970. Age specificity and ecological theory. Ecology, 51, 588–601. Forcada, J., Notarbartolo di Sciara, G. and Fabbri, F. 1995. Abundance of fin whales and striped dolphins summering in the Corso-Ligurian basin. Mammalia 59(1): 127-140

Fortuna, C.M., Filidei, E. jr. 2011. Annual Report on the implementation of Council Regulation (EC) 812/2004 - 2010. Rapporto tecnico preparato per il Ministero delle politiche agricole, alimentari e forestali, 10 pagine. Fortuna, C.M., Holcer, D., Filidei, E. jr, Tunesi, L., 2011 Relazione finale del progetto "Valutazione dell'impatto della mortalità causata da attività di pesca su Cetacei e tartarughe marine in Adriatico: primo survey per la stima

30/04/2014 10.57.58 Page 2 of 6

dell'abbondanza" (Prot. MIPAAF DG PEMAC n. 1690 del 10/02/2010 e al Prot. MATTM DPN n. 27623 del 23/12/2009), 51 pagine + Allegati.

Fortuna, C.M., Vallini, C., Filidei, E. jr, Ruffino, M., Consalvo, I., Di Muccio, S., Gion, C., Scacco, U., Tarulli, E., Giovanardi, O., Mazzola, A. 2010. Bycatch of cetaceans and other species of conservation concern during pair trawl fishing operations in the Adriatic Sea (Italy). Chemistry and Ecology 26(Supplement):65-76.

Fossi M. C., Urban J., Casini S., Maltese S., Spinsanti G.. Panti C., Porcelloni S., Panigada S., Lauriano G., Niño-Torres C., Rojas-Bracho L., Jimenez B., Muñoz-Arnanz J.and Marsili L. 2009. A Multi-Trial Diagnostic Tool In Fin Whale (Balaenoptera physalus) Skin Biopsies Of The Pelagos Sanctuary (Mediterranean Sea) And The Gulf Of California (Mexico). Marine environmental research (doi:10.1016/j.marenvres.2009.10.006)

Gannier A 1998. Les cétacés de Méditerranée nord-occidentale: nouveaux résultats sur leur distribution, la structure de leur peuplement et l'abondance relative des différentes espèces. Mésogée 56: 3-19.

Gannier A 2006. Le peuplement estival de cetaces dans le Sanctuaire Marin Pelagos (Mediterranee nord-occidentale): distribution et abondance. Mammalia 70: 17-27.

Gannier, A., Bourreau, S., Drouot, V., Gannier, O., Laran, S. 2002. Summer distribution of fin whale (Balaenoptera physalus) in the Mediterranean Sea. Mésogée, Bull. Mus. Hist. Nat. Marseille 60:17-25.

Laran S, Gannier A (2008) Spatial and temporal prediction of fin whale distribution in the northwestern Mediterranean Sea. ICES Journal of Marine Science, 65: 1260–1269.

Lauriano, G., Panigada, S., Fortuna, C.M., Holcer, D., Filidei, E. jr, Pierantonio, N. and Donovan, G. 2011. Monitoring density and abundance of cetaceans in the seas around Italy through aerial survey: a contribution to conservation and the future ACCOBAMS Survey. International Whaling Commission SC/63/SM6. Manca Zeichen, M., Finoia, M.G. and S. Donfrancesco, B., V. 2008. On the use of Optical Remote Sensing to Assess Phytoplankton Dynamics in the Pelagos Sanctuary (Ligurian-Provencal Sea) Ocean Optics XIX. 5–10 October. Marini, L., Consiglio, C., Angradi, A.M., Catalano, B., Sanna, A., Valentini, T., Finoia, M.G. & Villetti, G. 1006. Distribution, abundance and seasonality of

Finoia, M.G. & Villetti, G. 1996. Distribution, abundance and seasonality of cetaceans sighted during scheduled ferry crossings in the central Tyrrhenian Sea: 1989–1992. Italian Journal of Zoology 63:381–388.

Marini, L., Villetti, G. & Consiglio, C., 1995. Wintering areas of Fin whales (Balaenoptera physalus) in the Mediterranean Sea, a preliminary survey. European Research on Cetaceans 9:126-128.

Marsili, L. (2000) Lipophilic contaminants in marine mammals: review of the results of ten years' work at the Department of environmental biology, Siena University (Italy). International Journal of Environmental Pollution 13:416–452. Notarbartolo di Sciara, G., Zanardelli, M., Jahoda, M., Panigada, S., Airoldi, S. 2003. The Fin whale Balaenoptera physalus (L. 1758) in the Mediterranean Sea. Mammal Review 33:105-150.

Orsi Relini L. 2000. The Cetacean Sanctuary in the Ligurian Sea: a further reason. Biol. Mar. Mediterr., 7 (3): 117-126.

Orsi Relini L., Palandri G., Garibaldi F., Lanteri L. 2004. Note su alcuni parametri di popolazione della balenottera comune, Balaenoptera physalus (Linneo, 1758). Biol. Mar. Mediterr., 11 (2): 138-154.

Panigada S, Zanardelli M, MacKenzie M, Donovan C, Mélin F, et al. (2008) Modelling habitat preferences for fin whales and striped dolphins in the Pelagos Sanctuary (Western Mediterranean Sea) with physiographic and remote sensing

30/04/2014 10.57.58 Page 3 of 6

variables. Remote Sens Environ 112: 3400-3412.

Panigada, S., Lauriano, G., Burt, L., Pierantonio, N., Donovan, G. 2011. Monitoring winter and summer abundance of cetaceans in the Pelagos Sanctuary (Northwestern Mediterranean Sea) through aerial surveys. Plos one 6(7): e22878. Doi:10.1371/journalpone.0022878

Panigada, S., Pesante, G., Zanardelli, M., Capoulade, F., Gannier, A. and Weinrich, M. 2006. Medditerranean fin whales at risk from fatal ship strikes. Mar. Poll. Bull. 52: 1287-98.

Politi, E., Bearzi, M., Notarbartolo di Sciara, G., Cussino, E. & Gnone, G. 1992. Distribution and frequency of cetaceans in the waters adjacent to the Greek Ionian Islands. European Research on Cetaceans 6:75-78

RAC/SPA. 1998. Report of the Meeting of Experts on the Implementation of the Action Plans for Marine Mammals (Monk seal and Cetaceans) adopted within MAP. UNEP(OCA)/MED WG. 146/5, Tunis, annex 5, pp:55-57.

Rallo, G. 1979. I cetacei dell'Adriatico. WWF Veneto, 4, 3-4.

Zanardelli M., Panigada S., Airoldi S., Borsani J.F., Jahoda M., Notarbartolo di Sciara G. (1999) Site fidelity, seasonal residence and sex ratio of fin whales (Balaenoptera physalus) in the Ligurian Sea feeding grounds. European Research on Cetaceans, 12, 24.

Zanardelli, M., Airoldi, S., Beaubrun, P., Bérubé, M., Borsani, J.F., Gannier, A., Guinet, C., Hammond, P., Jahoda, M., Lauriano, G., Notarbartolo di Sciara, G., Panigada, S. 2011. Long-term photo-identification study of fin whales in the Pelagos Sanctuary (NW Mediterranean). European Research on Cetaceans, 25.

#### 2.3 Range

2.3.1 Surface area - Range (km²)

2.3.2 Method - Range surface area

2.3.3 Short-term trend period

2.3.4 Short-term trend direction

2.3.5 Short-term trend magnitude

2.3.6 Long-term trend period

2.3.7 Long-term trend direction

2.3.8 Long-term trend magnitude

2.3.9 Favourable reference range

352500

Estimate based on partial data with some extrapolation and/or modelling (2)

2000-2011

unknown (x)

min max

N/A

min max

area (km²)

operator N/A unkown Yes

method

2.3.10 Reason for change

Use of different method

#### 2.4 Population

2.4.1 Population size

(individuals or agreed exception)

Unit N/A

min max

2.4.2 Population size (other than individuals) Unit number of map 10x10 km grid cells (grids10x10)

min 775 max 775

2.4.3 Additional information

**Definition of locality** 

Conversion method

**Problems** 

It is not possible to convert grids into individuals

2.4.4 Year or period

2.4.5 Method – population size

2.4.6 Short-term trend period

2010-2011

Estimate based on partial data with some extrapolation and/or modelling (2)

2000-2011

10.57.58 30/04/2014 Page 4 of 6

<ul><li>2.4.7 Short term trend direction</li><li>2.4.8 Short-term trend magnitude</li><li>2.4.9 Short-term trend method</li><li>2.4.10 Long-term trend period</li></ul>	unknown( min Absent data	,	max	confidence interval
2.4.11 Long term trend direction 2.4.12 Long-term trend magnitude 2.4.13 Long-term trend method 2.4.14 Favourable reference population	N/A min N/A number operator unknown method	N/A Yes	max	confidence interval

2.4.15 Reason for change

Improved knowledge/more accurate data Use of different method

#### 2.5 Habitat for the Species

2.5.1 Surface area - Habitat (km²)

2.5.2 Year or period

2.5.3 Method used - habitat

2.5.4 a) Quality of habitat

2.5.4 b) Quality of habitat - method

2.5.5 Short term trend period

2.5.6 Short term trend direction

2.5.7 Long-term trend period

2.5.8 Long term trend direction

2.5.9 Area of suitable habitat (km²)

2.5.10 Reason for change

Absent data (0)

Unknown

expert based

2000-2011

unknown (x)

N/A

#### 2.6 Main Pressures

Pressure	ranking	pollution qualifier(s)
death or injury by collision (G05.11)	high importance (H)	N/A
Noise nuisance, noise pollution (H06.01)	medium importance (M)	N/A
marine macro-pollution (i.e. plastic bags, styrofoam) (H03.03)	low importance (L)	N/A

2.6.1 Method used – pressures mainly based on expert judgement and other data (2)

#### 2.7 Main Threats

Threat	ranking	pollution qualifier(s)
death or injury by collision (G05.11)	high importance (H)	N/A
Noise nuisance, noise pollution (H06.01)	medium importance (M)	N/A
marine macro-pollution (i.e. plastic bags, styrofoam) (H03.03)	low importance (L)	N/A

2.7.1 Method used – threats

expert opinion (1)

#### 2.8 Complementary Information

2.8.1 Justification of % thresholds for trends

2.8.2 Other relevant Information

The observed species distribution indicates that the species' range spans the entire region. The species distribution pattern is in line with its ecological traits, with the Ligurian Sea and Gulf of Lion as important summer feeding grounds, the Tyrrhenian Sea as potential corridor and Lampedusa Island as important winter feeding ground. There are no sufficient data to infer trends, although there is

30/04/2014 10.57.58 Page 5 of 6

some evidence that the species relative abundance is decreasing in a portion of the the Ligurian sea.

2.8.3 Trans-boundary assessment

#### 2.9 Conclusions (assessment of conservation status at end of reporting period)

2.9.1 Range assessment Unknown (XX) qualifiers N/A

2.9.2. Population assessment Unknown (XX)

qualifiers N/A

2.9.3. Habitat assessment Unknown (XX)

qualifiers N/A

assessment Unknown (XX)

qualifiers N/A

Unknown (XX)

2.9.5 Overall assessment of Conservation Status

Lonservation Status

2.9.4. Future prospects

2.9.5 Overall trend in Conservation Status

N/A

### 3. Natura 2000 coverage and conservation measures - Annex II species

#### 3.1 Population

3.1.1 Population Size Unit N/A

min max

3.1.2 Method used N/A

3.1.3 Trend of population size within N/A

#### **3.2 Conversation Measures**

30/04/2014 10.57.58 Page 6 of 6

### Notes

Species name: Balaenopte	era physalus (2621) Region code: MMED	
Field label	Note	User
2.4.7 Short term trend direction	There are not sufficent data to infer trends.	ISPRA_ AUNA
2.3.4 Range Trend	There are not sufficent data to infer trends.	ISPRA_ AUNA

30/04/2014 10.57.45 Page 1