0.1 Member State	ΙΤ
0.2.1 Species code	1312
0.2.2 Species name	Nyctalus noctula
0.2.3 Alternative species scientific name	N/A
0.2.4 Common name	N/A

1. National Level

1.1 Maps

1.1.1 Distribution Map	Yes
1.1.1a Sensitive species	No
1.1.2 Method used - map	Estimate based on partial data with some extrapolation and/or modelling (2)
1.1.3 Year or period	1985-2012
1.1.4 Additional map	No
1.1.5 Range map	Yes

2. Biogeographical Or Marine Level

2.1 Biogeographical Region

2.2 Published sources

Mediterranean (MED)

The present species assessment (fields 0.1-2.9) has been compiled by Daniele Paoloni, Cristiano Spilinga (Associazione Teriologica Italiana - ATIt) and Anna Alonzi, Piero Genovesi, Francesca Ronchi (Institute for Environmental Protection and Research - ISPRA). Information, unpublished data and experts' judgments have been provided by Paolo Agnelli, Mara Calvini, Luca Cistrone, Michele Ferretto, Danilo Russo, Dino Scaravelli, Martina Spada, Roberto Toffoli, Simone Vergari (Italian Group for bat Research).

Archivio Osservatorio Regionale per Biodiversità. Regione Umbria.

Calvini M., 2006. Monitoraggio dei chirotteri nella piana del Magra e Vallecchia (SP) (rapporto interno).

Calvini M., 2006. I Chirotteri della ZPS Beigua-Turchino e del Parco del Beigua; 70 pag. Ente Parco del Beigua, Regione Liguria.

Calvini M., 2007. Studio preliminare sulla chirotterofauna delle tre foreste demaniali del Parco dell'Aveto (rapporto interno).

Calvini M., 2007. I Chirotteri delle Alpi Liguri; 24 pag. Provincia di Imperia, Regione Liguria.

Calvini M., 2009. Indagine chirotterologica nei seguenti SIC della provincia di Savona: IT1323201, IT1324011, IT1323112 e IT1323203 (rapporto interno).

Calvini M., 2010. Monitoraggio delle colonie di chirotteri riproduttive e svernanti di particolare interesse conservazionistico note in Liguria (rapporto interno).

Database del Repertorio Naturalistico Toscano.

Regione Liguria, 2008, Carta della Biodiversità, www.ambienteinliguria.it

22/04/2014 16.13.50 Page 1 of 11

Ruffo S., Stoch F., 2005. Checklist e distribuzione della fauna italiana. Memorie del Museo Civico di storia naturale di Verona, 2.serie, Sezione scienze della Vita 16.

Spilinga C., Russo D., Carletti S., Jiménez Grijalva M.P., Sergiacomi U., Ragni B., (in stampa). Chirotteri dell'Umbria. Distribuzione geografica ed ecologica. Regione Umbria. Università degli Studi di Perugia.

Scaravelli D. & E. Guidi, Nyctalus noctula in Italia e rilievi sulla colonia riproduttiva di Cervia (RA). Riass. III Cong. Ital. Teriologia, San Remo 21-23 Settembre 2001: 100.

Scaravelli D. 2005. Observations on Nyctalus noctula breeding in Italy. Xth EBRS Galway. Bat Res. News, 46: 3.

Toffoli R., 2011. I Chirotteri del Parco Naturale delle Capanne di Marcarolo. Regione Piemonte-Parco Naturale delle Capanne di Marcarolo (rapporto interno).

2.3 Range

2.3.1 Surface area - Range (km²)

2.3.2 Method - Range surface area

2.3.3 Short-term trend period

2.3.4 Short-term trend direction

2.3.5 Short-term trend magnitude

2.3.6 Long-term trend period

2.3.7 Long-term trend direction

2.3.8 Long-term trend magnitude

2.3.9 Favourable reference range

24700

Estimate based on partial data with some extrapolation and/or modelling (2)

2001-2012

stable (0)

min max

N/A

min

max

area (km²)

operator approximately equal to (≈)

unkown

method Expert judgement

2.3.10 Reason for change

Improved knowledge/more accurate dataUse of different method

2.4 Population

2.4.1 Population size

(individuals or agreed exception)

Unit N/A

min max

2.4.2 Population size (other than individuals) Unit

number of map 10x10 km grid cells (grids10x10)

min

41 41 max

2.4.3 Additional information

Definition of locality

Conversion method

Problems Impossible to convert grids to individuals

2.4.4 Year or period

2.4.5 Method – population size

1985-2012

Estimate based on expert opinion with no or minimal sampling (1)

2.4.6 Short-term trend period

2001-2012 unknown (x)

2.4.7 Short term trend direction 2.4.8 Short-term trend magnitude

min max

confidence interval

2.4.9 Short-term trend method

Estimate based on expert opinion with no or minimal sampling (1)

2.4.10 Long-term trend period 2.4.11 Long term trend direction

N/A

22/04/2014 16.13.50 Page 2 of 11

2.4.12 Long-term trend magnitude confidence interval min max 2.4.13 Long-term trend method N/A number 2.4.14 Favourable reference population N/A operator unknown Yes method Expert judgement 2.4.15 Reason for change 2.5 Habitat for the Species 2.5.1 Surface area - Habitat (km²) 2.5.2 Year or period 2.5.3 Method used - habitat Absent data (0) 2.5.4 a) Quality of habitat Unknown 2.5.4 b) Quality of habitat - method **Expert based** 2.5.5 Short term trend period 2001-2012 2.5.6 Short term trend direction unknown (x) 2.5.7 Long-term trend period 2.5.8 Long term trend direction N/A 2.5.9 Area of suitable habitat (km²) 2.5.10 Reason for change

2.6 Main Pressures			
Pressure		ranking	pollution qualifier(s)
Forest and Plantation management &	use (B02)	high importance (H)	N/A
forestry clearance (B02.02)		high importance (H)	N/A
removal of dead and dying trees (B02.0	04)	high importance (H)	N/A
use of biocides, hormones and chemica	als (A07)	medium importance (M)	N/A
burning down (J01.01)		medium importance (M)	N/A
2.6.1 Method used – pressures	based only on ex	pert judgements (1)	

2.0.1 Method used – pressures	based only on expert Judgements (1)			
2.7 Main Threats				
Threat		ranking	pollution qualifier(s)	
Forest and Plantation management &	use (B02)	high importance (H)	N/A	
forestry clearance (B02.02)		high importance (H)	N/A	
removal of dead and dying trees (B02	04)	high importance (H)	N/A	
use of biocides, hormones and chemic	cals (A07)	medium importance (M)	N/A	
burning down (J01.01)		medium importance (M)	N/A	
wind energy production (C03.03)		medium importance (M)	N/A	

2.7.1 Method used – threats expert opinion (1)

2.8 Complementary Information

2 C Main Duasseurs

2.8.1 Justification of % thresholds for trends2.8.2 Other relevant Information

2.8.3 Trans-boundary assessment

2.9 Conclusions (assessment of conservation status at end of reporting period)

22/04/2014 16.13.50 Page 3 of 11

2.9.1 Range
2.9.2. Population
2.9.3. Habitat
2.9.4. Future prospects
2.9.5 Overall assessment of Conservation Status
2.9.5 Overall trend in

assessment Favourable (FV)
qualifiers N/A
assessment Unknown (XX)
qualifiers N/A
assessment Unknown (XX)
qualifiers N/A
assessment Unknown (XX)
qualifiers N/A
Unknown (XX)

N/A

3. Natura 2000 coverage and conservation measures - Annex II species

max

3.1 Population

Conservation Status

3.1.1 Population Size Unit N/A min

3.1.2 Method used N/A

3.1.3 Trend of population size within N/A

3.2 Conversation Measures

2. Biogeographical Or Marine Level

2.1 Biogeographical Region

2.2 Published sources

Continental (CON)

The present species assessment (fields 0.1-2.9) has been compiled by Daniele Paoloni, Cristiano Spilinga (Associazione Teriologica Italiana - ATIt) and Anna Alonzi, Piero Genovesi, Francesca Ronchi (Institute for Environmental Protection and Research - ISPRA). Information, unpublished data and experts' judgments have been provided by Paolo Agnelli, Mara Calvini, Luca Cistrone, Michele Ferretto, Danilo Russo, Dino Scaravelli, Martina Spada, Roberto Toffoli, Simone Vergari (Italian Group for bat Research).

Banca Dati Regionale Emilia Romagna (aggiornamento al 2010).

Calvini M., 2006. Monitoraggio dei chirotteri nella piana del Magra e Vallecchia (SP) (rapporto interno).

Calvini M., 2007. Studio preliminare sulla chirotterofauna delle tre foreste demaniali del Parco dell'Aveto (rapporto interno).

Calvini M., 2009. I Chirotteri del SIC IT1110022 Stagno di Oulx e IT1110020 Lago di Viverone. IPLA (rapporto interno).

Calvini M., 2010. Monitoraggio delle colonie di chirotteri riproduttive e svernanti di particolare interesse conservazionistico note in Liguria (rapporto interno). Calvini M., 2006. Monitoraggio dei chirotteri nella piana del Magra e Vallecchia (SP) (rapporto interno).

22/04/2014 16.13.50 Page 4 of 11

Calvini M., 2007. Studio preliminare sulla chirotterofauna delle tre foreste demaniali del Parco dell'Aveto (rapporto interno).

Calvini M., 2009. I Chirotteri del SIC IT1110022 Stagno di Oulx e IT1110020 Lago di Viverone. IPLA (rapporto interno).

Calvini M., 2010. Monitoraggio delle colonie di chirotteri riproduttive e svernanti di particolare interesse conservazionistico note in Liguria (rapporto interno).

Dall'Asta A., 1995-1996. Atlante preliminare dei Chirotteri (Chiroptera, Mammalia) della Regione Friuli-Venezia Giulia - Prima Sintesi Cartografica. Tesi di Laurea in Scienze Naturali, Fac. di Scienze MM. FF. NN. dell'Università degli Studi di Trieste, Relatori G. A. Amirante & S. Dolce: 1-103.

Database del Repertorio Naturalistico Toscano.

Insubria DataBat, 2012. Data base chirotteri dell'Università degli Studi dell'Insubria aggiornato al 2012.

Kryštufek B., Rešek Donev N., 2005. The Atlas of Slovenian Bats (Chiroptera). Scopolia, 55 (2005): 1-92.

Lapini L., Dall'Asta A., Dublo L., Spoto M., Venier E., 1996 (1995). Materiali per una teriofauna dell'Italia Nord - Orientale (Mammalia, Friuli-Venezia Giulia). Gortania 17: 149-248.

Lapini L., 1997. Animali in città. Anfibi, Rettili e Mammiferi del Comune di Udine. Comune di Udine, Settore Attività Culturali ed Educative ed., Udine: 1-149.

Ruffo S., Stoch F., 2005. Checklist e distribuzione della fauna italiana. Memorie del Museo Civico di storia naturale di Verona, 2.serie, Sezione scienze della Vita 16.

Sindaco R., Baratti N., Boano G., 1992. I Chirotteri del Piemonte e della Valle d'Aosta. Hystrix. (n.s.) 4 (1): 1-40.

Spada M., Preatoni G., Tosi G., Martinoli A., 2010. Piano di monitoraggio dei Vertebrati terrestri di interesse comunitario (Direttive 79/409/CEE e 92/43/CEE) in Lombardia. Il monitoraggio dei Chirotteri. Fondazione Lombardia per l'Ambiente, Università degli Studi dell'Insubria.

Toffoli R., 2009. I Chirotteri del SIC IT1160010 Bosco del Merlino (e successivi aggiornamenti). IPLA (rapporto interno).

Vigorita V., Cucè L., 2008. La fauna selvatica in Lombardia. Rapporto 2008 su distribuzione, abbondanza e stato di conservazione di uccelli e mammiferi. Regione Lombardia. Pp. 364.

2.3 Range

2.3.1 Surface area - Range (km²)

2.3.2 Method - Range surface area

2.3.3 Short-term trend period

35800

Estimate based on partial data with some extrapolation and/or modelling (2) 2001-2012

22/04/2014 16.13.50 Page 5 of 11

ii, iv alid v species (Alii	iek bj			
2.3.4 Short-term trend direction2.3.5 Short-term trend magnitude2.3.6 Long-term trend period2.3.7 Long-term trend direction	stable (0) min N/A	max		
2.3.8 Long-term trend magnitude	min	max		
2.3.9 Favourable reference range	area (km²)		hi agual ta ()	
	operator unkown	No	ly equal to (≈)	
	method	Expert judge		
2.3.10 Reason for change	Improved knowleds	ge/more accurat	te dataUse of different method	
2.4 Population				
2.4.1 Population size	Unit N/A			
(individuals or agreed exception)	min	max		
2.4.2 Population size		•	grid cells (grids10x10)	
(other than individuals)	min 87	max 87		
2.4.3 Additional information	Definition of locality			
	Conversion method Problems		a to convert gride into individuals	
2.4.4 Year or period	1985-2012	iiiibossinie	to convert grids into individuals	
2.4.5 Method – population size		expert opinion w	rith no or minimal sampling (1)	
2.4.6 Short-term trend period	2001-2012			
2.4.7 Short term trend direction	stable (0)			
2.4.8 Short-term trend magnitude2.4.9 Short-term trend method	min Estimate based on a	max	confidence interval vith no or minimal sampling (1)	
2.4.10 Long-term trend period	Littillate based off	expert opinion w		
2.4.11 Long term trend direction	N/A			
2.4.12 Long-term trend magnitude	min	max	confidence interval	
2.4.13 Long-term trend method 2.4.14 Favourable reference	N/A number			
population	operator approx	cimately equal to	0 (≈)	
	unknown No			
	•	judgement		
2.4.15 Reason for change	Improved knowledg	ge/more accurat	e data Use of different method	
2.5 Habitat for the Species				
2.5.1 Surface area - Habitat (km²)2.5.2 Year or period				
2.5.3 Method used - habitat2.5.4 a) Quality of habitat	Absent data (0) Good			
2.5.4 b) Quality of habitat - method	Expert based			
2.5.5 Short term trend period	2001-2012			
2.5.6 Short term trend direction	stable (0)			
2.5.7 Long-term trend period2.5.8 Long term trend direction	N/A			
2.5.0 Long term trend direction	N/A			

22/04/2014 16.13.50 Page 6 of 11

Improved knowledge/more accurate data Use of different method

2.5.9 Area of suitable habitat (km²)

2.5.10 Reason for change

2.6 Main Pressures			
Pressure		ranking	pollution qualifier(s)
Forest and Plantation management & use (B02)		high importance (H)	N/A
forestry clearance (B02.02)		high importance (H)	N/A
removal of dead and dying trees (B02.	04)	high importance (H)	N/A
use of biocides, hormones and chemicals (A07)		medium importance (M)	N/A
2.6.1 Method used – pressures	based only on expe	rt judgements (1)	
2.7 Main Threats			
Threat		ranking	pollution qualifier(s)
Forest and Plantation management &	use (B02)	high importance (H)	N/A
forestry clearance (B02.02)		high importance (H)	N/A
removal of dead and dying trees (B02.	04)	high importance (H)	N/A
use of biocides, hormones and chemicals (A07)		medium importance (M)	N/A
wind energy production (C03.03)		medium importance (M)	N/A
2.7.1 Method used – threats	expert opinion (1)		
2.8 Complementary Information			
2.8.1 Justification of % thresholds for trends			
2.8.2 Other relevant Information			
2.8.3 Trans-boundary assessment			
2.9 Conclusions (assessment of cor	nservation status at	end of reporting period)	
2.9.1 Range	assessment Favour qualifiers N/A	rable (FV)	
2.9.2. Population	assessment Favourable (FV)		
2.9.3. Habitat	qualifiers N/A assessment Favoui qualifiers N/A	rable (FV)	
2.9.4. Future prospects	assessment Favour qualifiers N/A	rable (FV)	
2.9.5 Overall assessment of	Favourable (FV)		

3. Natura 2000 coverage and conservation measures - Annex II species

N/A

3.1 Population

Conservation Status
2.9.5 Overall trend in

Conservation Status

3.1.1 Population Size	Unit	N/A	
	min		max
3.1.2 Method used	N/A		
3.1.3 Trend of population size within	N/A		

3.2 Conversation Measures

22/04/2014 16.13.50 Page 7 of 11

2. Biogeographical Or Marine Level

2.1 Biogeographical Region

2.2 Published sources

Alpine (ALP)

The present species assessment (fields 0.1-2.9) has been compiled by Daniele Paoloni, Cristiano Spilinga (Associazione Teriologica Italiana - ATIt) and Anna Alonzi, Piero Genovesi, Francesca Ronchi (Institute for Environmental Protection and Research - ISPRA). Information, unpublished data and experts' judgments have been provided by Paolo Agnelli, Mara Calvini, Luca Cistrone, Michele Ferretto, Danilo Russo, Dino Scaravelli, Martina Spada, Roberto Toffoli, Simone Vergari (Italian Group for bat Research).

Calvini M., 2007. I Chirotteri delle Alpi Liguri; 24 pag. Provincia di Imperia, Regione Liguria.

Calvini M., 2009. I Chirotteri del SIC IT1110022 Stagno di Oulx e IT1110020 Lago di Viverone. IPLA (rapporto interno).

Calvini M., 2009. Indagine sulla chirotterofauna nel SIC "Bric Tana-Bric Mongarda", comune di Millesimo (SV).

Calvini M., 2009. Indagine chirotterologica nei seguenti SIC della provincia di Savona: IT1323201, IT1324011, IT1323112 e IT1323203 (rapporto interno).

Calvini M., 2010. Monitoraggio delle colonie di chirotteri riproduttive e svernanti di particolare interesse conservazionistico note in Liguria (rapporto interno).

Dati AVK - Arbeitsgemeinschaft Vogelkunde Südtirol (1987-2009).

Debernardi P., Patriarca E., 2007. The bats of the Lake Maggiore Piedmont shore (NW Italy). Hystrix It. J. Mamm. (n.s.) 18 (1): 39-55.

Insubria DataBat, 2012. Data base chirotteri dell'Università degli Studi dell'Insubria aggiornato al 2012.

Spada M., Preatoni G., Tosi G., Martinoli A., 2010. Piano di monitoraggio dei Vertebrati terrestri di interesse comunitario (Direttive 79/409/CEE e 92/43/CEE) in Lombardia. Il monitoraggio dei Chirotteri. Fondazione Lombardia per l'Ambiente, Università degli Studi dell'Insubria.

Vigorita V., Cucè L., 2008. La fauna selvatica in Lombardia. Rapporto 2008 su distribuzione, abbondanza e stato di conservazione di uccelli e mammiferi. Regione Lombardia. Pp. 364.

Museo di S.N di Bolzano. Indagine sui pipistrelli dell'Alto Adige (1986-2003).

Provincia di Trento. Rilevamenti e monitoraggi popolazioni chirotteri della provincia di Trento nel periodo 1999-2012.

Ruffo S., Stoch F., 2005. Checklist e distribuzione della fauna italiana. Memorie del Museo Civico di storia naturale di Verona, 2.serie, Sezione scienze della Vita 16.

22/04/2014 16.13.50 Page 8 of 11

2.3 Range			
 2.3.1 Surface area - Range (km²) 2.3.2 Method - Range surface area 2.3.3 Short-term trend period 2.3.4 Short-term trend direction 	16700 Estimate based on partial data with some extrapolation and/or modelling (2) 2001-2012 stable (0)		
2.3.5 Short-term trend magnitude2.3.6 Long-term trend period	min	max	
2.3.7 Long-term trend direction	N/A		
2.3.8 Long-term trend magnitude2.3.9 Favourable reference range	min area (km²)	max	
2.3.9 Favourable reference range	operator	approximately equal t	.co (≈)
	unkown	No	
	method	Expert judgement	
2.3.10 Reason for change	Improved knowledge	e/more accurate dataUs	se of different method
2.4 Population			
2.4.1 Population size	Unit N/A		
(individuals or agreed exception)	min	max	
2.4.2 Population size	Unit number of i	map 10x10 km grid cells	(grids10x10)
(other than individuals)	min 35	max 35	,
2.4.3 Additional information	Definition of locality		
	Conversion method		
	Problems	Impossible to conv	ert grids into individuals
2.4.4 Year or period 2.4.5 Method – population size	1985-2012	xpert opinion with no or	e minimal campling (1)
2.4.6 Short-term trend period	2001-2012	cpert opinion with no or	milinai sampinig (1)
2.4.7 Short term trend direction	unknown (x)		
2.4.8 Short-term trend magnitude	min	max	confidence interval
2.4.9 Short-term trend method2.4.10 Long-term trend period	Estimate based on ex	opert opinion with no or	minimal sampling (1)
2.4.11 Long term trend direction	N/A		
2.4.12 Long-term trend magnitude	min	max	confidence interval
2.4.13 Long-term trend method 2.4.14 Favourable reference	N/A number		
population	operator N/A		
p o p moster.	unknown Yes method		
2.4.15 Reason for change	method		
2.5 Habitat for the Species			
2.5.1 Surface area - Habitat (km²)			
2.5.2 Year or period			
2.5.3 Method used - habitat	Absent data (0)		
2.5.4 a) Quality of habitat	Unknown		
2.5.4 b) Quality of habitat - method	Expert based		

22/04/2014 16.13.50 Page 9 of 11

2001-2012

unknown (x)

2.5.5 Short term trend period

2.5.7 Long-term trend period

2.5.6 Short term trend direction

2.5.8 Long term trend direction N/A 2.5.9 Area of suitable habitat (km²) 2.5.10 Reason for change 2.6 Main Pressures Pressure ranking pollution qualifier(s) Forest and Plantation management & use (B02) high importance (H) N/A forestry clearance (B02.02) high importance (H) N/A removal of dead and dying trees (B02.04) high importance (H) N/A use of biocides, hormones and chemicals (A07) medium importance (M) N/A 2.6.1 Method used – pressures based only on expert judgements (1) 2.7 Main Threats **Threat** ranking pollution qualifier(s) Forest and Plantation management & use (B02) high importance (H) N/A forestry clearance (B02.02) high importance (H) N/A removal of dead and dying trees (B02.04) high importance (H) N/A use of biocides, hormones and chemicals (A07) medium importance (M) N/A wind energy production (C03.03) medium importance (M) N/A 2.7.1 Method used – threats expert opinion (1) 2.8 Complementary Information

2.8.1 Justification of % thresholds for trends2.8.2 Other relevant Information

2.8.3 Trans-boundary assessment

2.9 Conclusions (assessment of conservation status at end of reporting period)

2.9.1 Range

assessment Favourable (FV)
qualifiers N/A

2.9.2. Population

assessment Unknown (XX)
qualifiers N/A

2.9.3. Habitat assessment Unknown (XX)

qualifiers N/A

assessment Unknown (XX)

qualifiers N/A

Unknown (XX)

2.9.5 Overall assessment of

Conservation Status

2.9.4. Future prospects

2.9.5 Overall trend in Conservation Status

N/A

3. Natura 2000 coverage and conservation measures - Annex II species

3.1 Population

3.1.1 Population Size Unit N/A

min max

22/04/2014 16.13.50 Page 10 of 11

3.1.2 Method used	N/A
3.1.3 Trend of population size within	N/A

3.2 Conversation Measures

22/04/2014 16.13.50 Page 11 of 11