0.1 Member State	Π
0.2.1 Species code	2029
0.2.2 Species name	Globicephala melas
0.2.3 Alternative species scientific name	N/A
0.2.4 Common name	Globicefalo

1. National Level

1.1 Maps

1.1.1 Distribution Map	Yes
1.1.1a Sensitive species	No
1.1.2 Method used - map	Estimate based on partial data with some extrapolation and/or modelling (2)
1.1.3 Year or period	2010-2011
1.1.4 Additional map	No
1.1.5 Range map	Yes

2. Biogeographical Or Marine Level

2.1 Biogeographical Region

2.2 Published sources

Marine Mediterranean (MMED)

The present species assessment (fields 0.1-2.9) has been compiled by Anna Alonzi, Piero Genovesi, Francesca Ronchi (ISPRA). Information and data have been extracted from MSFD Supporting document on the Initial Assessment on Cetaceans, including methodology, data used and results (ISPRA,2013). Contributing authors: Caterina Fortuna, Mario Acquarone, Aldo Annunziatellis, Antonella Arcangeli, Arianna Azzellino, Nicola Baccetti, Michela Bellingeri, Silvia Bonizzoni, Junio Fabrizio Borsani, Ilaria Caliani, Simonepietro Canese, Roberta Canneri, Nadia Cerioli, Andrea De Lucia, Salvatore Dimatteo, Carmelo Fanizza, Elio Filidei jr., Maria Cristina Fossi, Fulvio Garibaldi, Stefania Gaspari, Otello Giovanardi, Michela Giusti, Guido Gnone, Paolo Guidetti, Drasko Holcer, Giancarlo Lauriano, Letizia Marsili, Antonio Mazzola, Giulia Mo, Aurelie Moulins, Barbara Mussi, Giuseppe

Notarbartolo di Sciara, Lidia Orsi Relini, Daniela Silvia Pace, Simone Panigada, Gianni Pavan, Michela Podestà, Marina Pulcini, Sasa Raicevich, Ettore Randi, Teresa Romeo, Massimiliano Rosso, Antonello Sala, Paola Tepsich, Walter Zimmer e Nicola Zizzo. Expert judgements have been provided by Caterina Fortuna (ISPRA).

A.A.V.V. 2007. Workshop d'inquadramento del fenomeno delle interazioni tra cetacei e attività di pesca in Italia. Roma, 10-11 settembre 2007, Ministero delle politiche agricole, alimentari e forestali, Roma. Progetto «Programma Bycatch Italia-ACCOBAMS» (codice 6G23), finanziato dalla Direzione Generale per la Pesca marittima e Acquacoltura, MiPAAF.

Azzellino, A., Airoldi, S., Gaspari, S., Nani, B., 2008. Habitat use of cetaceans along the continental slope and Adjacent waters in the western Ligurian Sea. Deep Sea Res. 55, 296-323.

Azzellino, A., Lanfredi, C., D'Amico, A., Pavan, G., Podestà, M., Haun, J., 2011. Risk mapping for sensitive species to underwater anthropogenic sound emissions: model development and validation in two Mediterranean areas. Mar. Pollut. Bull. 63, 56-70.

Azzellino, A., Panigada, S., Lanfredi, C., Zanardelli, M., Airoldi, S., Notarbartolo di Sciara, G. 2012. Predictive habitat models for managing marine areas: Spatial and

30/04/2014 10.58.23 Page 1 of 5

temporal distribution of marine mammals within the Pelagos Sanctuary (Northwestern Mediterranean sea). Ocean & Coastal Management 67 (2012) 63-74.

Buckland ST, Anderson DR, Burnham KP, Laake JL, Borchers DL, Thomas L 2001. Introduction to distance sampling: estimating abundance of biological populations. Oxford University Press, Oxford.

Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., Borchers, D.L., Thomas, L. 2004. Advanced Distance Sampling. Oxford University Press.

Cañadas, A., R. Sagarminaga, R. de Stephanis, E. Urquiola and P.S. Hammond. 2005. Habitat selection models as a conservation tool: proposal of marine protected areas for cetaceans in Southern Spain. Aquatic Conservation: Marine and Freshwater Ecosystems 15:495-521.

Cañadas, A., Sagarminaga, R. and García-Tiscar, S. 2002. Cetacean distribution related with depth and slope in the Mediterranean waters off southern Spain. Deep Sea Research I 49: 2053-2073.

Caughley G. 1966. Mortality patterns in mammals. Ecology 47:906–918. Cerioni, S., Forni, L., Lo Tenero, A., Nannarelli, S. & Pulcini, M. 1996. A cetacean survey in the Taranto Gulf: work in progress. European Research on Cetaceans 9:102–103.

D'Ortenzio, F. and Ribera d'Alcala, M. 2008. On the trophic regimes of the Mediterranean Sea. Biosciences Discussions 5: 2959-2983.

Emelen J.M. 1970. Age specificity and ecological theory. Ecology, 51, 588–601. Fortuna, C.M., Filidei, E. jr. 2011. Annual Report on the implementation of Council Regulation (EC) 812/2004 - 2010. Rapporto tecnico preparato per il Ministero delle politiche agricole, alimentari e forestali, 10 pagine.

Fortuna, C.M., Holcer, D., Filidei, E. jr, Tunesi, L., 2011 Relazione finale del progetto "Valutazione dell'impatto della mortalità causata da attività di pesca su Cetacei e tartarughe marine in Adriatico: primo survey per la stima dell'abbondanza" (Prot. MIPAAF DG PEMAC n. 1690 del 10/02/2010 e al Prot. MATTM DPN n. 27623 del 23/12/2009), 51 pagine + Allegati.

Fortuna, C.M., Vallini, C., Filidei, E. jr, Ruffino, M., Consalvo, I., Di Muccio, S., Gion, C., Scacco, U., Tarulli, E., Giovanardi, O., Mazzola, A. 2010. Bycatch of cetaceans and other species of conservation concern during pair trawl fishing operations in the Adriatic Sea (Italy). Chemistry and Ecology 26(Supplement):65-76.

Gannier A 1998. Les cétacés de Méditerranée nord-occidentale: nouveaux résultats sur leur distribution, la structure de leur peuplement et l'abondance relative des différentes espèces. Mésogée 56: 3-19.

Gannier A 2006. Le peuplement estival de cetaces dans le Sanctuaire Marin Pelagos (Mediterranee nord-occidentale): distribution et abondance. Mammalia 70: 17-27.

Lauriano, G., Panigada, S., Fortuna, C.M., Holcer, D., Filidei, E. jr, Pierantonio, N. and Donovan, G. 2011b. Monitoring density and abundance of cetaceans in the seas around Italy through aerial survey: a contribution to conservation and the future ACCOBAMS Survey. International Whaling Commission SC/63/SM6. Marini, L., Consiglio, C., Angradi, A.M., Catalano, B., Sanna, A., Valentini, T., Finoia, M.G. & Villetti, G. 1996. Distribution, abundance and seasonality of cetaceans sighted during scheduled ferry crossings in the central Tyrrhenian Sea: 1989–1992. Italian Journal of Zoology 63:381–388.

Marsili, L. (2000) Lipophilic contaminants in marine mammals: review of the results of ten years' work at the Department of environmental biology, Siena University (Italy). International Journal of Environmental Pollution 13:416–452. Politi, E., Bearzi, M., Notarbartolo di Sciara, G., Cussino, E. & Gnone, G. 1992.

30/04/2014 10.58.23 Page 2 of 5

Distribution and frequency of cetaceans in the waters adjacent to the Greek Ionian Islands. European Research on Cetaceans 6:75–78

RAC/SPA. 1998. Report of the Meeting of Experts on the Implementation of the Action Plans for Marine Mammals (Monk seal and Cetaceans) adopted within MAP. UNEP(OCA)/MED WG. 146/5, Tunis, annex 5, pp:55-57.

Rallo, G. 1979. I cetacei dell'Adriatico. WWF Veneto, 4, 3-4.

2.3 Range

2.3.1 Surface area - Range (km²)

2.3.2 Method - Range surface area

2.3.3 Short-term trend period

2.3.4 Short-term trend direction

2.3.5 Short-term trend magnitude

2.3.6 Long-term trend period

2.3.7 Long-term trend direction

2.3.8 Long-term trend magnitude

2.3.9 Favourable reference range

45000

Estimate based on partial data with some extrapolation and/or modelling (2)

2000-2011

unknown (x)

min max

N/A

min max

area (km²)

N/A operator unkown Yes

method

Use of different method 2.3.10 Reason for change

2.4 Population

2.4.1 Population size

(individuals or agreed exception)

2.4.2 Population size

(other than individuals)

Unit N/A

min max

Unit number of map 10x10 km grid cells (grids10x10)

min 150 max 150

2.4.3 Additional information **Definition of locality**

Conversion method

Problems It is not possible o convert grids into individuals

2.4.4 Year or period

2.4.5 Method – population size

2.4.6 Short-term trend period

2.4.7 Short term trend direction

2.4.8 Short-term trend magnitude

2.4.9 Short-term trend method

2.4.10 Long-term trend period

2.4.11 Long term trend direction

2.4.12 Long-term trend magnitude

2.4.13 Long-term trend method

2.4.14 Favourable reference

population

2010-2011

Estimate based on partial data with some extrapolation and/or modelling (2)

2000-2011

unknown (x)

confidence interval min max

Absent data (0)

N/A

min

confidence interval max

N/A number

operator N/A Yes

unknown method

Use of different method 2.4.15 Reason for change

2.5 Habitat for the Species

2.5.1 Surface area - Habitat (km²)

30/04/2014 10.58.23 Page 3 of 5

2.5.2 Year or period	
2.5.3 Method used - habitat	Absent data (0)
2.5.4 a) Quality of habitat	Unknown
2.5.4 b) Quality of habitat - method	Expert opinion
2.5.5 Short term trend period	
2.5.6 Short term trend direction	unknown (x)
2.5.7 Long-term trend period	
2.5.8 Long term trend direction	N/A
2.5.9 Area of suitable habitat (km²)	
2.5.10 Reason for change	Use of different method
2.6 Main Prossures	

2.6 Main Pressures			
Pressure		ranking	pollution qualifier(s)
Marine water pollution (H03)		high importance (H)	Mixed pollutants (X)
netting (F02.01.02)		low importance (L)	N/A
2.6.1 Method used – pressures	mainly based on exp	pert judgement and other da	ata (2)
2.7 Main Threats			
Threat		ranking	pollution qualifier(s)
Marine water pollution (H03)		high importance (H)	Mixed pollutants (X)
netting (F02.01.02)		low importance (L)	N/A
2.7.1 Method used – threats	expert opinion (1)		
2.8 Complementary Information			

2.8.1 Justification of % thresholds for trends

2.8.2 Other relevant Information This pelagic species is present in very low numbers. Its distribution is currently unknown.

2.8.3 Trans-boundary assessment

2.9 Conclusions (assessment of conservation status at end of reporting period)

2.9.1 Kange	qualifiers N/A
2.9.2. Population	assessment Unknown (XX) qualifiers N/A
2.9.3. Habitat	assessment Unknown (XX) qualifiers N/A
2.9.4. Future prospects	assessment Unknown (XX) qualifiers N/A
2.9.5 Overall assessment of Conservation Status	Unknown (XX)
2.9.5 Overall trend in Conservation Status	N/A

3. Natura 2000 coverage and conservation measures - Annex II species

3.1 Population

30/04/2014 10.58.23 Page 4 of 5

3.1.1 Population Size	Unit min	N/A	max	
3.1.2 Method used	N/A			
3.1.3 Trend of population size within	N/A			
3.2 Conversation Measures				

30/04/2014 10.58.23 Page 5 of 5