0.1 Member State	IT
0.2.1 Species code	1350
0.2.2 Species name	Delphinus delphis
0.2.3 Alternative species scientific name	N/A
0.2.4 Common name	Delfino comune

1. National Level

1.1 Maps

1.1.1 Distribution Map	Yes
1.1.1a Sensitive species	No
1.1.2 Method used - map	Estimate based on partial data with some extrapolation and/or modelling (2)
1.1.3 Year or period	2010-2011
1.1.4 Additional map	No
1.1.5 Range map	Yes

2. Biogeographical Or Marine Level

2.1 Biogeographical Region

2.2 Published sources

Marine Mediterranean (MMED)

The present species assessment (fields 0.1-2.9) has been compiled by Anna Alonzi, Piero Genovesi, Francesca Ronchi (ISPRA). Information and data have been extracted from MSFD Supporting document on the Initial Assessment on Cetaceans, including methodology, data used and results (ISPRA,2013). Contributing authors: Caterina Fortuna, Mario Acquarone, Aldo Annunziatellis, Antonella Arcangeli, Arianna Azzellino, Nicola Baccetti, Michela Bellingeri, Silvia Bonizzoni, Junio Fabrizio Borsani, Ilaria Caliani, Simonepietro Canese, Roberta Canneri, Nadia Cerioli, Andrea De Lucia, Salvatore Dimatteo, Carmelo Fanizza, Elio Filidei jr., Maria Cristina Fossi, Fulvio Garibaldi, Stefania Gaspari, Otello Giovanardi, Michela Giusti, Guido Gnone, Paolo Guidetti, Drasko Holcer, Giancarlo Lauriano, Letizia Marsili, Antonio Mazzola, Giulia Mo, Aurelie Moulins, Barbara Mussi, Giuseppe

Notarbartolo di Sciara, Lidia Orsi Relini, Daniela Silvia Pace, Simone Panigada, Gianni Pavan, Michela Podestà, Marina Pulcini, Sasa Raicevich, Ettore Randi, Teresa Romeo, Massimiliano Rosso, Antonello Sala, Paola Tepsich, Walter Zimmer e Nicola Zizzo. Expert judgements have been provided by Caterina Fortuna (ISPRA).

A.A.V.V. 2007. Workshop d'inquadramento del fenomeno delle interazioni tra cetacei e attività di pesca in Italia. Roma, 10-11 settembre 2007, Ministero delle politiche agricole, alimentari e forestali, Roma. Progetto «Programma Bycatch Italia-ACCOBAMS» (codice 6G23), finanziato dalla Direzione Generale per la Pesca marittima e Acquacoltura, MiPAAF.

Azzellino, A., Airoldi, S., Gaspari, S., Nani, B., 2008. Habitat use of cetaceans along the continental slope and adjacent waters in the western Ligurian Sea. Deep Sea Res. 55, 296-323.

Azzellino, A., Lanfredi, C., D'Amico, A., Pavan, G., Podestà, M., Haun, J., 2011. Risk mapping for sensitive species to underwater anthropogenic sound emissions: model development and validation in two Mediterranean areas. Mar. Pollut. Bull. 63, 56-70.

Azzellino, A., Panigada, S., Lanfredi, C., Zanardelli, M., Airoldi, S., Notarbartolo di Sciara, G. 2012. Predictive habitat models for managing marine areas: Spatial and

30/04/2014 10.52.54 Page 1 of 5

temporal distribution of marine mammals within the Pelagos Sanctuary (Northwestern Mediterranean sea). Ocean & Coastal Management 67 (2012) 63-74.

Bearzi, G., Holcer, D., Notarbartolo di Sciara, G. 2004. The role of historical dolphin takes and habitat degradation in shaping the present status of northern Adriatic cetaceans. Aquatic Conservation: Marine and Freshwater Ecosystems 14:363–379.

Bearzi, G., Reeves, R.R., Notarbartolo di Sciara, G., Politi, E., Cañadas, A., Frantzis, A. and Mussi, B. 2003. Ecology, status and conservation of Short-beaked Common Dolphins (Delphinus delphis) in the Mediterranean Sea. Mammal Review. 33(34): 224-252.

Buckland ST, Anderson DR, Burnham KP, Laake JL, Borchers DL, Thomas L 2001. Introduction to distance sampling: estimating abundance of biological populations. Oxford University Press, Oxford.

Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., Borchers, D.L., Thomas, L. 2004. Advanced Distance Sampling. Oxford University Press.

Cañadas, A., Hammond, P.S. 2008. Abundance and habitat preferences of the short-beaked common dolphin (Delphinus delphis) in the South-western Mediterranean: implications for conservation. Endangered Species Research 4:309-331.

Cañadas, A., R. Sagarminaga, R. de Stephanis, E. Urquiola and P.S. Hammond. 2005. Habitat selection models as a conservation tool: proposal of marine protected areas for cetaceans in Southern Spain. Aquatic Conservation: Marine and Freshwater Ecosystems 15:495-521.

Cañadas, A., Sagarminaga, R. and García-Tiscar, S. 2002. Cetacean distribution related with depth and slope in the Mediterranean waters off southern Spain. Deep Sea Research I 49: 2053-2073.

Caughley G. 1966. Mortality patterns in mammals. Ecology 47:906–918. Cerioni, S., Forni, L., Lo Tenero, A., Nannarelli, S. & Pulcini, M. 1996. A cetacean survey in the Taranto Gulf: work in progress. European Research on Cetaceans 9:102–103.

D'Ortenzio, F. and Ribera d'Alcala, M. 2008. On the trophic regimes of the Mediterranean Sea. Biosciences Discussions 5: 2959-2983.

Emelen J.M. 1970. Age specificity and ecological theory. Ecology, 51, 588–601. Forcada, J. and Hammond, P.S. 1998. Geographical variation in abundance of striped and common dolphins of the western Mediterranean. J. Sea. Res. 39: 313–25.

Fortuna, C.M., Filidei, E. jr. 2011. Annual Report on the implementation of Council Regulation (EC) 812/2004 - 2010. Rapporto tecnico preparato per il Ministero delle politiche agricole, alimentari e forestali, 10 pagine. Fortuna, C.M., Holcer, D., Filidei, E. jr, Tunesi, L., 2011 Relazione finale del progetto "Valutazione dell'impatto della mortalità causata da attività di pesca su Cetacei e tartarughe marine in Adriatico: primo survey per la stima dell'abbondanza" (Prot. MIPAAF DG PEMAC n. 1690 del 10/02/2010 e al Prot. MATTM DPN n. 27623 del 23/12/2009), 51 pagine + Allegati. Fortuna, C.M., Vallini, C., Filidei, E. jr, Ruffino, M., Consalvo, I., Di Muccio, S., Gion, C., Scacco, U., Tarulli, E., Giovanardi, O., Mazzola, A. 2010. Bycatch of cetaceans and other species of conservation concern during pair trawl fishing operations in the Adriatic Sea (Italy). Chemistry and Ecology 26(Supplement):65-76.

Gannier A 1998. Les cétacés de Méditerranée nord-occidentale: nouveaux résultats sur leur distribution, la structure de leur peuplement et l'abondance relative des différentes espèces. Mésogée 56: 3-19.

30/04/2014 10.52.54 Page 2 of 5

Gannier A 2006. Le peuplement estival de cetaces dans le Sanctuaire Marin Pelagos (Mediterranee nord-occidentale): distribution et abondance. Mammalia 70: 17-27.

Lauriano, G., Panigada, S., Fortuna, C.M., Holcer, D., Filidei, E. jr, Pierantonio, N. and Donovan, G. 2011. Monitoring density and abundance of cetaceans in the seas around Italy through aerial survey: a contribution to conservation and the future ACCOBAMS Survey. International Whaling Commission SC/63/SM6. Marsili, L. (2000) Lipophilic contaminants in marine mammals: review of the results of ten years' work at the Department of environmental biology, Siena University (Italy). International Journal of Environmental Pollution 13:416–452. Orsi Relini L. 2000. The Cetacean Sanctuary in the Ligurian Sea: a further reason. Biol. Mar. Mediterr., 7 (3): 117-126.

Panigada, S., Lauriano, G., Burt, L., Pierantonio, N., Donovan, G. 2011. Monitoring winter and summer abundance of cetaceans in the Pelagos Sanctuary (Northwestern Mediterranean Sea) through aerial surveys. Plos one 6(7): e22878. Doi:10.1371/journalpone.0022878

Politi, E., Bearzi, M., Notarbartolo di Sciara, G., Cussino, E. & Gnone, G. 1992. Distribution and frequency of cetaceans in the waters adjacent to the Greek Ionian Islands. European Research on Cetaceans 6:75–78

RAC/SPA. 1998. Report of the Meeting of Experts on the Implementation of the Action Plans for Marine Mammals (Monk seal and Cetaceans) adopted within MAP. UNEP(OCA)/MED WG. 146/5, Tunis, annex 5, pp:55-57.

Rallo, G. 1979. I cetacei dell'Adriatico. WWF Veneto, 4, 3-4.

2.3 Range

2.3.1 Surface area - Range (km²)

2.3.2 Method - Range surface area

2.3.3 Short-term trend period

2.3.4 Short-term trend direction

2.3.5 Short-term trend magnitude

2.3.6 Long-term trend period

2.3.7 Long-term trend direction

2.3.8 Long-term trend magnitude

2.3.9 Favourable reference range

105000

Estimate based on partial data with some extrapolation and/or modelling (2)

2000-2011

unknown (x)

min max

N/A

min max

area (km²)

operator N/A unkown Yes

method

2.3.10 Reason for change

Use of different method

2.4 Population

2.4.1 Population size

(individuals or agreed exception)

Unit N/A

min max

2.4.2 Population size (other than individuals)

Unit number

number of map 10x10 km grid cells (grids10x10)

min 650 max 650

2.4.3 Additional information

Definition of locality

Conversion method

Problems it is not possible to convert grids into individuals

2.4.4 Year or period

2.4.5 Method – population size

2.4.6 Short-term trend period

2010-2011

Estimate based on partial data with some extrapolation and/or modelling (2)

2000-2011

30/04/2014 10.52.54 Page 3 of 5

2.4.7 Short term trend direction2.4.8 Short-term trend magnitude2.4.9 Short-term trend method2.4.10 Long-term trend period	unknown(min Absent dat		max	confidence interval
2.4.11 Long term trend direction 2.4.12 Long-term trend magnitude 2.4.13 Long-term trend method 2.4.14 Fayourable reference	N/A min N/A number		max	confidence interval
population	operator unknown method	N/A Yes		
2.4.15 Reason for change	Improved k	knowledge,	/more accurat	te data Use of different method

2.5 Habitat for the Species

2.5.1 Surface area - Habitat (km²)

2.5.2 Year or period

2.5.3 Method used - habitat

2.5.4 a) Quality of habitat

2.5.4 b) Quality of habitat - method

2.5.5 Short term trend period

2.5.6 Short term trend direction

2.5.7 Long-term trend period

2.5.8 Long term trend direction

2.5.9 Area of suitable habitat (km²)

2.5.10 Reason for change

Absent data (0)

Unknown

Expert opinion

N/A

N/A

2.6 Main Pressures

Pressure	ranking	pollution qualifier(s)
netting (F02.01.02)	high importance (H)	N/A
Marine water pollution (H03)	medium importance (M)	Mixed pollutants (X)
reduction of prey availability (including carcasses) (J03.01.01)	low importance (L)	N/A

2.6.1 Method used – pressures	mainly based on expert judgement and other data (2)

2.7 Main Threats

Threat	ranking	pollution qualifier(s)
netting (F02.01.02)	high importance (H)	N/A
Marine water pollution (H03)	medium importance (M)	Mixed pollutants (X)
reduction of prey availability (including carcasses) (J03.01.01)	low importance (L)	N/A

2.7.1 Method used – threats expert opinion (1)

2.8 Complementary Information

2.8.1 Justification of % thresholds for trends

2.8.2 Other relevant Information

This species is present in very low numbers. Currently, it is regularly present around the Pontinian and Flegrean archipelagos, the Pelagian Islands and south Sicily. Very rare in Adriatic sea. Here historical data suggest current geographical extinction.

2.8.3 Trans-boundary assessment

30/04/2014 10.52.54 Page 4 of 5

2.9 Conclusions (assessment of co	onservation status at end of reporting period)			
2.9.1 Range	assessment Unknown (XX) qualifiers N/A			
2.9.2. Population	assessment Unknown (XX) qualifiers N/A			
2.9.3. Habitat	assessment Unknown (XX) qualifiers N/A			
2.9.4. Future prospects	assessment Unknown (XX) qualifiers N/A			
2.9.5 Overall assessment of Conservation Status	Unknown (XX)			
2.9.5 Overall trend in Conservation Status	N/A			
3. Natura 2000 coverage and conservation measures - Annex II species				

3.1 Population				
3.1.1 Population Size	Unit min	N/A	max	
3.1.2 Method used3.1.3 Trend of population size within	N/A N/A			
3.2 Conversation Measures				

30/04/2014 10.52.54 Page 5 of 5

Notes

Species name: Delphinus o	delphis (1350) Region code: MMED	
Field label	Note	User
2.4.7 Short term trend direction	There are not sufficient data to infer trends.	ISPRA_ AUNA
2.3.4 Range Trend	There are not sufficient data to infer trends.	ISPRA_ AUNA

30/04/2014 10.52.31 Page 1