0.1 Member State	IT
0.2.1 Species code	1310
0.2.2 Species name	Miniopterus schreibersii
0.2.3 Alternative species scientific name	N/A
0.2.4 Common name	N/A

1. National Level

1.1 Maps

1.1.1 Distribution Map	Yes
1.1.1a Sensitive species	No
1.1.2 Method used - map	Estimate based on partial data with some extrapolation and/or modelling (2)
1.1.3 Year or period	1985-2012
1.1.4 Additional map	No
1.1.5 Range map	Yes

2. Biogeographical Or Marine Level

2.1 Biogeographical Region

2.2 Published sources

Mediterranean (MED)

The present species assessment (fields 0.1-2.9) has been compiled by Daniele Paoloni, Cristiano Spilinga (Associazione Teriologica Italiana - ATIt) and Anna Alonzi, Piero Genovesi, Francesca Ronchi (Institute for Environmental Protection and Research - ISPRA). Information, unpublished data and experts' judgments have been provided by Paolo Agnelli, Mara Calvini, Luca Cistrone, Michele Ferretto, Danilo Russo, Dino Scaravelli, Martina Spada, Roberto Toffoli, Simone Vergari (Italian Group for bat Research).

Distribution data for the following Nature 2000 sites have been inserted by the Ministry of Environment (source: Italian Nature 2000 database): IT5220018; IT5220022; IT5220016; IT8030014; IT8030020.

Distribution data for the following grid cells have been removed by the Ministry of Environment: 10kmE424N193; 10kmE421N193.

Distribution data for the following grid cells have been inserted by the Ministry of Environment: 10kmE418N186; 10kmE417N193; 10kmE422N197

Archivio Osservatorio Regionale per Biodiversità. Regione Umbria.

Bux M., Russo D. e Scillitani G. 2003. La chirotterofauna della Puglia. Hystrix, It. J. Mamm. (n. s.) supp.: 150.

Calvini M., 2006. Monitoraggio dei chirotteri nella piana del Magra e Vallecchia (SP) (rapporto interno).

Calvini M., 2006. I Chirotteri della ZPS Beigua-Turchino e del Parco del Beigua; 70 pag. Ente Parco del Beigua, Regione Liguria.

Calvini M., 2007. Studio preliminare sulla chirotterofauna delle tre foreste demaniali del Parco dell'Aveto (rapporto interno).

Calvini M., 2007. I Chirotteri delle Alpi Liguri; 24 pag. Provincia di Imperia,

22/04/2014 11.59.31 Page 1 of 12

Regione Liguria.

Calvini M., 2009. Indagine chirotterologica nei seguenti SIC della provincia di Savona: IT1323201, IT1324011, IT1323112 e IT1323203 (rapporto interno).

Calvini M., 2010. Monitoraggio delle colonie di chirotteri riproduttive e svernanti di particolare interesse conservazionistico note in Liguria (rapporto interno).

Capizzi et al. (2012) Progetto atlante dei Mammiferi del Lazio - Regione Lazio – ARP.

Database del Repertorio Naturalistico Toscano.

Fornasari L., Bani L., De Carli E., Gori E., Farina F., Violani C. & Zava B., 1999. Dati sulla distribuzione geografica e ambientale di Chirotteri nell'Italia continentale e peninsulare. Atti I° Conv. Ital. Sui Chirotteri (1999): 63-81.

Loy A., De Lisio L., Capula M., Ciucci P., Russo D., Sciarretta A., 2012. Rapporto finale - Convenzione stipulata tra la Regione Molise e la Unione Zoologica Italiana per la realizzazione dei piani di gestione dei Siti Natura 2000.n. 1393/2008. Unione Zoologica Italiana, Regione Molise.

Mancini M, Scaravelli D., Pellegrini M., 2003. Check-list, status e conservazione dei mammiferi in Molise e aree limitrofe. Hystrix (N.S.) suppl. 2003: 194.

Marsico A., 1999. Contributo alla conoscenza della chirotterofauna pugliese. Dati biogeografici, fenologici e morfologici. Tesi di laurea i Zoologia dei Vertebrati, Corso di laurea in Scienze Naturali, Univ. Di Bari.

Mucedda M., Murittu G., Oppes A., Pidinchedda E., 1995. Osservazioni sui Chirotteri troglofili della Sardegna. Boll. Soc. Sarda Sci. Nat., 30: 97-129.

Mucedda M., Bertelli M. L., Pidinchedda E., 1996. Note su Miniopterus schreibersi (Chiroptera, Vespertilionidae) della Sardegna. Boll. Gruppo Spel. Sassarese, 16: 52-54.

Mucedda M., Bertelli M. L., Pidinchedda E., 2001. Note su Myotis capaccinii (Chiroptera, Vespertilionidae) della Sardegna. Boll. Gruppo Spel. Sassarese, 18: 35-38.

Mucedda M., Vadacca M. e Ciccarese N. 2003. Osservazioni sui Chirotteri di alcune grotte costiere del Salento sud-orientale (Lecce). Thalassia Salentina 26: 237-240.

Regione Autonoma della Sardegna - Assessorato Difesa Ambiente, 2012 - "Servizio di monitoraggio dello stato di conservazione degli habitat e delle specie di importanza comunitaria presenti nei siti della Rete Natura 2000 in Sardegna – Linea 4. Redazione del Rapporto sullo stato di conservazione di habitat e specie.

Regione Autonoma della Sardegna - Assessorato Difesa Ambiente - 2008-2009. "Realizzazione del sistema di monitoraggio dello stato di conservazione degli habitat e delle specie di interesse comunitario della Regione Autonoma della Sardegna".

22/04/2014 11.59.31 Page 2 of 12

Regione Liguria, 2008, Carta della Biodiversità, www.ambienteinliguria.it

Ruffo S., Stoch F., 2005. Checklist e distribuzione della fauna italiana. Memorie del Museo Civico di storia naturale di Verona, 2.serie, Sezione scienze della Vita 16.

Scaravelli D. e Bertozzi M., 2001. Nota sui Chirotteri e micromammiferi delle gravine materane. Abstract III Conv. Ital. Di Teriologia.

Scaravelli D., G. Garofalo, M. Bertozzi, G. Aloise, M. Ruedi & M. Cagnin, 2002. Primi risultati del Progetto Chirotteri della Calabria. 63° Conv. Dell'Unione Zoologica Italiana, Cosenza, 22-26 settembre 2002: 91-92.

Spilinga C., Russo D., Carletti S., Jiménez Grijalva M.P., Sergiacomi U., Ragni B., (in stampa). Chirotteri dell'Umbria. Distribuzione geografica ed ecologica. Regione Umbria. Università degli Studi di Perugia.

2.3 Range

2.3.1 Surface area - Range (km²) 126200

2.3.2 Method - Range surface area Estimate based on partial data with some extrapolation and/or modelling (2)

2.3.3 Short-term trend period 2001-2012 2.3.4 Short-term trend direction stable (0)

2.3.5 Short-term trend magnitude min max

2.3.6 Long-term trend period

2.3.7 Long-term trend direction N/A

2.3.8 Long-term trend magnitude min max

2.3.9 Favourable reference range area (km²)

operator approximately equal to (≈)

unkown No

method Expert judgement

2.3.10 Reason for change Improved knowledge/more accurate dataUse of different method

2.4 Population

2.4.1 Population size Unit N/A

(individuals or agreed exception) min max

2.4.2 Population size Unit number of map 10x10 km grid cells (grids10x10)

(other than individuals) min 250 max 250

2.4.3 Additional information Definition of locality

Conversion method

Problems Impossible to convert grids to individuals

2.4.4 Year or period 1985-2012

2.4.5 Method – population size Estimate based on expert opinion with no or minimal sampling (1)

2.4.6 Short-term trend period 2001-2012

2.4.7 Short term trend direction decrease (-)

2.4.8 Short-term trend magnitude min max confidence interval

2.4.9 Short-term trend method Estimate based on expert opinion with no or minimal sampling (1)

2.4.11 Long term trend direction N/A

2.4.10 Long-term trend period

22/04/2014 11.59.31 Page 3 of 12

ii, it and t species (iiii	ick b _j				
2.4.12 Long-term trend magnitude	min	max		confidence i	nterval
2.4.13 Long-term trend method	N/A				
2.4.14 Favourable reference	number	magaa than (s.)			
population	operator unknown	more than (>) No			
	method	Expert judgeme	ent		
2.4.15 Reason for change	Improved k	nowledge/more a		se of differen	t method
2.5 Habitat for the Species					
2.5.1 Surface area - Habitat (km²)					
2.5.2 Year or period					
2.5.3 Method used - habitat	Absent dat	a (0)			
2.5.4 a) Quality of habitat	Moderate				
2.5.4 b) Quality of habitat - method	Expert base				
2.5.5 Short term trend period2.5.6 Short term trend direction	2001-2012 stable (0)				
2.5.7 Long-term trend period	stable (u)				
2.5.8 Long term trend direction	N/A				
2.5.9 Area of suitable habitat (km²)	•				
2.5.10 Reason for change	Improved k	knowledge/more	accurate data U	se of differen	t method
2.6 Main Pressures					
Pressure		rankin	g	pol	lution qualifier(s)
Mining and quarrying (C01)		high ir	mportance (H)	N/A	4
speleology (G01.04.02)		high ir	mportance (H)	N/A	4
recreational cave visits (G01.04.03)		high ir	mportance (H)	N/A	4
wind energy production (C03.03)		high ir	mportance (H)	N/A	4
closures of caves or galleries (G05.08)		high ir	mportance (H)	N/A	4
2.6.1 Method used – pressures	based only	on expert judgen	nents (1)		
2.7 Main Threats					
Threat		rankin	g	pol	lution qualifier(s)
Mining and quarrying (C01)		high ir	mportance (H)	N/A	4
speleology (G01.04.02)		high ir	mportance (H)	N/A	4
recreational cave visits (G01.04.03)		high ir	mportance (H)	N/A	4
wind energy production (C03.03)		high ir	mportance (H)	N/A	4
closures of caves or galleries (G05.08)		high ir	mportance (H)	N/A	4
2.7.1 Method used – threats	expert opir	nion (1)			
2.8 Complementary Information					
2.8.1 Justification of % thresholds for trends					
2.8.2 Other relevant Information					

2.9 Conclusions (assessment of conservation status at end of reporting period)

2.8.3 Trans-boundary assessment

22/04/2014 11.59.31 Page 4 of 12

2.9.1 Range2.9.2. Population2.9.3. Habitat2.9.4. Future prospects

2.9.5 Overall assessment of Conservation Status
2.9.5 Overall trend in

assessment Favourable (FV)
qualifiers N/A
assessment Inadequate (U1)
qualifiers N/A
assessment Inadequate (U1)
qualifiers N/A
assessment Bad (U2)
qualifiers N/A

declining (-)

Bad (U2)

3. Natura 2000 coverage and conservation measures - Annex II species

3.1 Population

Conservation Status

3.1.1 Population Size

Unit N/A

min max

3.1.2 Method used

Absent data (0)

N/A

3.2 Conversation Measures

3.2.1 Measure	3.2.2 Type	3.2.3 Ranking	3.2.4 Location	3.2.5 Broad Evaluation
Adapt forest management (3.2)	Administrative	medium importance (M)	Both	Maintain Long term
Establish protected areas/sites (6.1)	Legal	high importance (H)	Inside	Long term Unknown
Legal protection of habitats and species (6.3)	Legal	medium importance (M)	Both	Maintain Unknown
Specific single species or species group management measures (7.4)	Legal Administrative	high importance (H)	Both	Maintain

2. Biogeographical Or Marine Level

2.1 Biogeographical Region

2.2 Published sources

Continental (CON)

The present species assessment (fields 0.1-2.9) has been compiled by Daniele Paoloni, Cristiano Spilinga (Associazione Teriologica Italiana - ATIt) and Anna Alonzi, Piero Genovesi, Francesca Ronchi (Institute for Environmental Protection and Research - ISPRA). Information, unpublished data and experts' judgments have been provided by Paolo Agnelli, Mara Calvini, Luca Cistrone, Michele Ferretto, Danilo Russo, Dino Scaravelli, Martina Spada, Roberto Toffoli, Simone Vergari (Italian Group for bat Research).

Archivio Osservatorio Regionale per Biodiversità. Regione Umbria.

Banca Dati Regionale Emilia Romagna (aggiornamento al 2010).

Calvini M., 2006. Monitoraggio dei chirotteri nella piana del Magra e Vallecchia

22/04/2014 11.59.31 Page 5 of 12

(SP) (rapporto interno).

Calvini M., 2007. Studio preliminare sulla chirotterofauna delle tre foreste demaniali del Parco dell'Aveto (rapporto interno).

Calvini M., 2009. I Chirotteri del SIC IT1110022 Stagno di Oulx e IT1110020 Lago di Viverone. IPLA (rapporto interno).

Calvini M., 2010. Monitoraggio delle colonie di chirotteri riproduttive e svernanti di particolare interesse conservazionistico note in Liguria (rapporto interno).

Dall'Asta A., 1995-1996. Atlante preliminare dei Chirotteri (Chiroptera, Mammalia) della Regione Friuli-Venezia Giulia - Prima Sintesi Cartografica. Tesi di Laurea in Scienze Naturali, Fac. di Scienze MM. FF. NN. dell'Università degli Studi di Trieste, Relatori G. A. Amirante & S. Dolce: 1-103.

Database del Repertorio Naturalistico Toscano.

Debernardi P., Patriarca E., Toffoli R., 2010. Monitoraggio delle colonie di chirotteri riproduttive e svernanti di particolare interesse conservazionistico note in Piemonte e dati preliminari sull'attività di swarming. Stato delle conoscenze al 30 aprile 2010. CRC, Regione Piemonte - Direzione ambiente - Settore pianificazione e gestione aree naturali protette (relazione interna). Pp. 83.

Insubria DataBat, 2012. Data base chirotteri dell'Università degli Studi dell'Insubria aggiornato al 2012.

Lapini L., Dall'Asta A., Dublo L., Spoto M., Venier E., 1996 (1995). Materiali per una teriofauna dell'Italia Nord - Orientale (Mammalia, Friuli-Venezia Giulia). Gortania 17: 149-248

Kryštufek B., Rešek Donev N., 2005. The Atlas of Slovenian Bats (Chiroptera). Scopolia, 55 (2005): 1-92.

Patriarca E., Debernardi P., Toffoli R., 2012. Piano d'azione per i chirotteri del Piemonte. Regione Piemonte. Bozza. Versione pubblicata on line (231 pp.) su http://www.centroregionalechirotteri.org/

Ruffo S., Stoch F., 2005. Checklist e distribuzione della fauna italiana. Memorie del Museo Civico di storia naturale di Verona, 2.serie, Sezione scienze della Vita 16.

Scaravelli D., Gellini S., Matteucci C. & L. Cicognani (ed), 2001. Atlante Mammiferi Provincia di Ravenna, STERNA & Amm. Prov. Ravenna, 119 pp.

Scaravelli D., 1997. Onferno. Riserva Naturale Orientata. Regione Emilia Romagna e Giunti Editori.

Spada M., Preatoni G., Tosi G., Martinoli A., 2010. Piano di monitoraggio dei Vertebrati terrestri di interesse comunitario (Direttive 79/409/CEE e 92/43/CEE) in Lombardia. Il monitoraggio dei Chirotteri. Fondazione Lombardia per l'Ambiente, Università degli Studi dell'Insubria.

22/04/2014 11.59.31 Page 6 of 12

Spilinga C., Russo D., Carletti S., Jiménez Grijalva M.P., Sergiacomi U., Ragni B., (in stampa). Chirotteri dell'Umbria. Distribuzione geografica ed ecologica. Regione Umbria. Università degli Studi di Perugia.

Vigorita V., Cucè L., 2008. La fauna selvatica in Lombardia. Rapporto 2008 su distribuzione, abbondanza e stato di conservazione di uccelli e mammiferi. Regione Lombardia. Pp. 364.

2.3 Range

2.3.1 Surface area - Range (km²) 2.3.2 Method - Range surface area 2.3.3 Short-term trend period 2.3.4 Short-term trend direction

2.3.5 Short-term trend magnitude

2.3.6 Long-term trend period

2.3.7 Long-term trend direction 2.3.8 Long-term trend magnitude

2.3.9 Favourable reference range

47700

Estimate based on partial data with some extrapolation and/or modelling (2)

2001-2012 stable (0)

min max

N/A

min max

area (km²)

operator approximately equal to (≈)

unkown

method Expert judgement

2.3.10 Reason for change

Improved knowledge/more accurate dataUse of different method

2.4 Population

2.4.1 Population size

(individuals or agreed exception)

Unit N/A

min max

2.4.2 Population size (other than individuals) Unit number of map 10x10 km grid cells (grids10x10)

88 88 min max

2.4.3 Additional information

Definition of locality

Conversion method

Problems Impossible to convert grids to individuals

2.4.4 Year or period

1985-2012

Estimate based on expert opinion with no or minimal sampling (1)

2.4.5 Method - population size 2.4.6 Short-term trend period

2001-2012

2.4.7 Short term trend direction

decrease (-)

2.4.8 Short-term trend magnitude 2.4.9 Short-term trend method

max

confidence interval Estimate based on expert opinion with no or minimal sampling (1)

2.4.10 Long-term trend period

N/A

min

2.4.11 Long term trend direction

confidence interval min max

2.4.12 Long-term trend magnitude 2.4.13 Long-term trend method

N/A

2.4.14 Favourable reference

population

number

operator more than (>)

unknown No

method Expert judgement

2.4.15 Reason for change

Improved knowledge/more accurate data Use of different method

2.5 Habitat for the Species

22/04/2014 11.59.31 Page 7 of 12

ii, iv alid v species (Alii	ick bj	
 2.5.1 Surface area - Habitat (km²) 2.5.2 Year or period 2.5.3 Method used - habitat 2.5.4 a) Quality of habitat 2.5.4 b) Quality of habitat - method 2.5.5 Short term trend period 2.5.6 Short term trend direction 2.5.7 Long-term trend period 2.5.8 Long term trend direction 2.5.9 Area of suitable habitat (km²) 2.5.10 Reason for change 	Absent data (0) Moderate Expert based 2001-2012 stable (0) N/A Improved knowledge/more accurate data Use of	different method
2.6 Main Pressures		
Pressure	ranking	pollution qualifier(s)
Mining and quarrying (C01)	high importance (H)	N/A
speleology (G01.04.02)	high importance (H)	N/A
recreational cave visits (G01.04.03)	high importance (H)	N/A
wind energy production (C03.03)	high importance (H)	N/A
closures of caves or galleries (G05.08)	high importance (H)	N/A
2.6.1 Method used – pressures	based only on expert judgements (1)	
2.7 Main Threats		
Threat	ranking	pollution qualifier(s)
Mining and quarrying (C01)	high importance (H)	N/A
speleology (G01.04.02)	high importance (H)	N/A
recreational cave visits (G01.04.03)	high importance (H)	N/A
wind energy production (C03.03)	high importance (H)	N/A
closures of caves or galleries (G05.08)	high importance (H)	N/A
2.7.1 Method used – threats	expert opinion (1)	
2.8 Complementary Information		
2.8.1 Justification of % thresholds for trends		
2.8.2 Other relevant Information		
2.8.3 Trans-boundary assessment		
2.9 Conclusions (assessment of con	servation status at end of reporting period)	
2.9.1 Range	assessment Favourable (FV) qualifiers N/A	
2.9.2. Population	assessment Inadequate (U1) qualifiers N/A	
2.9.3. Habitat	assessment Inadequate (U1) qualifiers N/A	
2.9.4. Future prospects	assessment Bad (U2)	

22/04/2014 11.59.31 Page 8 of 12

qualifiers N/A

Bad (U2)

2.9.5 Overall assessment of

Conservation Status

2.9.5 Overall trend in Conservation Status

3.1 Population

declining (-)

3. Natura 2000 coverage and conservation measures - Annex II species

3.1.1 Population Size Unit N/A min max

3.1.2 Method used Absent data (0)

3.1.3 Trend of population size within N/A

3.2 Conversation Measures

3.2.1 Measure	3.2.2 Type	3.2.3 Ranking	3.2.4 Location	3.2.5 Broad Evaluation
Other agriculture-related measures (2.0)	Administrative	medium importance (M)	Inside	No effect
Other forestry-related measures (3.0)	Administrative	medium importance (M)	Inside	No effect
Other spatial measures (6.0)	Administrative	medium importance (M)	Both	Maintain Long term
Legal protection of habitats and species (6.3)	s Legal	high importance (H)	Both	Not evaluated

2. Biogeographical Or Marine Level

2.1 Biogeographical Region

2.2 Published sources

Alpine (ALP)

The present species assessment (fields 0.1-2.9) has been compiled by Daniele Paoloni, Cristiano Spilinga (Associazione Teriologica Italiana - ATIt) and Anna Alonzi, Piero Genovesi, Francesca Ronchi (Institute for Environmental Protection and Research - ISPRA). Information, unpublished data and experts' judgments have been provided by Paolo Agnelli, Mara Calvini, Luca Cistrone, Michele Ferretto, Danilo Russo, Dino Scaravelli, Martina Spada, Roberto Toffoli, Simone Vergari (Italian Group for bat Research).

Distribution data for the following Nature 2000 sites have been inserted by the Ministry of Environment (source: Italian Nature 2000 database): IT1160057.

Calvini M., 2007. I Chirotteri delle Alpi Liguri; 24 pag. Provincia di Imperia, Regione Liguria.

Calvini M., 2009. I Chirotteri del SIC IT1110022 Stagno di Oulx e IT1110020 Lago di Viverone. IPLA (rapporto interno).

Calvini M., 2009. Indagine sulla chirotterofauna nel SIC "Bric Tana-Bric Mongarda", comune di Millesimo (SV).

Calvini M., 2009. Indagine chirotterologica nei seguenti SIC della provincia di Savona: IT1323201, IT1324011, IT1323112 e IT1323203 (rapporto interno).

Calvini M., 2010. Monitoraggio delle colonie di chirotteri riproduttive e svernanti

22/04/2014 11.59.31 Page 9 of 12

di particolare interesse conservazionistico note in Liguria (rapporto interno).

Dall'Asta A., 1995-1996. Atlante preliminare dei Chirotteri (Chiroptera, Mammalia) della Regione Friuli-Venezia Giulia - Prima Sintesi Cartografica. Tesi di Laurea in Scienze Naturali, Fac. Di Scienze MM. FF. NN. Dell'Università degli Studi di Trieste, Relatori G. A. Amirante & S. Dolce: 1-103.

Lapini L., Dall'Asta A., Dublo L., Spoto M., Venier E., 1996 (1995). Materiali per una teriofauna dell'Italia Nord - Orientale (Mammalia, Friuli-Venezia Giulia). Gortania 17: 149-248

Ruffo S., Stoch F., 2005. Checklist e distribuzione della fauna italiana. Memorie del Museo Civico di storia naturale di Verona, 2. serie, Sezione scienze della Vita 16.

2.3 Range

2.3.1 Surface area - Range (km²)

2.3.2 Method - Range surface area

2.3.3 Short-term trend period

2.3.4 Short-term trend direction

2.3.5 Short-term trend magnitude

2.3.6 Long-term trend period

2.3.7 Long-term trend direction

2.3.8 Long-term trend magnitude

2.3.9 Favourable reference range

10300

Estimate based on partial data with some extrapolation and/or modelling (2)

2001-2012

stable (0)

min max

N/A

min max

area (km²)

operator approximately equal to (≈)

unkown

method Expert judgement

2.3.10 Reason for change

Improved knowledge/more accurate dataUse of different method

2.4 Population

2.4.1 Population size

(individuals or agreed exception)

2.4.2 Population size

(other than individuals)

2.4.3 Additional information

Unit N/A

min max

Unit number of map 10x10 km grid cells (grids10x10)

min max

Definition of locality

Problems Impossible to convert grids into individuals

2.4.4 Year or period

2.4.5 Method – population size

2.4.6 Short-term trend period

2.4.7 Short term trend direction

2.4.8 Short-term trend magnitude

2.4.9 Short-term trend method

2.4.10 Long-term trend period

2.4.11 Long term trend direction

Conversion method

1985-2012

Estimate based on expert opinion with no or minimal sampling (1)

2001-2012

decrease (-)

min max confidence interval

Estimate based on expert opinion with no or minimal sampling (1)

N/A

22/04/2014 11.59.31 Page 10 of 12

2.4.12 Long-term trend magnitude	min	max	confidence in	terval
2.4.13 Long-term trend method 2.4.14 Favourable reference	N/A number			
population	operator unknown	more than (>) No		
	method	Expert judgement		
2.4.15 Reason for change		nowledge/more accurat	e data Use of different	method
2.5 Habitat for the Species	·	<i>.</i>		
2.5.1 Surface area - Habitat (km²) 2.5.2 Year or period 2.5.3 Method used - habitat 2.5.4 a) Quality of habitat	Absent dat Moderate	a (0)		
2.5.4 b) Quality of habitat - method 2.5.5 Short term trend period 2.5.6 Short term trend direction	Expert base 2001-2012 stable (0)			
 2.5.7 Long-term trend period 2.5.8 Long term trend direction 2.5.9 Area of suitable habitat (km²) 	N/A			
2.5.10 Reason for change	Improved l	knowledge/more accurat	e data Use of different	method
2.6 Main Pressures				
Pressure		ranking	poll	ution qualifier(s)
Mining and quarrying (C01)		high importa	nce (H) N/A	
speleology (G01.04.02)		high importa	nce (H) N/A	
recreational cave visits (G01.04.03)		high importa	nce (H) N/A	
wind energy production (C03.03)		low importar	nce (L) N/A	
closures of caves or galleries (G05.08)		high importa	nce (H) N/A	
2.6.1 Method used – pressures 2.7 Main Threats	based only	on expert judgements (2	1)	
Threat		ranking	poll	ution qualifier(s)
Mining and quarrying (C01)		high importa	nce (H) N/A	
speleology (G01.04.02)		high importa	nce (H) N/A	
recreational cave visits (G01.04.03)		high importa	nce (H) N/A	
wind energy production (C03.03)		low importar	nce (L) N/A	
closures of caves or galleries (G05.08)		high importa	nce (H) N/A	
2.7.1 Method used – threats	expert opir	nion (1)		
2.8 Complementary Information				
2.8.1 Justification of % thresholds for trends				
2.8.2 Other relevant Information				

2.9 Conclusions (assessment of conservation status at end of reporting period)

2.8.3 Trans-boundary assessment

22/04/2014 11.59.31 Page 11 of 12

2.9.1 Range

2.9.2. Population

2.9.3. Habitat

2.9.4. Future prospects

2.9.5 Overall assessment of Conservation Status

2.9.5 Overall trend in Conservation Status

assessment qualifiers

Bad (U2)

declining (-)

assessment Favourable (FV)
qualifiers N/A
assessment Inadequate (U1)
qualifiers N/A
assessment Inadequate (U1)
qualifiers N/A
assessment Bad (U2)
qualifiers N/A
Bad (U2)

3. Natura 2000 coverage and conservation measures - Annex II species

3.1 Population						
3.1.1 Population Size		Unit N	/A			
		min	max			
3.1.2 Method used		Absent data (0)				
3.1.3 Trend of population size within		N/A				
3.2 Conversation Measu	ires					
3.2.1 Measure	3.2.2 Type		3.2.3 Ranking	3.2.4 Location	3.2.5 Broad Evaluation	
Other agriculture-related measures (2.0)	Contractua		medium importance (M)	Inside	No effect	
Other forestry-related measures (3.0)	Contractua		medium importance (M)	Inside	No effect	

22/04/2014 11.59.31 Page 12 of 12