CODE: 91B0

NAME: Thermophilous Fraxinus angustifolia woods

#### 1. National Level

#### **1.1 Maps**

1.1.1 Distribution Map

1.1.2 Distribution Method

1.1.3 Year or period

1.1.4 Additional map

1.1.5 Range Map

Yes

Estimate based on expert opinion with no or minimal sampling (1)

2005-2012

No

Yes

#### 2. Biogeographical Or Marine Level

2.1 Biogeographical Region

2.2 Published

#### **Mediterranean (MED)**

The present Habitat assessment (fields 0.1-3.1) has been compiled by Pierangela Angelini (ISPRA). Published and unpublished data, information and experts' judgments have been provided by Edoardo Biondi, Liliana Zivkovic and Giovanni Spampinato(SBI), Pietro Massimiliano Bianco and Pierangela Angelini (ISPRA, field 2.7.1).

Angelini P., Augello R., Bianco P.M., Gennaio R., La Ghezza V., Lavarra P., Marrese M., Papallo O., Perrino V. M., Sani R., M. Stelluti. 2012. Carta degli habitat della Regione Puglia per il sistema informativo di Carta della Natura alla scala 1:50.000. ISPRA - Arpa Puglia

Biondi E, Blasi C, Burrascano S, Casavecchia S, Copiz R, Del Vico E, Galdenzi D, Gigante D, Lasen C, Spampinato G, Venanzoni R, Zivkovic L (2009a) Italian interpretation Manual of the habitats (92/43/EEC Directive). Ministero dell'Ambiente e della Tutela del Territorio e del Mare. http://vnr.unipg.it/habitat/Blasi et al., 2010. La Vegetazione d'Italia con Carta delle Serie di Vegetazione in scala 1:500000. Palombi ed.,

Casella L., Agrillo E., Bianco P.M., Cardillo A., Carbone M., Cattena C., Laureti L., Lugari A., Spada F., 2008. Carta degli habitat della Regione Lazio per il sistema informativo di Carta della Natura alla scala 1:50.000. ISPRA - Università degli Studi di Roma "La Sapienza" - Regione Lazio

ISPRA, 2011. Dati del sistema informativo di Carta della Natura alla scala 1:50.000.

ISPRA, Corine land cover 2006 IV livello. Dati della Rete del sistema Informativo Nazionale Ambientale - SINAnet

ISPRA, 2005. Dati del sistema informativo di Carta della Natura alla scala 1:50.000.

Papini F., Gianguzzi L., Brullo S., Bianco P. M., Angelini P., 2006. Carta degli habitat della Regione Sicilia per il sistema informativo di Carta della Natura alla scala 1:50.000. Dipartimento di Scienze Botaniche dell'Università degli Studi di Palermo - Dipartimento di Botanica dell'Università degli Studi di Catania -Regione Sicilia – ISPRA

Pedrotti F., 1980. Foreste ripariali lungo la costa adriatica dell'Italia. Coll. Phtosoc. IX: 143-154.

Pedrotti F., Gafta D., 1996. Ecologia delle foreste ripariali e paludose dell'Italia. L'uomo e l'ambiente, 23

11/04/2014 09:37:45 Page 1 of 7

2.3 Range of the habitat type in the biogeographical region or marine region

2.3.1 Surface area - Range (km²) 9300

2.3.2 Range method used Estimate based on expert opinion with no or minimal sampling (1)

2.3.3 Short-term trend period 2001-2012 2.3.4 Short-term trend direction decrease (-)

2.3.5 Short-term trend magnitude min max

2.3.6 Long-term trend period

2.3.7 Long-term trend direction N/A

2.3.8 Long-term trend magnitude min max

2.3.9 Favourable reference range area (km²)

operator much more than (>>)

unkown No

method

2.3.10 Reason for change Improved knowledge/more accurate data Use of different method

#### 2.4 Area covered by Habitat

2.4.1 Surface area (km²) 6,55

2.4.2 Year or period 2005-2012

2.4.3 Method used Estimate based on expert opinion with no or minimal sampling (1)

2.4.4 Short-term trend period 2001-2012 2.4.5 Short-term trend direction decrease (-)

2.4.6 Short-term trend magnitude min max

2.4.8 Long-term trend period

2.4.9 Long-term trend direction N/A

2.4.10 Long-term trend magnitude min max

2.4.11 Long term trend method used N/A

2.4.12 Favourable reference area area (km)

operator much more than (>>)

unknown No

method

2.4.13 Reason for change Improved knowledge/more accurate data Use of different method

#### 2.5 Main Pressures

Pressure	ranking	pollution qualifier(s)
roads, motorways (D01.02)	medium importance (M)	N/A
burning down (J01.01)	medium importance (M)	N/A
motorised vehicles (G01.03)	medium importance (M)	N/A
removal of forest undergrowth (B02.03)	low importance (L)	N/A
discontinuous urbanisation (E01.02)	low importance (L)	N/A
dispersed habitation (E01.03)	medium importance (M)	N/A
artificial planting on open ground (non-native trees) (B01.02)	medium importance (M)	N/A
Discharges (E03)	low importance (L)	N/A
electricity and phone lines (D02.01)	medium importance (M)	N/A
forest exploitation without replanting or natural regrowth (B03)	medium importance (M)	N/A

11/04/2014 09:37:45 Page 2 of 7

invasive non-native species (IO1)	medium importance (M)	N/A	
· · · · ·		•	
	ert judgement and other data (	۷)	
2.6 Main Threats Threat	ranking	pollution qualifier(s)	
roads, motorways (D01.02)	medium importance (M)	N/A	
burning down (J01.01)	medium importance (M)	N/A	
motorised vehicles (G01.03)	· · · · · · · · · · · · · · · · · · ·	<u> </u>	
,	medium importance (M)	N/A	
removal of forest undergrowth (B02.03)	low importance (L)	N/A	
discontinuous urbanisation (E01.02)	low importance (L)	N/A	
dispersed habitation (E01.03)	medium importance (M)	N/A	
artificial planting on open ground (non-native trees) (B01.02)	medium importance (M)	N/A	
Discharges (E03)	low importance (L)	N/A	
electricity and phone lines (D02.01)	medium importance (M)	N/A	
forest exploitation without replanting or natural regrowth (B03)	medium importance (M)	N/A	
invasive non-native species (I01)	medium importance (M)	N/A	
2.6.1 Method used – threats expert opinion (1)			
2.7 Complementary Information			
2.7.1 Species			
Carex pendula			
Carex remota			
Carex riparia			
Carex otrubae			
Fraxinus angustifolia ssp. Oxycarpa			
Fraxinus angustifolia ssp. Angustifolia			
Iris foetidisssima			
Iris pseudacorus			
Lycopus europaeus			
Lythrum salicaria			
Juncus conglomeratus			
Lysimachia vulgaris			
Oenanthe aquatica			
Ranunculus ficaria			
Rumex sanguineus			
Veronica scutellata			
Humulus lupulus			
Glyceria fluitans			

11/04/2014 09:37:45 Page 3 of 7

2.7.2 Species method used

Selected by ISPRA's expert from bibliographical and field research

2.7.3 Justification of % - thresholds for trends

2.7.4 Structure and functions - methods used

2.7.5 Other relevant information

Estimate based on expert opinion with no or minimal sampling (1)

#### 2.8 Conclusions (assessment of conservation status at end of reporting period)

2.8.1 Range

assessment Bad (U2) qualifiers N/A

2.8.2 Area

assessment Bad (U2)

qualifiers N/A assessment Bad (U2)

2.8.3 Specific structures and functions (incl Species)

qualifiers N/A

2.8.4 Future prospects

assessment Inadequate (U1)

2.6.4 Future prospects

qualifiers N/A

2.8.5 Overall assessment of

Conservation Status

Bad (U2)

2.8.5 Overall trend in Conservation Status

unknown (x)

### 3. Natura 2000 coverage conservation measures - Annex I habitat types on biogeographical level

#### 3.1 Area covered by habitat

3.1.1 Surface area (km²) min 2,8293 max 2,8293

3.1.2 Method used Complete survey/Complete survey or a statistically robust estimate (3)

3.1.3. Trend of surface area N/A

#### **3.2 Conversation Measures**

3.2.1 Measure	3.2.2 Type	3.2.3 Ranking	3.2.4 Location	3.2.5 Broad Evaluation
Restoring/improving forest habitats (3.1)	Recurrent	high importance (H)	Both	Enhance
Adapt forest management (3.2)	Administrative Recurrent	high importance (H)	Both	Maintain Enhance Long term
Establish protected areas/sites (6.1)	Legal	high importance (H)	Inside	Not evaluated

#### 2.1 Biogeographical Region

2.2 Published

#### Continental (CON)

The present Habitat assessment (fields 0.1-3.1) has been compiled by Pierangela Angelini (ISPRA). Published and unpublished data, information and experts' judgments have been provided by Edoardo Biondi and Liliana Zivkovic(SBI), Pietro Massimiliano Bianco and Pierangela Angelini (ISPRA, field 2.7.1). Allegrezza M., Mentoni M. & Tesei G., 2010. Geomorfologia e paesaggio vegetale: l'esempio della grande frana di Pescacci (Comune di Serra San Quirico-Appennino centrale). Fitosocioligia 47(2): 57-97. Biondi E, Blasi C, Burrascano S, Casavecchia S, Copiz R, Del Vico E, Galdenzi D,

Biondi E, Blasi C, Burrascano S, Casavecchia S, Copiz R, Del Vico E, Galdenzi D Gigante D, Lasen C, Spampinato G, Venanzoni R, Zivkovic L (2009a) Italian

11/04/2014 09:37:45 Page 4 of 7

interpretation Manual of the habitats (92/43/EEC Directive). Ministero dell'Ambiente e della Tutela del Territorio e del Mare. http://vnr.unipg.it/habitat Biondi E. & Allegrezza M., 2004. L'Ambiente della Selva di Gallignano. I Quaderni della Selva. Vol. II.

Blasi et al., 2010. La Vegetazione d'Italia con Carta delle Serie di Vegetazione in scala 1:500000. Palombi ed., ISPRA, Corine land cover 2006 IV livello. Dati della Rete del sistema Informativo Nazionale Ambientale - SINAnet

Pedrotti F., 1980. Foreste ripariali lungo la costa adriatica dell'Italia. Coll.

phtosoc. IX: 143-154.

Pedrotti F., Gafta D., 1996. Ecologia delle foreste ripariali e paludose dell'Italia.

L'uomo e l'ambiente, 23

#### 2.3 Range of the habitat type in the biogeographical region or marine region

2.3.1 Surface area - Range (km²) 2100

2.3.2 Range method used Estimate based on expert opinion with no or minimal sampling (1)

2.3.3 Short-term trend period 2001-2012 2.3.4 Short-term trend direction decrease (-)

2.3.5 Short-term trend magnitude min max

2.3.6 Long-term trend period

2.3.7 Long-term trend direction N/A

2.3.8 Long-term trend magnitude min max

2.3.9 Favourable reference range area (km²)

operator much more than (>>)

unkown No

method

2.3.10 Reason for change

Improved knowledge/more accurate data Use of different method

#### 2.4 Area covered by Habitat

2.4.1 Surface area (km²) 1,14

2.4.2 Year or period 2005-2012

2.4.3 Method used Estimate based on expert opinion with no or minimal sampling (1)

2.4.4 Short-term trend period 2001-2012 2.4.5 Short-term trend direction decrease (-)

2.4.6 Short-term trend magnitude min max

2.4.7 Short term trend method used Estimate based on expert opinion with no or minimal sampling (1)

2.4.8 Long-term trend period

2.4.9 Long-term trend direction N/A

2.4.10 Long-term trend magnitude min max

2.4.11 Long term trend method used

N/A

2.4.12 Favourable reference area area (km)

operator much more than (>>)

unknown No

method

2.4.13 Reason for change Improved knowledge/more accurate data Use of different method

#### 2.5 Main Pressures

11/04/2014 09:37:45 Page 5 of 7

nabitat types (Annex D)		
Pressure	ranking	pollution qualifier(s)
roads, motorways (D01.02)	medium importance (M)	N/A
burning down (J01.01)	medium importance (M)	N/A
motorised vehicles (G01.03)	medium importance (M)	N/A
removal of forest undergrowth (B02.03)	low importance (L)	N/A
discontinuous urbanisation (E01.02)	low importance (L)	N/A
dispersed habitation (E01.03)	medium importance (M)	N/A
artificial planting on open ground (non-native trees) (B01.02)	medium importance (M)	N/A
Discharges (E03)	low importance (L)	N/A
electricity and phone lines (D02.01)	medium importance (M)	N/A
forest exploitation without replanting or natural regrowth (B03)	medium importance (M)	N/A
invasive non-native species (I01)	medium importance (M)	N/A
2.5.1 Method used – pressures mainly based on exp	ert judgement and other data (2	2)
2.6 Main Threats		
Threat	ranking	pollution qualifier(s)
roads, motorways (D01.02)	medium importance (M)	N/A
burning down (J01.01)	medium importance (M)	N/A
motorised vehicles (G01.03)	medium importance (M)	N/A
removal of forest undergrowth (B02.03)	low importance (L)	N/A
discontinuous urbanisation (E01.02)	low importance (L)	N/A
dispersed habitation (E01.03)	medium importance (M)	N/A
artificial planting on open ground (non-native trees) (B01.02)	medium importance (M)	N/A
Discharges (E03)	low importance (L)	N/A
electricity and phone lines (D02.01)	medium importance (M)	N/A
forest exploitation without replanting or natural regrowth (B03)	medium importance (M)	N/A
invasive non-native species (I01)	medium importance (M)	N/A
2.6.1 Method used – threats expert opinion (1)		
2.7 Complementary Information		
2.7.1 Species		
Carex otrubae		
Carex pendula		
Carex remota		
Clematis viticella		
Fraxinus angustifolia ssp. Angustifolia		
Fraxinus angustifolia ssp. Oxycarpa		
Glyceria fluitans		
Humulus lupulus		
<u> </u>		

11/04/2014 09:37:45 Page 6 of 7

Iris foetidissima

habitat types (Annex I	D)
Iris pseudacorus	
Juncus conglomeratus	
Lycopus europaeus	
Lysimachia vulgaris	
Lythrum salicaria	
Oenanthe aquatica	
Ranunculus ficaria	
Rumex sanguineus	
Veronica scutellata	
2.7.2 Species method used	Selected by ISPRA's expert from bibliographical and field research
2.7.3 Justification of % - thresholds for trends	
2.7.4 Structure and functions - methods used	Estimate based on expert opinion with no or minimal sampling (1)
2.7.5 Other relevant information	
2.8 Conclusions (assessment of co	onservation status at end of reporting period)
2.8.1 Range	assessment Bad (U2) qualifiers N/A
2.8.2 Area	assessment Bad (U2) qualifiers N/A
2.8.3 Specific structures	assessment Bad (U2)
and functions (incl Species) 2.8.4 Future prospects	qualifiers N/A
2.8.4 Future prospects	assessment Inadequate (U1) qualifiers N/A
2.8.5 Overall assessment of Conservation Status	Bad (U2)
2.8.5 Overall trend in Conservation Status	unknown (x)
3. Natura 2000 coverage Annex I habitat types on 3.1 Area covered by habitat	conservation measures - biogeographical level
3.1.1 Surface area (km²)	min 1,14 max 1,14
3.1.2 Method used	Complete survey/Complete survey or a statistically robust estimate (3)
3.1.3. Trend of surface area	N/A
3.2 Conversation Measures	
3.2.1 Measure 3.2.2 Typ	pe 3.2.3 Ranking 3.2.4 Location 3.2.5 Broad Evaluation
No measure known/ impossible to carry out specific measures (1.3)	()

11/04/2014 09:37:45 Page 7 of 7