

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

CODE: 3170

NAME: Mediterranean temporary ponds

1. National Level

1.1 Maps

1.1.1 Distribution Map	Yes
1.1.2 Distribution Method	Estimate based on expert opinion with no or minimal sampling (1)
1.1.3 Year or period	2005-2012
1.1.4 Additional map	No
1.1.5 Range Map	Yes

2. Biogeographical Or Marine Level

2.1 Biogeographical Region

2.2 Published

Mediterranean (MED)

The present Habitat assessment (fields 0.1-3.1) has been compiled by Pierangela Angelini (ISPRA). Published and unpublished data, information and experts' judgments have been provided by Edoardo Biondi, Liliana Zivkovic and Giovanni Spampinato(SBI), Pietro Massimiliano Bianco and Pierangela Angelini (ISPRA, field 2.7.1).

Copiz R., Zavattero L., 2009. Rete ecologica del Parco Nazionale del Circeo: analisi dello status e della distribuzione di specie e habitat e definizione degli elementi della rete. Università di Roma La Sapienza, Dip.to di Biologia Vegetale. Inedito.

Blasi C., Manes F. (a cura di), 2001. Studi propedeutici alla stesura del piano del Parco Nazionale del Circeo: componenti flora, vegetazione e unità di paesaggio. Università di Roma La Sapienza, Dip.to di Biologia Vegetale. Inedito.

Alfonso G., Belmonte G., Ernandes P., Zuccarello V., 2011. Stagni temporanei mediterranei in Puglia. Biodiversità e aspetti di un habitat poco conosciuto. Ed. Grifo.

Biondi E. & Bagella S., 2005. Vegetazione e paesaggio vegetale dell'arcipelago di La Maddalena (Sardegna nord-orientale). Fitosociologia 42(2) suppl.1.

Ernandes P., 2011. Il genere *Isoetes* (Pteridophyta, Lycopsidea): note tassonomiche, ecologia e distribuzione in Puglia. Ann. Mus. Civ. Rovereto. Sez.: Arch., St., Sc. nat. vol. 26 (2010): 347-358.

Bagella S., Gascon S., Caria M.C., Sala J., Mariani M.A., Boix D., 2010. Identifying key environmental factors related to plant and crustacean assemblages in Mediterranean temporary ponds. Biodivers Conserv 19:1749–1768. DOI 10.1007/s10531-010-9801-5

Bagella S., Gascon S., Caria M.C., Sala J., Boix D., 2011. Cross-taxon congruence in Mediterranean temporary wetlands: vascular plants, crustaceans, and coleopterans. Community Ecology 12(1): 40-50.

Bagella S., Caria M.C., Zuccarello V., 2010. Patterns of emblematic habitat types in Mediterranean temporary wetlands. C. R. Biologies 333 (2010) 694–700.

Bagella S. & Caria M.C., 2012. Diversity and ecological characteristics of vascular flora in Mediterranean temporary pools. C. R. Biologies 335 (2012) 69–76

Bagella S., Caria M.C., Farris E. & Filigheddu R., 2007. Issues related to the classification of Mediterranean temporary wet habitats according with the European Union Habitats Directive. Fitosociologia vol. 44 (2) suppl. 1: 245-249

Bagella S., Caria M.C., Farris E., Filigheddu R., 2009. Phytosociological analysis in

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

- Sardinian Mediterranean temporary wet habitats. *Fitosociologia* vol. 46 (1): 11-26
- Bagella S., Caria M.C., 2011. Vegetation series: a tool for the assessment of grassland ecosystem services in Mediterranean large-scale grazing systems. *Fitosociologia* vol. 48 (2) suppl. 1: 47-54
- Bagella S., Caria M.C., Farris E., Filigheddu R., 2009. Spatial-time variability and conservation relevance of plant communities in Mediterranean temporary wet habitats: A case study in Sardinia (Italy). *Plant Biosystems*, Vol. 143, No. 3: 435–442
- Bagella S., Caria M.C., Molins A., Rosselló J.A., 2011. Different spore structures in sympatric *Isoetes histrix* populations and their relationship with gross morphology, chromosome number, and ribosomal nuclear ITS sequences. *Flora* 206: 451–457.
- Angelini P., Augello R., Bianco P.M., Gennaio R., La Ghezza V., Lavarra P., Marrese M., Papallo O., Perrino V. M., Sani R., M. Stelluti. 2012. Carta degli habitat della Regione Puglia per il sistema informativo di Carta della Natura alla scala 1:50.000. ISPRA - Arpa Puglia
- Bianco P.M., Laureti L., Papallo O. , Perfetti D. 2012 Carta degli habitat della Regione Umbria per il sistema informativo di Carta della Natura alla scala 1:50.000. ISPRA
- Biondi E, Blasi C, Burrascano S, Casavecchia S, Copiz R, Del Vico E, Galdenzi D, Gigante D, Lasen C, Spampinato G, Venanzoni R, Zivkovic L (2009a) Italian interpretation Manual of the habitats (92/43/EEC Directive). Ministero dell'Ambiente e della Tutela del Territorio e del Mare. <http://vnr.unipg.it/habitat/>
- Blasi et al., 2010. La Vegetazione d'Italia con Carta delle Serie di Vegetazione in scala 1:500000. Palombi ed.
- Camarda I., Carta L., Brunu A., Brundu G. , Laureti L. , Angelini P., Bagnaia R., 2011. Carta degli habitat della Regione Sardegna per il sistema informativo di Carta della Natura alla scala 1:50.000. Dipartimento di Scienze Botaniche Ecologiche e Geologiche dell'Università degli Studi di Sassari - ISPRA - Regione Sardegna
- Casella L., Agrillo E., Bianco P.M., Cardillo A., Carbone M., Cattena C., Laureti L., Lugari A., Spada F., 2008. Carta degli habitat della Regione Lazio per il sistema informativo di Carta della Natura alla scala 1:50.000. ISPRA - Università degli Studi di Roma "La Sapienza" - Regione Lazio
- ISPRA, 2011. Dati del sistema informativo di Carta della Natura alla scala 1:50.000.
- ISPRA, Corine land cover 2006 IV livello. Dati della Rete del sistema Informativo Nazionale Ambientale - SINAnet
- ISPRA, 2005. Dati del sistema informativo di Carta della Natura alla scala 1:50.000.
- Papini F., Gianguzzi L., Brullo S., Bianco P. M., Angelini P., 2006. Carta degli habitat della Regione Sicilia per il sistema informativo di Carta della Natura alla scala 1:50.000. Dipartimento di Scienze Botaniche dell'Università degli Studi di Palermo - Dipartimento di Botanica dell'Università degli Studi di Catania -Regione Sicilia – ISPRA

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

2.3 Range of the habitat type in the biogeographical region or marine region

2.3.1 Surface area - Range (km ²)	64300	
2.3.2 Range method used	Estimate based on expert opinion with no or minimal sampling (1)	
2.3.3 Short-term trend period	2001-2012	
2.3.4 Short-term trend direction	decrease (-)	
2.3.5 Short-term trend magnitude	min	max
2.3.6 Long-term trend period		
2.3.7 Long-term trend direction	N/A	
2.3.8 Long-term trend magnitude	min	max
2.3.9 Favourable reference range	area (km ²)	
	operator	more than (>)
	unknown	No
	method	
2.3.10 Reason for change	genuine change	No
	improved knowledge	Yes
	different method	Yes

2.4 Area covered by Habitat

2.4.1 Surface area (km ²)	26,83		
2.4.2 Year or period	2005-2012		
2.4.3 Method used	Estimate based on expert opinion with no or minimal sampling (1)		
2.4.4 Short-term trend period	2001-2012		
2.4.5 Short-term trend direction	decrease (-)		
2.4.6 Short-term trend magnitude	min	max	confidence interval
2.4.7 Short term trend method used	Estimate based on expert opinion with no or minimal sampling (1)		
2.4.8 Long-term trend period			
2.4.9 Long-term trend direction	N/A		
2.4.10 Long-term trend magnitude	min	max	confidence interval
2.4.11 Long term trend method used	N/A		
2.4.12 Favourable reference area	area (km)		
	operator	more than (>)	
	unknown	No	
	method		
2.4.13 Reason for change	Improved knowledge/more accurate dataUse of different method		

2.5 Main Pressures

Pressure	ranking	pollution qualifier(s)
use of biocides, hormones and chemicals (A07)	medium importance (M)	N/A
Pollution to surface waters (limnic & terrestrial, marine & brackish) (H01)	low importance (L)	N/A
Fertilisation (A08)	low importance (L)	N/A
Landfill, land reclamation and drying out, general (J02.01)	medium importance (M)	N/A
Soil pollution and solid waste (excluding discharges) (H05)	medium importance (M)	N/A
modifying structures of inland water courses (J02.05.02)	high importance (H)	N/A

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

Water abstractions from groundwater (J02.07)	low importance (L)	N/A
--	--------------------	-----

2.5.1 Method used – pressures	Estimate based on partial data with some extrapolation and/or modelling(2)
-------------------------------	---

2.6 Main Threats

Threat	ranking	pollution qualifier(s)
use of biocides, hormones and chemicals (A07)	medium importance (M)	N/A
Pollution to surface waters (limnic & terrestrial, marine & brackish) (H01)	medium importance (M)	N/A
Fertilisation (A08)	medium importance (M)	N/A
Landfill, land reclamation and drying out, general (J02.01)	low importance (L)	N/A
modifying structures of inland water courses (J02.05.02)	medium importance (M)	N/A
Soil pollution and solid waste (excluding discharges) (H05)	medium importance (M)	N/A
Water abstractions from groundwater (J02.07)	high importance (H)	N/A

2.6.1 Method used – threats	Estimate based on expert opinion with no or minimal sampling(1)
-----------------------------	--

2.7 Complementary Information

2.7.1 Species

Schenkia spicata (=Centaurium spicatum)

Anagallis arvensis subsp. Parviflora

Centaurium pulchellum

Cicendia filiformis

Crypsis spp.

Cyperus flavescens

Cyperus fuscus

Elatine spp.

Exaculum pusillum

Gnaphalium uliginosum

Isolepis cerbua

Isoëtes spp.

Juncus bufonius

Juncus capitatus

Juncus tenageja

Myosotis caespitosa

Radiola linoides

Riccia spp.

Serapias lingua

Serapias vomeracea

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

2.7.2 Species method used

Selected by ISPRA's expert from bibliographical and field research

2.7.3 Justification of % - thresholds for trends

2.7.4 Structure and functions - methods used

Estimate based on expert opinion with no or minimal sampling(1)

2.7.5 Other relevant information

2.8 Conclusions (assessment of conservation status at end of reporting period)

2.8.1 Range

assessment Inadequate(U1)
qualifiers N/A

2.8.2 Area

assessment Inadequate(U1)
qualifiers N/A

2.8.3 Specific structures and functions (incl Species)

assessment Inadequate(U1)
qualifiers N/A

2.8.4 Future prospects

assessment Inadequate(U1)
qualifiers N/A

2.8.5 Overall assessment of Conservation Status

Inadequate(U1)

2.8.5 Overall trend in Conservation Status

stable(=)

3. Natura 2000 coverage conservation measures - Annex I habitat types on biogeographical level

3.1 Area covered by habitat

3.1.1 Surface area (km²)

min 26,0936 max 26,0936

3.1.2 Method used

Complete survey/Complete survey or a statistically robust estimate (3)

3.1.3. Trend of surface area

N/A

3.2 Conversation Measures

2.1 Biogeographical Region

2.2 Published

Continental (CON)

The present Habitat assessment (fields 0.1-3.1) has been compiled by Pierangela Angelini (ISPRA). Published and unpublished data, information and experts' judgments have been provided by Edoardo Biondi and Liliana Zivkovic(SBI), Pietro Massimiliano Bianco and Pierangela Angelini (ISPRA, field 2.7.1). Bianco P.M., Laureti L., Papallo O. , Perfetti D. 2012 Carta degli habitat della Regione Umbria per il sistema informativo di Carta della Natura alla scala 1:50.000. ISPRA

Biondi E, Blasi C, Burrascano S, Casavecchia S, Copiz R, Del Vico E, Galdenzi D, Gigante D, Lasen C, Spampinato G, Venanzoni R, Zivkovic L (2009a) Italian

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

interpretation Manual of the habitats (92/43/EEC Directive). Ministero dell'Ambiente e della Tutela del Territorio e del Mare. <http://vnr.unipg.it/habitat/>
 Blasi et al., 2010. La Vegetazione d'Italia con Carta delle Serie di Vegetazione in scala 1:500000. Palombi ed.
 Casella L., Agrillo E., Bianco P.M., Cardillo A., Carbone M., Cattena C., Laureti L., Lugari A., Spada F., 2008. Carta degli habitat della Regione Lazio per il sistema informativo di Carta della Natura alla scala 1:50.000. ISPRA - Università degli Studi di Roma "La Sapienza" - Regione Lazio
 ISPRA, 2011. Dati del sistema informativo di Carta della Natura alla scala 1:50.000.
 ISPRA, Corine land cover 2006 IV livello. Dati della Rete del sistema Informativo Nazionale Ambientale - SINAnet
 PIANO DI GESTIONE del SIC-zps IT4070002 "BARDELLO". Rapporto tecnico non pubblicato.

2.3 Range of the habitat type in the biogeographical region or marine region

2.3.1 Surface area - Range (km ²)	7300	
2.3.2 Range method used	Estimate based on expert opinion with no or minimal sampling (1)	
2.3.3 Short-term trend period	2001-2012	
2.3.4 Short-term trend direction	decrease (-)	
2.3.5 Short-term trend magnitude	min	max
2.3.6 Long-term trend period	N/A	
2.3.7 Long-term trend direction	N/A	
2.3.8 Long-term trend magnitude	min	max
2.3.9 Favourable reference range	area (km ²)	
	operator	much more than (>>)
	unknown	No
	method	
2.3.10 Reason for change	genuine change	No
	improved knowledge	Yes
	different method	Yes

2.4 Area covered by Habitat

2.4.1 Surface area (km ²)	1,93	
2.4.2 Year or period	2005-2012	
2.4.3 Method used	Estimate based on expert opinion with no or minimal sampling (1)	
2.4.4 Short-term trend period	2001-2012	
2.4.5 Short-term trend direction	decrease (-)	
2.4.6 Short-term trend magnitude	min	max confidence interval
2.4.7 Short term trend method used	Estimate based on expert opinion with no or minimal sampling (1)	
2.4.8 Long-term trend period	N/A	
2.4.9 Long-term trend direction	N/A	
2.4.10 Long-term trend magnitude	min	max confidence interval
2.4.11 Long term trend method used	N/A	
2.4.12 Favourable reference area	area (km)	
	operator	much more than (>>)
	unknown	No

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

method

2.4.13 Reason for change

Improved knowledge/more accurate data Use of different method

2.5 Main Pressures

Pressure	ranking	pollution qualifier(s)
use of biocides, hormones and chemicals (A07)	high importance (H)	N/A
Modification of hydrographic functioning, general (J02.05)	medium importance (M)	N/A
Pollution to surface waters (limnic & terrestrial, marine & brackish) (H01)	medium importance (M)	N/A
infilling of ditches, dykes, ponds, pools, marshes or pits (J02.01.03)	low importance (L)	N/A

2.5.1 Method used – pressures

Estimate based on partial data with some extrapolation and/or modelling(2)

2.6 Main Threats

Threat	ranking	pollution qualifier(s)
use of biocides, hormones and chemicals (A07)	high importance (H)	N/A
Modification of hydrographic functioning, general (J02.05)	medium importance (M)	N/A
Pollution to surface waters (limnic & terrestrial, marine & brackish) (H01)	medium importance (M)	N/A
infilling of ditches, dykes, ponds, pools, marshes or pits (J02.01.03)	low importance (L)	N/A

2.6.1 Method used – threats

Estimate based on expert opinion with no or minimal sampling(1)

2.7 Complementary Information

2.7.1 Species

Cicendia filiformis
Crypsis spp
Cyperus flavescent
Cyperus fuscus
Gnaphalium uliginosum
Illecebrum verticillatum
Juncus bufonius
Juncus capitatus
Juncus pygmaeus
Juncus tenageja
Lythrum spp.
Serapias spp.
Centaurium spp.
Isolepis cernua
Isolepis setacea
Mentha pulegium

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

Peplis portula

Radiola linoides

Riccia spp.

Schenkia spicata (=Centaurium spicatum)

2.7.2 Species method used Selected by ISPRA's expert from bibliographical and field research

2.7.3 Justification of % - thresholds for trends

2.7.4 Structure and functions - methods used Estimate based on expert opinion with no or minimal sampling(1)

2.7.5 Other relevant information

2.8 Conclusions (assessment of conservation status at end of reporting period)

2.8.1 Range assessment Bad(U2)
qualifiers N/A

2.8.2 Area assessment Bad(U2)
qualifiers N/A

2.8.3 Specific structures and functions (incl Species) assessment Inadequate(U1)
qualifiers N/A

2.8.4 Future prospects assessment Inadequate(U1)
qualifiers N/A

2.8.5 Overall assessment of Conservation Status Bad(U2)

2.8.5 Overall trend in Conservation Status declining(-)

3. Natura 2000 coverage conservation measures - Annex I habitat types on biogeographical level

3.1 Area covered by habitat

3.1.1 Surface area (km²) min 1,9071 max 1,9071

3.1.2 Method used Complete survey/Complete survey or a statistically robust estimate (3)

3.1.3. Trend of surface area N/A

3.2 Conversation Measures

2.1 Biogeographical Region

2.2 Published

Alpine (ALP)

The present Habitat assessment (fields 0.1-3.1) has been compiled by Pierangela Angelini (ISPRA). Published and unpublished data, information and experts' judgments have been provided by Edoardo Biondi, Liliana Zivkovic and Cesare Lasen(SBI), Pietro Massimiliano Bianco and Pierangela Angelini (ISPRA, field

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

2.7.1).

Biondi E, Blasi C, Burrascano S, Casavecchia S, Copiz R, Del Vico E, Galdenzi D, Gigante D, Lasen C, Spampinato G, Venanzoni R, Zivkovic L (2009a) Italian interpretation Manual of the habitats (92/43/EEC Directive). Ministero dell'Ambiente e della Tutela del Territorio e del Mare. <http://vnr.unipg.it/habitat/>
Blasi et al., 2010. La Vegetazione d'Italia con Carta delle Serie di Vegetazione in scala 1:500000. Palombi ed.
Casella L., Agrillo E., Bianco P.M., Cardillo A., Carbone M., Cattena C., Laureti L., Lugari A., Spada F., 2008. Carta degli habitat della Regione Lazio per il sistema informativo di Carta della Natura alla scala 1:50.000. ISPRA - Università degli Studi di Roma "La Sapienza" - Regione Lazio
ISPRA, 2011. Dati del sistema informativo di Carta della Natura alla scala 1:50.000.
ISPRA, Corine land cover 2006 IV livello. Dati della Rete del sistema Informativo Nazionale Ambientale - SINAnet

2.3 Range of the habitat type in the biogeographical region or marine region

2.3.1 Surface area - Range (km ²)	2800
2.3.2 Range method used	Estimate based on expert opinion with no or minimal sampling (1)
2.3.3 Short-term trend period	2001-2012
2.3.4 Short-term trend direction	unknown (x)
2.3.5 Short-term trend magnitude	min max
2.3.6 Long-term trend period	
2.3.7 Long-term trend direction	N/A
2.3.8 Long-term trend magnitude	min max
2.3.9 Favourable reference range	area (km ²) operator N/A unkown Yes method
2.3.10 Reason for change	genuine change No improved knowledge Yes different method Yes

2.4 Area covered by Habitat

2.4.1 Surface area (km ²)	9,5
2.4.2 Year or period	2005-2012
2.4.3 Method used	Estimate based on expert opinion with no or minimal sampling (1)
2.4.4 Short-term trend period	2001-2012
2.4.5 Short-term trend direction	unknown (x)
2.4.6 Short-term trend magnitude	min max confidence interval
2.4.7 Short term trend method used	Estimate based on expert opinion with no or minimal sampling (1)
2.4.8 Long-term trend period	
2.4.9 Long-term trend direction	N/A
2.4.10 Long-term trend magnitude	min max confidence interval
2.4.11 Long term trend method used	N/A

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

2.4.12 Favourable reference area

area (km)
operator N/A
unknown Yes
method

2.4.13 Reason for change

Improved knowledge/more accurate dataUse of different method

2.5 Main Pressures

Pressure	ranking	pollution qualifier(s)
canalisation (J02.03.02)	low importance (L)	N/A

2.5.1 Method used – pressures

Estimate based on expert opinion with no or minimal sampling(1)

2.6 Main Threats

Threat	ranking	pollution qualifier(s)
canalisation (J02.03.02)	low importance (L)	N/A

2.6.1 Method used – threats

Estimate based on expert opinion with no or minimal sampling(1)

2.7 Complementary Information

2.7.1 Species

Schenkia spicata (= Centaurium spicatum)

Isoëtes spp.

Juncus tenageja

Lythrum tribracteatum

Serapias spp.

Centaurium pulchellum

Isolepis cernua

Isolepis setacea

Lythrum hyssopifolia

Mentha pulegium

Myosotis caespitosa

Peplis portula

Radiola linoides

Ranunculus muricatus

Riccia spp.

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

2.7.2 Species method used	Selected by ISPRA's expert from bibliographical and field research
2.7.3 Justification of % - thresholds for trends	
2.7.4 Structure and functions - methods used	Estimate based on expert opinion with no or minimal sampling(1)
2.7.5 Other relevant information	

2.8 Conclusions (assessment of conservation status at end of reporting period)

2.8.1 Range	assessment Unknown(XX) qualifiers N/A
2.8.2 Area	assessment Unknown(XX) qualifiers N/A
2.8.3 Specific structures and functions (incl Species)	assessment Unknown(XX) qualifiers N/A
2.8.4 Future prospects	assessment Unknown(XX) qualifiers N/A
2.8.5 Overall assessment of Conservation Status	Unknown(XX)
2.8.5 Overall trend in Conservation Status	N/A

3. Natura 2000 coverage conservation measures - Annex I habitat types on biogeographical level

3.1 Area covered by habitat

3.1.1 Surface area (km²)	min 9,4999 max 9,4999
3.1.2 Method used	Complete survey/Complete survey or a statistically robust estimate (3)
3.1.3. Trend of surface area	N/A

3.2 Conversation Measures

Notes

Habitat code: 3170 Region code: ALP

Field label	Note	User
2.1 Region	Secondo gli esperti ISPRA la presenza dell'habitat in questa regione biogeografica è dubbia	ISPRA_habitat

Habitat code: 3170 Region code: MED

Field label	Note	User
2.4.5 Short time trend direction	Le azioni intraprese con diversi progetti di salvaguardia rendono le prospettive future di questo habitat medie stabili.	ISPRA_habitat
2.4.9 Long term trend direction	Le azioni intraprese con diversi progetti di salvaguardia rendono le prospettive future di questo habitat medie stabili.	ISPRA_habitat
2.4.1 Surface area	In Sicilia la distribuzione dell'habitat è sovradimensionata in relazione al fatto che per i SIC che non riportano dati regionali ma citano nella scheda Natura 2000 la presenza dell'habitat sono stati considerati tutti i quadranti che si sovrappongono ai confini del SIC.	ISPRA_habitat