

# The Fermi Hole

Ivan Kulesh    Martijn Papendrecht

Delft University of Technology, The Netherlands

March 5, 2018

## Some properties of Fermi gas

Particles dwell in a large but finite volume  $\mathcal{V}$ .

Periodical boundary conditions.

Restriction on values of the wave vector:

$$\mathbf{k} = 2\pi \left( \frac{n_x}{L_x}, \frac{n_y}{L_y}, \frac{n_z}{L_z} \right) \quad (1)$$


The volume of the unit cell in k-space:

$$dk_x dk_y dk_z = d^3k = \frac{(2\pi)^3}{\mathcal{V}} \quad (2)$$

## Some properties of Fermi gas

Particles occupy all energy levels below Fermi energy. Volume in k-space:

$$V_k = \frac{4}{3}\pi k_F^3 \quad (3)$$



## Some properties of Fermi gas

Particles occupy all energy levels below Fermi energy. Volume in k-space:


$$V_k = \frac{4}{3}\pi k_F^3 \quad (3)$$

Number of particles:

$$N = g_\sigma \frac{V_k}{d^3k} = \frac{2 \frac{4}{3}\pi k_F^3 \mathcal{V}}{8\pi^3} = \frac{k_F^3 \mathcal{V}}{3\pi^2} \implies \quad (4)$$
$$k_F^3 = \frac{3\pi^2 N}{\mathcal{V}}$$

## Fermionic ground state

In the ground state  $|g\rangle$  fermions occupy all available levels in the  $k$ -space up to  $k_F$ :



## Removing particle

We remove a particle at **position**  $\mathbf{x}$  with a spin  $\sigma$

$$|\phi_\sigma(\mathbf{x})\rangle = \hat{\psi}_\sigma(\mathbf{x}) |g\rangle$$

Particle density at this state:

$$\rho(\mathbf{x}', \sigma') = \langle \phi_\sigma(\mathbf{x}) | \hat{\psi}_{\sigma'}^\dagger(\mathbf{x}') \hat{\psi}_{\sigma'}(\mathbf{x}') | \phi_\sigma(\mathbf{x}) \rangle$$

We will show that the density equals to

$$\left( \frac{N}{2\mathcal{V}} \right)^2 g_{\sigma\sigma'}(\mathbf{x} - \mathbf{x}')$$

## Removing particle

We remove a particle at **position**  $\mathbf{x}$  with a spin  $\sigma$

$$|\phi_\sigma(\mathbf{x})\rangle = \hat{\psi}_\sigma(\mathbf{x}) |g\rangle$$

Particle density at this state:

$$\rho(\mathbf{x}', \sigma') = \langle \phi_\sigma(\mathbf{x}) | \hat{\psi}_{\sigma'}^\dagger(\mathbf{x}') \hat{\psi}_{\sigma'}(\mathbf{x}') | \phi_\sigma(\mathbf{x}) \rangle$$

We will show that the density equals to

$$\left( \frac{N}{2\mathcal{V}} \right)^2 g_{\sigma\sigma'}(\mathbf{x} - \mathbf{x}')$$

## Removing particle

We remove a particle at **position**  $\mathbf{x}$  with a spin  $\sigma$

$$|\phi_\sigma(\mathbf{x})\rangle = \hat{\psi}_\sigma(\mathbf{x}) |g\rangle$$

Particle density at this state:

$$\rho(\mathbf{x}', \sigma') = \langle \phi_\sigma(\mathbf{x}) | \hat{\psi}_{\sigma'}^\dagger(\mathbf{x}') \hat{\psi}_{\sigma'}(\mathbf{x}') | \phi_\sigma(\mathbf{x}) \rangle$$

We will show that the density equals to

$$\left(\frac{N}{2\mathcal{V}}\right)^2 g_{\sigma\sigma'}(\mathbf{x} - \mathbf{x}')$$



## Removing particle

The state after removing a particle. In coordinate space:

$$|\phi_{\sigma}(\mathbf{x})\rangle = \hat{\psi}_{\sigma}(\mathbf{x}) |g\rangle$$

But we want to use momentum space:

Fourier transform

$$\hat{\psi}_{\mathbf{k},\sigma}(\mathbf{x}) = \sum_{\mathbf{k}} \hat{c}_{\mathbf{k},\sigma} \psi_{\mathbf{k},\sigma}(\mathbf{x})$$

$$\psi_{\mathbf{k},\sigma}(\mathbf{x}) = \frac{1}{\sqrt{\mathcal{V}}} e^{i\mathbf{k}\cdot\mathbf{x}} \sigma$$

## Density in momentum space

$$\rho(\mathbf{x}', \sigma') =$$

$$\begin{aligned} \langle g | \sum_{\mathbf{k}} \hat{c}_{\mathbf{k},\sigma}^\dagger \psi_{\mathbf{k},\sigma}^*(\mathbf{x}) \sum_{\mathbf{l}} \hat{c}_{\mathbf{l},\sigma'}^\dagger \psi_{\mathbf{l},\sigma'}^*(\mathbf{x}') \sum_{\mathbf{m}} \hat{c}_{\mathbf{m},\sigma'} \psi_{\mathbf{m},\sigma'}(\mathbf{x}') \sum_{\mathbf{n}} \hat{c}_{\mathbf{n},\sigma} \psi_{\mathbf{n},\sigma}(\mathbf{x}) | g \rangle = \\ \sum_{\mathbf{k},\mathbf{l},\mathbf{m},\mathbf{n}} \langle g | \hat{c}_{\mathbf{k},\sigma}^\dagger \hat{c}_{\mathbf{l},\sigma'}^\dagger \hat{c}_{\mathbf{m},\sigma'} \hat{c}_{\mathbf{n},\sigma} | g \rangle \psi_{\mathbf{k},\sigma}^*(\mathbf{x}) \psi_{\mathbf{l},\sigma'}^*(\mathbf{x}') \psi_{\mathbf{m},\sigma'}(\mathbf{x}') \psi_{\mathbf{n},\sigma}(\mathbf{x}) \end{aligned}$$

## Density in momentum space

$$\rho(\mathbf{x}', \sigma') =$$

$$\begin{aligned} \langle g | \sum_{\mathbf{k}} \hat{c}_{\mathbf{k},\sigma}^\dagger \psi_{\mathbf{k},\sigma}^*(\mathbf{x}) \sum_{\mathbf{l}} \hat{c}_{\mathbf{l},\sigma'}^\dagger \psi_{\mathbf{l},\sigma'}^*(\mathbf{x}') \sum_{\mathbf{m}} \hat{c}_{\mathbf{m},\sigma'} \psi_{\mathbf{m},\sigma'}(\mathbf{x}') \sum_{\mathbf{n}} \hat{c}_{\mathbf{n},\sigma} \psi_{\mathbf{n},\sigma}(\mathbf{x}) | g \rangle = \\ \sum_{\mathbf{k},\mathbf{l},\mathbf{m},\mathbf{n}} \langle g | \hat{c}_{\mathbf{k},\sigma}^\dagger \hat{c}_{\mathbf{l},\sigma'}^\dagger \hat{c}_{\mathbf{m},\sigma'} \hat{c}_{\mathbf{n},\sigma} | g \rangle \psi_{\mathbf{k},\sigma}^*(\mathbf{x}) \psi_{\mathbf{l},\sigma'}^*(\mathbf{x}') \psi_{\mathbf{m},\sigma'}(\mathbf{x}') \psi_{\mathbf{n},\sigma}(\mathbf{x}) \end{aligned}$$


## Allowed operators

We will focus on the expression

$$\langle g | \hat{c}_{\mathbf{k},\sigma}^\dagger \hat{c}_{\mathbf{l},\sigma'}^\dagger \hat{c}_{\mathbf{m},\sigma'} \hat{c}_{\mathbf{n},\sigma} | g \rangle$$

Removing two particles:

$$|g\rangle$$




# Allowed operators

We will focus on the expression

$$\langle g | \hat{c}_{\mathbf{k},\sigma}^\dagger \hat{c}_{\mathbf{l},\sigma'}^\dagger \hat{c}_{\mathbf{m},\sigma'} \hat{c}_{\mathbf{n},\sigma} | g \rangle$$

Removing two particles:

$$\hat{c}_{\mathbf{n},\sigma} | g \rangle$$




## Allowed operators

We will focus on the expression

$$\langle g | \hat{c}_{\mathbf{k},\sigma}^\dagger \hat{c}_{\mathbf{l},\sigma'}^\dagger \hat{c}_{\mathbf{m},\sigma'} \hat{c}_{\mathbf{n},\sigma} | g \rangle$$

Removing two particles:

$$\hat{c}_{\mathbf{m},\sigma'} \hat{c}_{\mathbf{n},\sigma} | g \rangle$$




# Allowed operators

We will focus on the expression

$$\langle g | \hat{c}_{\mathbf{k},\sigma}^\dagger \hat{c}_{\mathbf{l},\sigma'}^\dagger \hat{c}_{\mathbf{m},\sigma'} \hat{c}_{\mathbf{n},\sigma} | g \rangle$$

Restoring the state:

$$\hat{c}_{\mathbf{k},\sigma}^\dagger \hat{c}_{\mathbf{l},\sigma'}^\dagger \hat{c}_{\mathbf{m},\sigma'} \hat{c}_{\mathbf{n},\sigma} | g \rangle$$



# Allowed operators

We will focus on the expression

$$\langle g | \hat{c}_{\mathbf{k},\sigma}^\dagger \hat{c}_{\mathbf{l},\sigma'}^\dagger \hat{c}_{\mathbf{m},\sigma'} \hat{c}_{\mathbf{n},\sigma} | g \rangle$$

Allowed indices

$$\begin{cases} \mathbf{k}, \sigma = \mathbf{m}, \sigma' \\ \mathbf{l}, \sigma' = \mathbf{n}, \sigma \\ \mathbf{k}, \sigma \neq \mathbf{l}, \sigma' \end{cases} \quad \begin{cases} \mathbf{k}, \sigma = \mathbf{n}, \sigma \\ \mathbf{l}, \sigma' = \mathbf{m}, \sigma' \\ \mathbf{k}, \sigma \neq \mathbf{l}, \sigma' \end{cases}$$



## Density for different spin

If spins are different

$$\sigma \neq \sigma' \implies \text{only } \begin{cases} \mathbf{k} = \mathbf{n} \\ \mathbf{l} = \mathbf{m} \end{cases}$$

We can simplify expression for density:

$$\rho(\mathbf{x}', \sigma') = \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}, \sigma}^\dagger \hat{c}_{\mathbf{l}, \sigma'}^\dagger \hat{c}_{\mathbf{l}, \sigma'} \hat{c}_{\mathbf{k}, \sigma} | g \rangle \psi_{\mathbf{k}, \sigma}^*(\mathbf{x}) \psi_{\mathbf{l}, \sigma'}^*(\mathbf{x}') \psi_{\mathbf{l}, \sigma'}(\mathbf{x}') \psi_{\mathbf{k}, \sigma}(\mathbf{x})$$

## Density for different spin

If spins are different

$$\sigma \neq \sigma' \implies \text{only } \begin{cases} \mathbf{k} = \mathbf{n} \\ \mathbf{l} = \mathbf{m} \end{cases}$$

We can simplify expression for density:

$$\rho(\mathbf{x}', \sigma') = \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}, \sigma}^\dagger \hat{c}_{\mathbf{l}, \sigma'}^\dagger \hat{c}_{\mathbf{l}, \sigma'} \hat{c}_{\mathbf{k}, \sigma} | g \rangle \psi_{\mathbf{k}, \sigma}^*(\mathbf{x}) \psi_{\mathbf{l}, \sigma'}^*(\mathbf{x}') \psi_{\mathbf{l}, \sigma'}(\mathbf{x}') \psi_{\mathbf{k}, \sigma}(\mathbf{x})$$

### Wave functions

$$\psi_{\mathbf{l}, \sigma'}(\mathbf{x}') = \frac{1}{\sqrt{\mathcal{V}}} e^{i\mathbf{l} \cdot \mathbf{x}'} \sigma'$$

$$\psi_{\mathbf{l}, \sigma'}^*(\mathbf{x}') = \frac{1}{\sqrt{\mathcal{V}}} e^{-i\mathbf{l} \cdot \mathbf{x}'} \sigma'$$

## Density for different spin

If spins are different

$$\sigma \neq \sigma' \implies \text{only } \begin{cases} \mathbf{k} = \mathbf{n} \\ \mathbf{l} = \mathbf{m} \end{cases}$$

We can simplify expression for density:

$$\rho(\mathbf{x}', \sigma') = \frac{1}{\mathcal{V}} \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}, \sigma}^\dagger \hat{c}_{\mathbf{l}, \sigma'}^\dagger \hat{c}_{\mathbf{l}, \sigma'} \hat{c}_{\mathbf{k}, \sigma} | g \rangle \psi_{\mathbf{k}, \sigma}^*(\mathbf{x}) \psi_{\mathbf{k}, \sigma}(\mathbf{x})$$

## Density for different spin

If spins are different

$$\sigma \neq \sigma' \implies \text{only } \begin{cases} \mathbf{k} = \mathbf{n} \\ \mathbf{l} = \mathbf{m} \end{cases}$$

We can simplify expression for density:

$$\rho(\mathbf{x}', \sigma') = \frac{1}{V^2} \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}, \sigma}^\dagger \hat{c}_{\mathbf{l}, \sigma'}^\dagger \hat{c}_{\mathbf{l}, \sigma'} \hat{c}_{\mathbf{k}, \sigma} | g \rangle$$

## Density for different spin

If spins are different

$$\sigma \neq \sigma' \implies \text{only } \begin{cases} \mathbf{k} = \mathbf{n} \\ \mathbf{l} = \mathbf{m} \end{cases}$$

We can simplify expression for density:

$$\rho(\mathbf{x}', \sigma') = \frac{1}{\mathcal{V}^2} \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}, \sigma}^\dagger \hat{c}_{\mathbf{l}, \sigma'}^\dagger \hat{c}_{\mathbf{l}, \sigma'} \hat{c}_{\mathbf{k}, \sigma} | g \rangle$$

### Commutation for fermions

$$\{\hat{c}_{\mathbf{l}, \sigma'}, \hat{c}_{\mathbf{k}, \sigma}\} = 0 \implies \hat{c}_{\mathbf{l}, \sigma'} \hat{c}_{\mathbf{k}, \sigma} = -\hat{c}_{\mathbf{k}, \sigma} \hat{c}_{\mathbf{l}, \sigma'}$$

## Density for different spin

If spins are different

$$\sigma \neq \sigma' \implies \text{only } \begin{cases} \mathbf{k} = \mathbf{n} \\ \mathbf{l} = \mathbf{m} \end{cases}$$

We can simplify expression for density:

$$\rho(\mathbf{x}', \sigma') = \frac{-1}{V^2} \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}, \sigma}^\dagger \hat{c}_{\mathbf{l}, \sigma'}^\dagger \hat{c}_{\mathbf{k}, \sigma} \hat{c}_{\mathbf{l}, \sigma'} | g \rangle$$

### Commutation for fermions

$$\{\hat{c}_{\mathbf{l}, \sigma'} \hat{c}_{\mathbf{k}, \sigma}\} = 0 \implies \hat{c}_{\mathbf{l}, \sigma'} \hat{c}_{\mathbf{k}, \sigma} = -\hat{c}_{\mathbf{k}, \sigma} \hat{c}_{\mathbf{l}, \sigma'}$$

## Density for different spin

If spins are different

$$\sigma \neq \sigma' \implies \text{only } \begin{cases} \mathbf{k} = \mathbf{n} \\ \mathbf{l} = \mathbf{m} \end{cases}$$

We can simplify expression for density:

$$\rho(\mathbf{x}', \sigma') = \frac{-1}{\mathcal{V}^2} \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}, \sigma}^\dagger \hat{c}_{\mathbf{l}, \sigma'}^\dagger \hat{c}_{\mathbf{k}, \sigma} \hat{c}_{\mathbf{l}, \sigma'} | g \rangle$$

### Commutation for fermions

$$\{\hat{c}_{\mathbf{l}, \sigma'}^\dagger, \hat{c}_{\mathbf{k}, \sigma}\} = \delta_{\mathbf{k}, \mathbf{l}} \implies \hat{c}_{\mathbf{l}, \sigma'}^\dagger \hat{c}_{\mathbf{k}, \sigma} = -\hat{c}_{\mathbf{k}, \sigma} \hat{c}_{\mathbf{l}, \sigma'}^\dagger + \delta_{\mathbf{k}, \mathbf{l}}$$

## Density for different spin

If spins are different

$$\sigma \neq \sigma' \implies \text{only } \begin{cases} \mathbf{k} = \mathbf{n} \\ \mathbf{l} = \mathbf{m} \end{cases}$$

We can simplify expression for density:

$$\rho(\mathbf{x}', \sigma') = \frac{1}{\mathcal{V}^2} \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}, \sigma}^\dagger \hat{c}_{\mathbf{k}, \sigma} \hat{c}_{\mathbf{l}, \sigma'}^\dagger \hat{c}_{\mathbf{l}, \sigma'} | g \rangle$$

### Commutation for fermions

$$\{\hat{c}_{\mathbf{l}, \sigma'}^\dagger, \hat{c}_{\mathbf{k}, \sigma}\} = \delta_{\mathbf{k}, \mathbf{l}} \implies \hat{c}_{\mathbf{l}, \sigma'}^\dagger \hat{c}_{\mathbf{k}, \sigma} = -\hat{c}_{\mathbf{k}, \sigma} \hat{c}_{\mathbf{l}, \sigma'}^\dagger + \delta_{\mathbf{k}, \mathbf{l}}$$



## Density for different spin

If spins are different

$$\sigma \neq \sigma' \implies \text{only } \begin{cases} \mathbf{k} = \mathbf{n} \\ \mathbf{l} = \mathbf{m} \end{cases}$$

We can simplify expression for density:

$$\begin{aligned} \rho(\mathbf{x}', \sigma') &= \frac{1}{\mathcal{V}^2} \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}, \sigma}^\dagger \hat{c}_{\mathbf{k}, \sigma} \hat{c}_{\mathbf{l}, \sigma'}^\dagger \hat{c}_{\mathbf{l}, \sigma'} | g \rangle - \\ &\quad - \delta_{\sigma, \sigma'} \frac{1}{\mathcal{V}^2} \sum_{\mathbf{k}} \langle g | \hat{c}_{\mathbf{k}, \sigma}^\dagger \hat{c}_{\mathbf{k}, \sigma} \hat{c}_{\mathbf{k}, \sigma'}^\dagger \hat{c}_{\mathbf{k}, \sigma'} | g \rangle \end{aligned}$$

## Density for different spin

Insert identity operator:

$$\rho(\mathbf{x}', \sigma') = \frac{1}{\mathcal{V}^2} \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}, \sigma}^\dagger \hat{c}_{\mathbf{k}, \sigma} \hat{\mathcal{I}} \hat{c}_{\mathbf{l}, \sigma'}^\dagger \hat{c}_{\mathbf{l}, \sigma'} | g \rangle$$

## Density for different spin

Insert identity operator:

$$\rho(\mathbf{x}', \sigma') = \frac{1}{\mathcal{V}^2} \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}, \sigma}^\dagger \hat{c}_{\mathbf{k}, \sigma} \hat{\mathcal{I}} \hat{c}_{\mathbf{l}, \sigma'}^\dagger \hat{c}_{\mathbf{l}, \sigma'} | g \rangle$$

Identity

$$\hat{\mathcal{I}} = \sum_n |n\rangle \langle n|$$

# Density for different spin

Insert identity operator:

$$\rho(\mathbf{x}', \sigma') = \frac{1}{\mathcal{V}^2} \sum_{\mathbf{k}, l} \langle g | \hat{c}_{\mathbf{k}, \sigma}^\dagger \hat{c}_{\mathbf{k}, \sigma} \sum_n |n\rangle \langle n| \hat{c}_{l, \sigma'}^\dagger \hat{c}_{l, \sigma'} |g\rangle$$

Identity

$$\hat{\mathcal{I}} = \sum_n |n\rangle \langle n|$$

Fock space

$$\langle n | g \rangle = \delta_{n, g}$$

# Density for different spin

Insert identity operator:

$$\rho(\mathbf{x}', \sigma') = \frac{1}{\mathcal{V}^2} \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}, \sigma}^\dagger \hat{c}_{\mathbf{k}, \sigma} | g \rangle \langle g | \hat{c}_{\mathbf{l}, \sigma'}^\dagger \hat{c}_{\mathbf{l}, \sigma'} | g \rangle$$

Fock space

$$\langle n | g \rangle = \delta_{n, g}$$

Number of particles operator

$$\hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} = \hat{n}_{\mathbf{k}\sigma}$$

## Density for different spin

Insert identity operator:

$$\rho(\mathbf{x}', \sigma') = \frac{1}{\mathcal{V}^2} \sum_{\mathbf{k}, l} \langle g | \hat{c}_{\mathbf{k}, \sigma}^\dagger \hat{c}_{\mathbf{k}, \sigma} | g \rangle \langle g | \hat{c}_{l, \sigma'}^\dagger \hat{c}_{l, \sigma'} | g \rangle$$

Number of particles operator

$$\hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} = \hat{n}_{\mathbf{k}\sigma}$$

$$\rho(\mathbf{x}', \sigma') = \frac{1}{\mathcal{V}^2} \left[ \sum_{\mathbf{k}} \langle g | \hat{n}_{\mathbf{k}\sigma} | g \rangle \right] \cdot \left[ \sum_l \langle g | \hat{n}_{l\sigma'} | g \rangle \right] = \left( \frac{N}{2\mathcal{V}} \right)^2$$

## Density for equal spin

If spins are equal

$$\sigma = \sigma' \implies \begin{cases} \mathbf{k} = \mathbf{n} \\ \mathbf{l} = \mathbf{m} \end{cases} \quad \text{or} \quad \begin{cases} \mathbf{k} = \mathbf{m} \\ \mathbf{l} = \mathbf{n} \end{cases} \quad \text{and} \quad \mathbf{k} \neq \mathbf{l}$$

We can simplify expression for density:

$$\rho(\mathbf{x}', \sigma) = \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{l}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \hat{c}_{\mathbf{l}\sigma} | g \rangle \psi_{\mathbf{k}\sigma}^*(\mathbf{x}) \psi_{\mathbf{l}\sigma}^*(\mathbf{x}') \psi_{\mathbf{k}\sigma}(\mathbf{x}') \psi_{\mathbf{l}\sigma}(\mathbf{x})$$

## Density for equal spin

If spins are equal

$$\sigma = \sigma' \implies \begin{cases} \mathbf{k} = \mathbf{n} \\ \mathbf{l} = \mathbf{m} \end{cases} \quad \text{or} \quad \begin{cases} \mathbf{k} = \mathbf{m} \\ \mathbf{l} = \mathbf{n} \end{cases} \quad \text{and} \quad \mathbf{k} \neq \mathbf{l}$$

We can simplify expression for density:

$$\rho(\mathbf{x}', \sigma) = \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{l}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \hat{c}_{\mathbf{l}\sigma} | g \rangle \psi_{\mathbf{k}\sigma}^*(\mathbf{x}) \psi_{\mathbf{l}\sigma}^*(\mathbf{x}') \psi_{\mathbf{k}\sigma}(\mathbf{x}') \psi_{\mathbf{l}\sigma}(\mathbf{x})$$

### Commutation for fermions

$$\{\hat{c}_{\mathbf{l}\sigma}^\dagger, \hat{c}_{\mathbf{k}\sigma}\} = \delta_{\mathbf{k}, \mathbf{l}} \implies \hat{c}_{\mathbf{l}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} = -\hat{c}_{\mathbf{k}\sigma} \hat{c}_{\mathbf{l}\sigma}^\dagger + \delta_{\mathbf{k}, \mathbf{l}}$$



## Density for equal spin

If spins are equal

$$\sigma = \sigma' \implies \begin{cases} \mathbf{k} = \mathbf{n} \\ \mathbf{l} = \mathbf{m} \end{cases} \quad \text{or} \quad \begin{cases} \mathbf{k} = \mathbf{m} \\ \mathbf{l} = \mathbf{n} \end{cases} \quad \text{and} \quad \mathbf{k} \neq \mathbf{l}$$

We can simplify expression for density:

$$\rho(\mathbf{x}', \sigma) = - \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \hat{c}_{\mathbf{l}\sigma}^\dagger \hat{c}_{\mathbf{l}\sigma} | g \rangle \psi_{\mathbf{k}\sigma}^*(\mathbf{x}) \psi_{\mathbf{k}\sigma}(\mathbf{x}') \psi_{\mathbf{l}\sigma}^*(\mathbf{x}') \psi_{\mathbf{l}\sigma}(\mathbf{x}) + \Delta$$

### Wave functions

$$\psi_{\mathbf{k}\sigma}(\mathbf{x}') = \frac{1}{\sqrt{\mathcal{V}}} e^{i\mathbf{k} \cdot \mathbf{x}'} \sigma$$

$$\psi_{\mathbf{k}\sigma}^*(\mathbf{x}) = \frac{1}{\sqrt{\mathcal{V}}} e^{-i\mathbf{k} \cdot \mathbf{x}} \sigma$$

## Density for equal spin

If spins are equal

$$\sigma = \sigma' \implies \begin{cases} \mathbf{k} = \mathbf{n} \\ \mathbf{l} = \mathbf{m} \end{cases} \quad \text{or} \quad \begin{cases} \mathbf{k} = \mathbf{m} \\ \mathbf{l} = \mathbf{n} \end{cases} \quad \text{and} \quad \mathbf{k} \neq \mathbf{l}$$

We can simplify expression for density:

$$\begin{aligned} \rho(\mathbf{x}', \sigma) = & -\frac{1}{V^2} \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \hat{c}_{\mathbf{l}\sigma}^\dagger \hat{c}_{\mathbf{l}\sigma} | g \rangle e^{-i\mathbf{k}(\mathbf{x}-\mathbf{x}')} e^{-i\mathbf{l}(\mathbf{x}'-\mathbf{x})} + \\ & + \frac{1}{V^2} \sum_{\mathbf{k}} \langle g | \hat{c}_{\mathbf{k},\sigma}^\dagger \hat{c}_{\mathbf{k},\sigma} \hat{c}_{\mathbf{k},\sigma}^\dagger \hat{c}_{\mathbf{k},\sigma} | g \rangle e^{-i\mathbf{k}\cdot\mathbf{x}} e^{i\mathbf{k}\cdot\mathbf{x}'} e^{i\mathbf{k}\cdot\mathbf{x}} e^{-i\mathbf{k}\cdot\mathbf{x}'} \end{aligned}$$

## Density for equal spin

If spins are equal

$$\sigma = \sigma' \implies \begin{cases} \mathbf{k} = \mathbf{n} \\ \mathbf{l} = \mathbf{m} \end{cases} \quad \text{or} \quad \begin{cases} \mathbf{k} = \mathbf{m} \\ \mathbf{l} = \mathbf{n} \end{cases} \quad \text{and} \quad \mathbf{k} \neq \mathbf{l}$$

We can simplify expression for density:

$$\rho(\mathbf{x}', \sigma) = -\frac{1}{\mathcal{V}^2} \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \hat{c}_{\mathbf{l}\sigma}^\dagger \hat{c}_{\mathbf{l}\sigma} | g \rangle e^{-i\mathbf{k} \cdot (\mathbf{x} - \mathbf{x}')} e^{-i\mathbf{l} \cdot (\mathbf{x}' - \mathbf{x})}$$

## Density for equal spin

Insert identity operator:

$$\rho(\mathbf{x}', \sigma) = -\frac{1}{V^2} \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \hat{\mathcal{I}} \hat{c}_{\mathbf{l}\sigma}^\dagger \hat{c}_{\mathbf{l}\sigma} | g \rangle e^{-i\mathbf{k}(\mathbf{x}-\mathbf{x}')} e^{-i\mathbf{l}(\mathbf{x}'-\mathbf{x})}$$

Identity

$$\hat{\mathcal{I}} = \sum_n |n\rangle \langle n|$$

## Density for equal spin

Insert identity operator:

$$\rho(\mathbf{x}', \sigma) = -\frac{1}{V^2} \sum_{\mathbf{k}, \mathbf{l}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \sum_n |n\rangle \langle n| \hat{c}_{\mathbf{l}\sigma}^\dagger \hat{c}_{\mathbf{l}\sigma} |g\rangle e^{-i\mathbf{k}(\mathbf{x}-\mathbf{x}')} e^{-i\mathbf{l}(\mathbf{x}'-\mathbf{x})}$$

Fock space

$$\langle n | g \rangle = \delta_{n,g}$$

## Density for equal spin

$$\rho(\mathbf{x}', \sigma) =$$

$$= -\frac{1}{\mathcal{V}^2} \left[ \sum_{\mathbf{k}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} | g \rangle e^{-i\mathbf{k}(\mathbf{x}' - \mathbf{x})} \right] \left[ \sum_{\mathbf{l}} \langle g | \hat{c}_{\mathbf{l}\sigma}^\dagger \hat{c}_{\mathbf{l}\sigma} | g \rangle e^{-i\mathbf{l}(\mathbf{x}' - \mathbf{x})} \right] + \left( \frac{N}{2\mathcal{V}} \right)^2$$

## Calculating sum

To continue we need to find

$$F(\mathbf{k}, \mathbf{r}) = \frac{1}{\mathcal{V}} \sum_{\mathbf{k}} e^{-i\mathbf{k}\mathbf{r}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} | g \rangle$$

## Calculating sum

To continue we need to find

$$F(\mathbf{k}, \mathbf{r}) = \frac{1}{\mathcal{V}} \sum_{\mathbf{k}} e^{-i\mathbf{k}\mathbf{r}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} | g \rangle$$



## Calculating sum

To continue we need to find

$$F(\mathbf{k}, \mathbf{r}) = \frac{1}{\mathcal{V}} \sum_{\mathbf{k}} e^{-i\mathbf{k}\mathbf{r}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} | g \rangle$$

Number of particles operator

$$\hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} = \hat{n}_{\mathbf{k}\sigma}$$

$$\langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} | g \rangle = \langle g | \hat{n}_{\mathbf{k}\sigma} | g \rangle = \begin{cases} 1 & k \leq k_F \\ 0 & k > k_F \end{cases}$$

## Calculating sum

To continue we need to find

$$F(\mathbf{k}, \mathbf{r}) = \frac{1}{\mathcal{V}} \sum_{\mathbf{k}} e^{-i\mathbf{k}\mathbf{r}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} | g \rangle$$

Heaviside step function

$$\langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} | g \rangle = H(k_F - |\mathbf{k}|)$$

## Calculating sum

To continue we need to find

$$F(\mathbf{k}, \mathbf{r}) = \frac{1}{\mathcal{V}} \sum_{\mathbf{k}} e^{-i\mathbf{k}\mathbf{r}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} | g \rangle$$

Change to summation inside the Fermi sphere

$$F(\mathbf{k}, \mathbf{r}) = \frac{1}{\mathcal{V}} \sum_{|\mathbf{k}| \leq k_F} e^{-i\mathbf{k}\mathbf{r}}$$

# From summation to integration

Multiply and divide by unit cell volume

$$F(\mathbf{k}, \mathbf{r}) = \frac{1}{\mathcal{V}} \sum_{|\mathbf{k}| \leq k_F} e^{-i\mathbf{k}\mathbf{r}} = \frac{1}{\mathcal{V}} \frac{\mathcal{V}}{8\pi^3} \sum_{|\mathbf{k}| \leq k_F} e^{-i\mathbf{k}\mathbf{r}} d^3\mathbf{k}$$


In the limit  $\mathcal{V} \rightarrow \infty$

$$F(\mathbf{k}, \mathbf{r}) = \frac{1}{8\pi^3} \sum_{|\mathbf{k}| \leq k_F} e^{-i\mathbf{k}\mathbf{r}} d^3\mathbf{k} \rightarrow \frac{1}{8\pi^3} \int_{|\mathbf{k}| \leq k_F} e^{-i\mathbf{k}\mathbf{r}} d^3\mathbf{k}$$

# Evaluation of the integral

We will integrate in spherical coordinates


$$F(\mathbf{k}, \mathbf{r}) = \frac{1}{8\pi^3} \int_{|\mathbf{k}| \leq k_F} e^{-i\mathbf{k}\mathbf{r}} d^3k$$



# Evaluation of the integral

We will integrate in spherical coordinates

$$F(\mathbf{k}, \mathbf{r}) = \frac{1}{8\pi^3} \int_{k=0}^{k_F} \int_{\theta=0}^{\pi} \int_{\phi=0}^{2\pi} e^{-i\mathbf{k}\mathbf{r} \cos \theta} k^2 \sin \theta \, d\phi d\theta dk$$



# Evaluation of the integral

Integration over  $\phi$

$$F(\mathbf{k}, \mathbf{r}) = \frac{1}{8\pi^3} \int_{k=0}^{k_F} \int_{\theta=0}^{\pi} \int_{\phi=0}^{2\pi} e^{-i\mathbf{k}\mathbf{r}\cos\theta} k^2 \sin\theta \, d\phi d\theta dk$$

$$F(\mathbf{k}, \mathbf{r}) = \frac{1}{4\pi^2} \int_{k=0}^{k_F} \int_{\theta=0}^{\pi} e^{-i\mathbf{k}\mathbf{r}\cos\theta} k^2 \sin\theta \, d\theta dk$$

# Evaluation of the integral

Integration over  $\theta$

$$F(\mathbf{k}, \mathbf{r}) = \frac{1}{4\pi^2} \int_{k=0}^{k_F} \int_{\theta=0}^{\pi} e^{-ikr \cos \theta} k^2 \sin \theta \, d\theta \, dk$$

$$F(\mathbf{k}, \mathbf{r}) = \frac{1}{4\pi^2} \int_{k=0}^{k_F} \int_{\theta=0}^{\pi} e^{-ikr \cos \theta} k^2 -d \cos \theta \, d\theta \, dk$$

$$F(\mathbf{k}, \mathbf{r}) = \frac{1}{4\pi^2} \int_{k=0}^{k_F} \int_{\eta=-1}^1 e^{ikr \eta} k^2 \, d\eta \, dk$$



# Evaluation of the integral

Integration over  $\theta$

$$F(k, \mathbf{r}) = \frac{1}{4\pi^2} \int_{k=0}^{k_F} \int_{\theta=0}^{\pi} e^{-ikr \cos \theta} k^2 \sin \theta \, d\theta \, dk$$

$$F(k, \mathbf{r}) = \frac{1}{4\pi^2} \int_{k=0}^{k_F} \int_{\theta=0}^{\pi} e^{-ikr \cos \theta} k^2 -d \cos \theta \, d\theta \, dk$$

$$F(k, \mathbf{r}) = \frac{1}{4\pi^2} \int_{k=0}^{k_F} \frac{2}{kr} \sin(kr) k^2 \, dk$$

# Evaluation of the integral

Integration over  $k$

$$F(k, \mathbf{r}) = \frac{1}{4\pi^2} \int_{k=0}^{k_F} \frac{2}{kr} \sin(kr) k^2 dk = \frac{1}{2\pi^2 r^3} \int_{k=0}^{k_F} \sin(kr) kr dk$$

$$F(k, \mathbf{r}) = \frac{1}{2\pi^2 r^3} \{ \sin(k_F r) - k_F r \cos(k_F r) \}$$

# Evaluation of the integral

Integration over  $k$

$$F(k, \mathbf{r}) = \frac{1}{4\pi^2} \int_{k=0}^{k_F} \frac{2}{kr} \sin(kr) k^2 dk = \frac{1}{2\pi^2 r^3} \int_{k=0}^{k_F} \sin(kr) kr dk$$

$$F(k, \mathbf{r}) = \frac{k_F^3}{2\pi^2 r^3 k_F^3} \{ \sin(k_F r) - k_F r \cos(k_F r) \}$$

Fermi wavenumber

$$k_F^3 = \frac{3\pi^2 N}{\mathcal{V}}$$

## Evaluation of the integral

$$\frac{1}{\mathcal{V}} \sum_{\mathbf{k}} e^{-i\mathbf{k}\mathbf{r}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} | g \rangle = \left( \frac{N}{2\mathcal{V}} \right) \frac{3 \{ \sin(k_F r) - k_F r \cos(k_F r) \}}{(k_F r)^3}$$

No divergence at zero: at  $k_F r \rightarrow 0 \iff r \ll \lambda_F$

$$3 \frac{\sin(k_F r) - k_F r \cos(k_F r)}{(k_F r)^3} \approx 3 \frac{x - x^3/6 - x(1 - x^2/2)}{x^3} = \frac{3}{3} = 1$$

## Bringing it together

$$\rho(\mathbf{x}', \sigma) = -\frac{1}{\mathcal{V}^2} \sum_{\mathbf{k}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} | g \rangle e^{i\mathbf{k} \cdot (\mathbf{x} - \mathbf{x}')} \sum_{\mathbf{l}} \langle g | \hat{c}_{\mathbf{l}\sigma}^\dagger \hat{c}_{\mathbf{l}\sigma} | g \rangle e^{i\mathbf{l} \cdot (\mathbf{x}' - \mathbf{x})}$$

$$\frac{1}{\mathcal{V}} \sum_{\mathbf{k}} e^{-i\mathbf{k}\mathbf{r}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} | g \rangle = \left( \frac{N}{2\mathcal{V}} \right) \frac{3 \{ \sin(k_F r) - k_F r \cos(k_F r) \}}{(k_F r)^3}$$

## Bringing it together

$$\rho(\mathbf{x}', \sigma) = -\frac{1}{\mathcal{V}^2} \sum_{\mathbf{k}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} | g \rangle e^{i\mathbf{k} \cdot (\mathbf{x} - \mathbf{x}')} \sum_{\mathbf{l}} \langle g | \hat{c}_{\mathbf{l}\sigma}^\dagger \hat{c}_{\mathbf{l}\sigma} | g \rangle e^{i\mathbf{l} \cdot (\mathbf{x}' - \mathbf{x})}$$

$$\frac{1}{\mathcal{V}} \sum_{\mathbf{k}} e^{-i\mathbf{k} \cdot \mathbf{r}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} | g \rangle = \left( \frac{N}{2\mathcal{V}} \right) \frac{3 \{ \sin(k_F r) - k_F r \cos(k_F r) \}}{(k_F r)^3}$$

## Bringing it together

$$\rho(\mathbf{x}', \sigma) = -\frac{1}{\mathcal{V}^2} \sum_{\mathbf{k}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} | g \rangle e^{i\mathbf{k} \cdot (\mathbf{x} - \mathbf{x}')} \sum_{\mathbf{l}} \langle g | \hat{c}_{\mathbf{l}\sigma}^\dagger \hat{c}_{\mathbf{l}\sigma} | g \rangle e^{i\mathbf{l} \cdot (\mathbf{x}' - \mathbf{x})}$$

$$\frac{1}{\mathcal{V}} \sum_{\mathbf{k}} e^{-i\mathbf{k}\mathbf{r}} \langle g | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} | g \rangle = \left( \frac{N}{2\mathcal{V}} \right) \frac{3 \{ \sin(k_F r) - k_F r \cos(k_F r) \}}{(k_F r)^3}$$

$$\rho(\mathbf{x}', \sigma) = \left( \frac{N}{2\mathcal{V}} \right)^2 - \left( \frac{N}{2\mathcal{V}} \right)^2 \left( \frac{3 \{ \sin(k_F r) - k_F r \cos(k_F r) \}}{(k_F r)^3} \right)^2$$

## Bringing it all together

$$\left(\frac{N}{2\mathcal{V}}\right)^2 g_{\sigma\sigma}(\mathbf{x}-\mathbf{x}') = \left(\frac{N}{2\mathcal{V}}\right)^2 - \delta_{\sigma\sigma'} \left(\frac{N}{2\mathcal{V}}\right)^2 \left(\frac{3\{\sin(k_F r) - k_F r \cos(k_F r)\}}{(k_F r)^3}\right)^2$$



## Bringing it all together

$$g_{\sigma\sigma}(\mathbf{x} - \mathbf{x}') = 1 - \delta_{\sigma\sigma'} \left( \frac{3 \{ \sin(k_F r) - k_F r \cos(k_F r) \}}{(k_F r)^3} \right)^2$$

## Visualization $\sigma = \sigma'$

