

TABLE e-1. SUMMARY OF STUDIES

Author and year (N=Number of patients)	Setting and time period	Type of study	Age in months	Sex	Dose and duration (treatment protocol)	Additional details	Definition of resolution	Probability of resolution
ADRENOCORTICOTROPIC HORMONE (ACTH)								
Hrachovy et al, 1983¹ (N=12)	Single center study at Houston, TX Period not specified, study published in 1983	Double-blind placebo-controlled study	Range: 3.5-24	Not specified	IM natural ACTH 20 IU/day	No prior treatment with ACTH or steroids	Spasm cessation and resolution of hypsarrhythmia at 2 weeks after treatment initiation	Clinical: 7/12 (0.5833) [Clinical + hypsarrhythmia: 7/12 (0.5833)]
Snead et al, 1989² (N=15)	Single center study at Birmingham, AL, USA 1984-1985	Prospective open trial	Mean: 7.5 Range: 2-18	M: 8 (53%) F: 7 (47%)	IM natural ACTH 150 IU/m ² /day	7 patients had other treatments at the time of ACTH	No spasms and normal EEG at 14 days	Clinical: 12/15 (0.8000) [Clinical + hypsarrhythmia: 12/15 (0.8000)]
Baram et al, 1996³ (N=15)	Single center study at Children's Hospital of Los Angeles, CA, USA Period not specified, study published in 1996	Randomized single-blinded study of IM ACTH high dose versus oral prednisone	Median (p ₂₅ -p ₇₅): 5 (3-6)	M: 4 (27%) F: 11 (73%)	IM natural ACTH 150 IU/m ² /day	2 patients had TSC	Spasm resolution 14 days after treatment initiation and resolution of hypsarrhythmia on EEG including a full sleep-wake cycle	Clinical: 14/15 (0.9333) [Clinical + hypsarrhythmia: 13/15 (0.8667)]
Vigevano & Cilio, 1997⁴ (N=19) [CLINICAL OUTCOME AT 20 DAYS]	Single center study at Rome, Italy 1992-1995	Randomized not blinded clinical trial (followed by a cross-over phase)	Mean (range): 5.3 (2-9)	M: 7 (37%) F: 12 (63%)	IM synthetic ACTH 10 IU once a day	1 patient discontinued ACTH because of side effects	Spasm resolution 20 days from treatment initiation	Clinical: 14/19 (0.7368) [Clinical + hypsarrhythmia: unknown]
Yanagaki et al, 1999⁵ (N=13)	Single center study at Tokyo, Japan 1991-1997	Randomized controlled trial high-dose versus low-dose ACTH	Mean (SD): Cryptogenic 5.8±2.6 Symptomatic 7.4±3.3	M: 8 (62%) F: 5 (39%)	IM synthetic ACTH 0.025 mg/Kg/day	Excluded patients previously treated with ACTH, corticosteroids, or intravenous immunoglobulin	Spasm resolution and disappearance of the hypsarrhythmia pattern within 2 weeks of treatment initiation	Clinical: 11/13 (0.8462) [Clinical + hypsarrhythmia: 11/13 (0.8462)]
Lux et al, 2004⁶ (N=25)	Multicenter study in the UK 1999-2002	Randomized non-blinded clinical trial	Median (p ₂₅ -p ₇₅): 5 (3.5-7)	M: 14 (56%) F: 11 (44%)	IM synthetic ACTH 0.5-0.75 mg every other day	Excluded patients with diagnosis or suspicion of TSC; 2 patients allocated to ACTH received prednisolone 4 patients received	Reported spasm resolution for at least 48 hours including day 13 and 14 of treatment	Clinical: 19/25 (0.7600) [Clinical + hypsarrhythmia: unknown]

Cost-Effectiveness Infantile Spasms

						lower dose due to adverse effects		
Lin et al, 2006 ⁷ (N=53)	Single-center study at National Taiwan University Hospital 1987-1998	Retrospective study	Age at spasm onset: Mean (SD): 8.2 (1.2) Age at treatment: Mean (SD): 11.1 (1.4)	M: 25 (47%) F: 28 (53%)	IM synthetic ACTH 2.5 IU/Kg/day	Excluded patients treated with ACTH or steroids before. Every patient received a three-day trial of 30-50 mg/Kg/day pyridoxine before ACTH was administered. 7 patients had TSC	Cessation of spasms and disappearance of hypsarrhythmia at 2 weeks	Clinical: 46/53 (0.8679) [Clinical + hypsarrhythmia: 46/53 (0.8679)]
Kossoff et al, 2008 ⁸ (N=20)	Single-center study at Johns Hopkins, Baltimore, MD 1996-2007	Retrospective study	Age at treatment: Median (p25-p75): 6.6 (5.9-7.9)	M: 13 (65%) F: 7 (35%)	IM natural ACTH 150 IU/m ² /day for 1 week and ACTH 75 IU/m ² /day for the second week	Inclusion of patients with other seizures prior to spasms unless they used an ASM for spasms	Complete absence of visible spasms by the family for at least a 24 hour period	Clinical: 17/20 (0.8500) [Clinical + hypsarrhythmia: 12/15 (0.8000)]
Cohen-Sadan et al, 2009 ⁹ (N=14)	Multicenter study in Israel 1985-2002	Multicenter retrospective study	Mean: 5 Range: 2-6.5	M: 9 (64%) F: 5 (36%)	IM natural ACTH 100 IU every other day or ACTH 20-40 IU every day	Patients with no abnormalities on MRI, normal metabolic work-up, no history of hypoxic-ischemic encephalopathy, and normal development prior to onset of spasms	Resolution of spasms at 2 weeks	Clinical: 11/14 (0.7857) [Clinical + hypsarrhythmia: 12/14 (0.8571)]
Wanigasinghe et al, 2015 ¹⁰ (N=49)	Single center study at Colombo, Sri Lanka 2010-2014	Single-center single-blind randomized trial	Mean (SD): 9.9 (8.7)	M: 31 (63%) F: 18 (37%)	IM synthetic ACTH 40-60 IU/every other day	13/49 patients had prior treatment with ASMs	Spasm freedom for at least 48 hours at day 14	Clinical: 18/49 (0.3674) [Clinical + hypsarrhythmia: 9/49 (0.1837)]
Knupp et al, 2016 ¹¹ (N=97)	Multicenter study in the USA 2012-2014	Prospective cohort study	Median: 6	M: 54 (56%) F: 43 (44%)	IM natural ACTH 150 IU/m ² /day (recommended dose)	Included patients with different etiologies, including TSC	Absence of spasms at 2 weeks from treatment initiation	Clinical: 66/97 (0.6804) [Clinical + hypsarrhythmia: unknown]
Hodgeman et al, 2016 ¹² (N=57)	Single-center study at Boston Children's Hospital, Boston, MA	Retrospective review	Median (p25-p75) age at onset: 6 (4.8-8.3)	M: 25 (44%) F: 32 (56%)	IM natural ACTH 150 IU/m ²	12 patients had prior treatments	Absence of spasms at day 14	Clinical: 40/57 (0.7018) [Clinical + hypsarrhythmia: unknown]

	2011-2015		Median (p25-p75) delay to treatment: 16 (7-50) days					mia: 40/57 (0.7018)]
Yin et al, 2017¹³ (N=111)	Single-center study at Xiangya Hospital, Changsha, China 2010-2016	Retrospective review	Age of onset: Mean: 5.9 SD: 4.4	M: 69 (62%) F: 42 (38%)	IM natural ACTH 2-4 IU/Kg/day (maximum 40 IU/day)	Excluded patients with a prior use of hormonal treatments	Absence of spasms at day 14	Clinical: 52/111 (0.4685) [Clinical + hypsarrhythmia: 46/111 (0.4144)]
Gowda et al, 2019¹⁴ (N=18)	Single center study at Indira Gandhi Institute of Child Health, Bangalore, India 2013-2015	Single-center prospective non-blinded randomized trial	Age of onset: Mean: 8.6 SD: 6.3 Age of treatment: Mean: 9.4 SD: 5.3	M: 12 (67%) F: 6 (33%)	IM natural ACTH 100 IU/m ² /day	No prior steroids	Absence of reported spasms at least 48 hours including days 13 and 14.	Clinical: 9/18 (0.5000) [Clinical + hypsarrhythmia: unknown]
Dressler et al, 2019¹⁵ (N=16) [CLINICAL OUTCOME AT 28 DAYS]	Single center study at Vienna, Austria 2008-2017	Randomized controlled trial of ACTH versus ketogenic diet	Median: 6	M: 10 (63%) F: 6 (38%)	IM synthetic ACTH 150 IU/m ² /day	Excluded patients with prior treatment with ketogenic diet or steroids	Spasm resolution and disappearance of the hypsarrhythmia at day 28 from treatment initiation	Clinical: 11/16 (0.6875) [Clinical + hypsarrhythmia: 11/16 (0.6875)]
PREDNISONE OR PREDNISOLONE								
Hrachovy et al, 1983¹ (N=12) [LOW DOSE]	Single center study at Houston, TX	Double-blind placebo-controlled study	Range: 3.5-24	Not specified	Oral prednisone 2 mg/Kg/day	No prior treatment with ACTH or steroids	Spasm cessation and resolution of hypsarrhythmia at 2 weeks after treatment initiation	Clinical: 5/12 (0.4167) [Clinical + hypsarrhythmia: 5/12 (0.4167)]
Baram et al, 1996³ (N=14) [LOW DOSE]	Single center study at Children's Hospital of Los Angeles, CA, USA	Randomized single-blinded study of im ACHT high dose versus oral prednisone	Median (p25-p75): 7 (5-8.75)	M: 8 (57%) F: 6 (43%)	Oral prednisone 2 mg/Kg/day	2 patients had TSC	Spasm resolution 14 days after treatment initiation and resolution of hypsarrhythmia on EEG including a full sleep-wake cycle	Clinical: 4/14 (0.2857) [Clinical + hypsarrhythmia: 4/14 (0.2857)]
Lux et al, 2004⁶ (N=30)	Multicenter study in the UK 1999-2002	Randomized non-blinded clinical trial	Median (p25-p75): 5 (4-6)	M: 18 (60%) F: 12 (40%)	Oral prednisolone 40-60mg/day	Excluded patients with diagnosis or suspicion of TSC 5 patients received lower dose due to adverse effects	Reported spasm resolution for at least 48 hours including day 13 and 14 of treatment	Clinical: 21/30 (0.7000) [Clinical + hypsarrhythmia: unknown]
Ware et al, 2012¹⁶ (N=17)	Single center study at Melbourne,	Single-center retrospective review	In the whole study of	In the whole study of	Oral prednisolone 40-	New-onset epileptic spasms	Absence of spasms at 2 weeks	Clinical: 13/17 (0.7647)

	Victoria, Australia 2007-2009		28 infants Range: 3-14	28 infants M: 17 (61%) F: 11 (39%)	60mg/day		No patients with TSC in this group	[Clinical + hypsarrhythmia: unknown]
Adhami & Harini, 2013 ¹⁷ (N=7)	Two center study at University of Massachusetts Medical Center and Boston Children's Hospital, Boston, MA 2011-2013	Two-center retrospective review	Age of treatment: Median (p ₂₅ -p ₇₅): 7.5 (6.9-9.3)	M: 2 (29%) F: 5 (71%)	Oral prednisolone minimum 4 mg/Kg/day (maximum 60 mg/day)		Absence of spasms on day 14	Clinical: 7/7 (1) [Clinical + hypsarrhythmia: 7/7 (1)]
Chellamuthu et al, 2014 ¹⁸ (N=32) [LOW DOSE]	Single center study at New Delhi, India 2012-2013	Single-center prospective randomized open-label trial	Median (p ₂₅ -p ₇₅): 10.5 (8-14.5)	M: 23 (72%) F: 9 (28%)	Oral prednisolone 2 mg/Kg/day	No patients with TSC	Spasm freedom for at least 48 hours at day 14	Clinical: 8/32 (0.2500) [Clinical + hypsarrhythmia: 7/32 (0.2188)]
Chellamuthu et al, 2014 ¹⁸ (N=31) High-dose	Single center study at New Delhi, India 2012-2013	Single-center prospective randomized open-label trial	Median (p ₂₅ -p ₇₅): 12 (9-18)	M: 21 (68%) F: 10 (32%)	Oral prednisolone 4 mg/Kg/day	No patients with TSC	Spasm freedom for at least 48 hours at day 14	Clinical: 16/31 (0.5161) [Clinical + hypsarrhythmia: 12/30 (0.4000)]
Wanigasinge et al, 2015 ¹⁰ (N=48)	Single center study at Colombo, Sri Lanka 2010-2014	Single-center single-blind randomized trial	Mean (SD): 8.3 (6.2)	M: 25 (52%) F: 23 (48%)	Oral prednisolone 40-60mg/day	12/48 patients had prior treatment with ASMs	Spasm freedom for at least 48 hours at day 14	Clinical: 28/48 (0.5833) [Clinical + hypsarrhythmia: 21/48 (0.4375)]
Yi et al, 2015 ¹⁹ (N=20)	Single center study at Jiangxi Children's Hospital, Nanchang, Jiangxi province, China 2011-2012	Single-center descriptive retrospective study	Mean: 16.5 Range: 3-53 Age of onset: Mean: 8.9 Range: 2.6-32	M: 17 (85%) F: 3 (15%)	Oral prednisone 40mg/day	No patients with TSC 6 patients had received at least two ASMs prior to prednisone	Absence of spasms at 2 weeks	Clinical: 16/20 (0.8000) [Clinical + hypsarrhythmia: 12/20 (0.6000)]
Knupp et al, 2016 ¹¹ (N=54)	Multicenter study in the USA 2012-2014	Prospective cohort study	Median: 7	M: 26 (48%) F: 28 (52%)	Oral corticosteroids: Prednisolone 40mg/day (recommended dose)	Included patients with different etiologies, including TSC	Absence of spasms at 2 weeks from treatment initiation	Clinical: 30/54 (0.5556) [Clinical + hypsarrhythmia: unknown]
Gonzalez-Giraldo et al, 2018 ²⁰ (N=87) First 15 patients also	Single center study at Johns Hopkins Hospital, Baltimore, MD, USA 2006-2016	Single-center retrospective review	Age of onset: Median: 6 Range: 1-32	M: 49 (56%) F: 38 (44%)	Oral prednisolone 40-60 mg/day	Prednisolone was the first treatment or the second treatment after failure of 2 weeks of	Absence of spasms and absence of hypsarrhythmia at 2 weeks	Clinical: 62/87 (0.7126) [Clinical + hypsarrhythmia:]

reported in Kossoff et al, 2009 ²¹						ketogenic diet		56/87 (0.6437)]
Eliyan et al, 2019 ²² (N=102) First 27 patients also reported in Hussain et al, 2014 ²³	Single center study at Mattel Children's Hospital, Los Angeles, CA 2009-2017	Single-center retrospective review	Age of onset: Median (p ₂₅ -p ₇₅): 7.1 (4.9-12.1)	M: 61 (60%) F: 41 (40%)	Oral prednisolone 8 mg/Kg/day with a maximum of 60 mg/day	5 patients had TSC Prior treatment with short or low-dose treatment with prednisolone or ACTH in 12 patients. Prior treatment with VGB in 36 patients	Absence of spasms and absence of hypsarrhythmia (with overnight EEG) at 2 weeks and no recurrence of spasms over 28 days	Clinical: 60/102 (0.5882) [Clinical + hypsarrhythmia: 60/102 (0.5882)]
Yi et al, 2019 ²⁴ (N=39)	Single center study at Jiangxi Children's Hospital, Nanchang, Jiangxi province, China 2015-2016	Single-center prospective randomized trial comparing prednisone with prednisone and topiramate	Age of onset: Median: 6 Range: 2-39 Age of treatment: Median: 9.2 Range: 3.5-40	M: 26 (67%) F: 13 (33%)	Oral prednisone 40mg/day (increased to 60mg/day if spasms sustained by day 7)	No prior hormone therapy 1/39 (3%) patient had TSC	Absence of spasms for at least 48 hours at 2 weeks	Clinical: 28/39 (0.7180) [Clinical + hypsarrhythmia: 21/39 (0.5385)]
Gowda et al, 2019 ¹⁴ (N=15)	Single center study at Indira Gandhi Institute of Child Health, Bangalore, India 2013-2015	Single-center prospective non-blinded randomized trial	Age of onset: Mean: 6 SD: 5.2 Age of treatment: Mean (SD): 13.9 (9.2)	M: 9 (60%) F: 6 (40%)	Oral prednisolone 4mg/Kg/day (maximum 60mg/day)	No prior steroids	Absence of reported spasms at least 48 hours including days 13 and 14.	Clinical: 5/15 (0.3333) [Clinical + hypsarrhythmia: 4/15 (0.2667)]

Data in rows marked as [LOW DOSE] were not used for the main analysis as they were considered underdosed with the drug as explained at https://ivansanchezfernandez.github.io/InfantileSpasms_low_dose_steroids/. Data in rows marked as [CLINICAL OUTCOME AT 20 DAYS] or [CLINICAL OUTCOME AT 28 DAYS] were not used for the main analysis as these studies did not report clinical resolution of spasms at 14 days. When there were several doses used for ACTH in the same study, we only considered the highest one.

Legend: ACTH: Adrenocorticotrophic hormone. ASM: Anti-seizure medication. EEG: Electroencephalogram. F: Female. IM: Intramuscular. IU: International units. Kg: Kilogram. M: Male.

mg: milligram. **MRI:** Magnetic resonance imaging. **SD:** Standard deviation. **TSC:** Tuberous sclerosis complex. **UK:** United Kingdom. **USA:** United States of America. **VGB:** Vigabatrin.

REFERENCES

1. Hrachovy RA, Frost JD, Jr., Kellaway P, Zion TE. Double-blind study of ACTH vs prednisone therapy in infantile spasms. *J Pediatr* 1983;103:641-645.
2. Snead OC, 3rd, Benton JW, Jr., Hosey LC, et al. Treatment of infantile spasms with high-dose ACTH: efficacy and plasma levels of ACTH and cortisol. *Neurology* 1989;39:1027-1031.
3. Baram TZ, Mitchell WG, Tournay A, Snead OC, Hanson RA, Horton EJ. High-dose corticotropin (ACTH) versus prednisone for infantile spasms: a prospective, randomized, blinded study. *Pediatrics* 1996;97:375-379.
4. Vigeveno F, Cilio MR. Vigabatrin versus ACTH as first-line treatment for infantile spasms: a randomized, prospective study. *Epilepsia* 1997;38:1270-1274.
5. Yanagaki S, Oguni H, Hayashi K, et al. A comparative study of high-dose and low-dose ACTH therapy for West syndrome. *Brain Dev* 1999;21:461-467.
6. Lux AL, Edwards SW, Hancock E, et al. The United Kingdom Infantile Spasms Study comparing vigabatrin with prednisolone or tetracosactide at 14 days: a multicentre, randomised controlled trial. *Lancet* 2004;364:1773-1778.
7. Lin HC, Young C, Wang PJ, Lee WT, Shen YZ. ACTH therapy for Taiwanese children with West syndrome -- efficacy and impact on long-term prognosis. *Brain Dev* 2006;28:196-201.
8. Kossoff EH, Hedderick EF, Turner Z, Freeman JM. A case-control evaluation of the ketogenic diet versus ACTH for new-onset infantile spasms. *Epilepsia* 2008;49:1504-1509.
9. Cohen-Sadan S, Kramer U, Ben-Zeev B, et al. Multicenter long-term follow-up of children with idiopathic West syndrome: ACTH versus vigabatrin. *European journal of neurology* 2009;16:482-487.
10. Wanigasinghe J, Arambepola C, Sri Ranganathan S, Sumanasena S, Attanapola G. Randomized, Single-Blind, Parallel Clinical Trial on Efficacy of Oral Prednisolone Versus Intramuscular Corticotropin on Immediate and Continued Spasm Control in West Syndrome. *Pediatric neurology* 2015;53:193-199.
11. Knupp KG, Coryell J, Nickels KC, et al. Response to treatment in a prospective national infantile spasms cohort. *Ann Neurol* 2016;79:475-484.
12. Hodgeman RM, Kapur K, Paris A, et al. Effectiveness of once-daily high-dose ACTH for infantile spasms. *Epilepsy & behavior : E&B* 2016;59:4-8.
13. Yin J, Lu Q, Yin F, et al. Effectiveness and Safety of Different Once-Daily Doses of Adrenocorticotrophic Hormone for Infantile Spasms. *Paediatric drugs* 2017;19:357-365.
14. Gowda VK, Narayanaswamy V, Shivappa SK, Benakappa N, Benakappa A. Corticotrophin-ACTH in Comparison to Prednisolone in West Syndrome - A Randomized Study. *Indian journal of pediatrics* 2019;86:165-170.
15. Dressler A, Benninger F, Trimmel-Schwahofner P, et al. Efficacy and tolerability of the ketogenic diet versus high-dose adrenocorticotrophic hormone for infantile spasms: A single-center parallel-cohort randomized controlled trial. *Epilepsia* 2019;60:441-451.
16. Ware TL, Mackay MT, Harvey AS, Freeman JL. Epileptic spasms: experience with a high-dose oral corticosteroid protocol. *Journal of paediatrics and child health* 2012;48:985-989.
17. Adhami S, Harini C. High dose oral steroid as successful initial treatment for infantile spasms: follow-up of seven cases. *Epilepsy Currents* 2013;13:389-390.
18. Chellamuthu P, Sharma S, Jain P, Kaushik JS, Seth A, Aneja S. High dose (4 mg/kg/day) versus usual dose (2 mg/kg/day) oral prednisolone for treatment of infantile spasms: an open-label, randomized controlled trial. *Epilepsy Res* 2014;108:1378-1384.
19. Yi ZS, Wu HP, Yu XY, Chen Y, Zhong JM. Efficacy and tolerability of high-dose prednisone in Chinese children with infantile spasms. *Brain Dev* 2015;37:23-28.
20. Gonzalez-Giraldo E, Stafstrom CE, Stanfield AC, Kossoff EH. Treating Infantile Spasms with High-Dose Oral Corticosteroids: A Retrospective Review of 87 Children. *Pediatric neurology* 2018;87:30-35.

21. Kossoff EH, Hartman AL, Rubenstein JE, Vining EP. High-dose oral prednisolone for infantile spasms: an effective and less expensive alternative to ACTH. *Epilepsy & behavior : E&B* 2009;14:674-676.
22. Eliyan Y, Heesch J, Alayari A, Rajaraman RR, Sankar R, Hussain SA. Very-High-Dose Prednisolone Before ACTH for Treatment of Infantile Spasms: Evaluation of a Standardized Protocol. *Pediatric neurology* 2019.
23. Hussain SA, Shinnar S, Kwong G, et al. Treatment of infantile spasms with very high dose prednisolone before high dose adrenocorticotrophic hormone. *Epilepsia* 2014;55:103-107.
24. Yi Z, Wu H, Yu X, et al. High-dose prednisone therapy for infantile spasms and late-onset epileptic spasms in China: The addition of topiramate provides no benefit. *Seizure* 2019;71:174-178.