SEARCH STRATEGY

Inclusion and exclusion criteria for studies.

PICO	Inclusion Criteria	Exclusion Criteria
Population	• Infants (0 to 24 months of age)	• Older children (> 24 months)
	with infantile spasms	Studies only on specific subgroups
		(only Down syndrome, only tuberous
		sclerosis complex, etc.)
Intervention	Oral prednisone or prednisolone	Other steroids
Comparison	Intramuscular ACTH	Other steroids
Outcomes	Cessation of clinical spasms at	 Lack of information on clinical
	14 days after initiation of	outcome
	treatment	

<u>Systematic search of the literature.</u> Our search strategy was registered at PROSPERO (https://www.crd.york.ac.uk/PROSPEROFILES/152181 STRATEGY 20191129.pdf) ¹ and performed on November, 29, 2019. The search strategy included:

-PubMed ("Spasms, Infantile"[mesh] OR infantile spasm*[tiab] OR west syndrome[tiab] OR neonatal spasm*[tiab]) AND ("Adrenocorticotropic Hormone"[Mesh] OR corticotrophin[tiab] OR corticotrophin[tiab] OR adrenocorticotrophin[tiab] OR adrenocorticotrophin[tiab] OR adrenocorticotrophin[tiab] OR "Prednisolone"[Mesh:noexp] OR "Prednisone"[Mesh] OR Predniso*[tiab] OR dacortin[tiab])

-Embase ('infantile spasm'/exp OR 'infantile spasm*':ab,ti OR 'west syndrome':ab,ti OR 'neonatal spasm*':ab,ti) AND ('prednisolone'/exp OR 'corticotropin'/exp OR 'prednisone'/exp OR corticotrophin:ab,ti OR corticotropin:ab,ti OR 'adrenocroticotrophic hormone':ab,ti OR adrenocorticotropin:ab,ti OR acth:ab,ti OR predniso*:ab,ti OR dacortin:ab,ti)

-Web of Science (TS=("infantile spasm*" OR "west syndrome" OR "neonatal spasm*") AND

TS=("corticotrophin" OR "corticotropin" OR "adrenocroticotrophic hormone" OR "adrenocorticotropin"

OR "adrenocorticotrophin" OR "acth" OR "predniso*" OR "dacortin"))




Figure. Search strategy.

REFERENCES

1. National Institute for Health Research. PROSPERO. International prospective register of systematic reviews. URL for our search strategy: https://www.crd.york.ac.uk/PROSPEROFILES/152181_STRATEGY_20191129.pdf. 2019.