MediFind PH: OTC Medicine Product Leaflets

Volume 1: Pain, Fever and Muscle Aches

29 product leaflets compiled in registration-file order. Each leaflet is included unchanged after this contents page and has its own bookmark.

No.	Product	Reg. No.	Page
1	Advil (Ibuprofen) 200 mg Softgel Capsule	DR-XY32938	2
2	Advil (Ibuprofen) 100 mg/5 mL Suspension	DR-XY29056	3
3	Alaxan (Ibuprofen + Paracetamol) 200 mg/325 mg Tablet	DR-XY37938	4
4	Alaxan FR (Ibuprofen + Paracetamol) 200 mg / 325 mg Capsule	DR-XY37939	5
5	Biogesic (Paracetamol) 250 mg / 5 mL Syrup (Orange Flavor)	DR-XY45639	6
6	Biogesic (Paracetamol) 250 mg/5mL Syrup (Melon Flavor) Syrup	DR-XY46366	12
7	Biogesic (Paracetamol) 500 mg Tablet	DR-153	18
8	Biogesic 325 (Paracetamol) 325 mg Tablet	DRP-8763	19
9	Calpol (Paracetamol) 250 mg / 5 mL Suspension	DRHR-1964	20
10	Calpol (Paracetamol) 120mg/5mL Suspension (Strawberry Flavor)	DRHR-2131	21
11	Dolfenal (Mefenamic Acid) 250 mg Film-Coated Tablet	DRP-8716	22
12	Efficascent (Menthol + Methyl Salicylate + Camphor) 140 mg/100 mg/35 mg per gram Ointment	DRHR-2201	29
13	Efficascent Oil (Methyl Salicylate + Camphor + Menthol) 168.73 mg / 38.46 mg / 4.23 mg per mL Liniment	DRHR-2119	31
14	Flanax (Naproxen Sodium) 220 mg Film-Coated Tablet	DR-XY46695	33
15	Flanax (Naproxen Sodium) 275 mg Film-Coated Tablet	DR-XY46696	38
16	FlanaxACE (Menthol + Tocopherol Acetate (Vitamin E) + Glycol Salicylate + dl-Camphor) 4.98/28/1.48/900 mg per 100 g Transdermal Patch	DRHR-2412	39
17	Katinko (Isopropyl Alcohol) 70% (v/v) Solution	DRHR-2243	40
18	KATINKO (Menthol) 42.7 mg per 100 mg Ointment Stick	DRHR-1935	50
19	Katinko (Methyl Salicylate+Menthol+Camphor) 8mg/7.5mg/2.6mg per 100mg Ointment Stick	DRHR-1151	54
20	Medicol (Ibuprofen) 200 mg Tablet	DR-XY24958	58
21	Medicol I-400 (Ibuprofen) 400 mg Tablet	DR-XY24952	59
22	Rexidol Forte (Paracetamol + Caffeine) 500 mg/ 65 mg Tablet	DR-XY46238	60
23	Salonpas (Methyl Salicylate + l-Menthol + Tocopherol Acetate (Vitamin E) + dl-Camphor) 36 mg/ 33 mg/ 12 mg/ 7.1 mg Medicated Patch	DRHR-821	61
24	Salonpas Jet Spray (Methyl Salicylate + Menthol) 100 mg 30 mg Aerosol Spray	DRHR-2304	62
25	Saridon (Paracetamol + Caffeine) 500 mg/50 mg Tablet	DR-XY46456	63
26	Saridon Triple Action (Paracetamol + Propyphenazone + Caffeine) 250 mg/150 mg/50 mg Tablet	DR-XY36639	68
27	Saridon Max (Naproxen Sodium) 275 mg Film-Coated Tablet	DRP-1872-02	73
28	Tempra (Paracetamol) 325 mg Tablet	DR-4396	74
29	Tempra Blizz (Paracetamol) 500 mg Effervescent Tablet (Orange Flavor)	DR-XY48911	76

Advil (Ibuprofen)

200 mg Softgel Capsule | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY32938

Advil: Official FDA Philippines registration record

Brand name	Advil
Generic name	Ibuprofen
Dosage strength and form	200 mg Softgel Capsule
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Anti-inflammatory and Antirheumatic, Non-steroids (Propionic Acid derivative)
Registration number	DR-XY32938
Manufacturer	Catalent Pharma Solutions, LLC

Advil: OTC forms registered under the Advil brand (capsule availability)

OTC capsule form registered under the Advil brand: YES. 2 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Advil	100 mg/5 mL Suspension	Over-the-Counter (OTC) Drug	DR-XY29056
Advil	200 mg Softgel Capsule	Over-the-Counter (OTC) Drug	DR-XY32938

Advil: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY32938 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Advil"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Advil (Ibuprofen)

100 mg/5 mL Suspension | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY29056

Advil: Official FDA Philippines registration record

Brand name	Advil
Generic name	Ibuprofen
Dosage strength and form	100 mg/5 mL Suspension
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Analgesic/Antipyretic
Registration number	DR-XY29056
Manufacturer	Interphil Laboratories, Inc.

Advil: OTC forms registered under the Advil brand (capsule availability)

OTC capsule form registered under the Advil brand: YES. 2 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Advil	100 mg/5 mL Suspension	Over-the-Counter (OTC) Drug	DR-XY29056
Advil	200 mg Softgel Capsule	Over-the-Counter (OTC) Drug	DR-XY32938

Advil: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY29056 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Advil"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Alaxan (Ibuprofen + Paracetamol)

200 mg/325 mg Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY37938

Alaxan: Official FDA Philippines registration record

Brand name	Alaxan
Generic name	Ibuprofen + Paracetamol
Dosage strength and form	200 mg/325 mg Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Non-Steroidal Anti-Inflammatory Drug (NSAID)/Analgesic/Antipyretic
Registration number	DR-XY37938
Manufacturer	PT. Medifarma Laboratories

Alaxan: OTC forms registered under the Alaxan brand (capsule availability)

OTC capsule form registered under the Alaxan brand: YES. 3 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Alaxan FR	200 mg / 325 mg Capsule	Over-the-Counter (OTC) Drug	DR-XY37939
Alaxan	200 mg/325 mg Tablet	Over-the-Counter (OTC) Drug	DR-XY37938
AlaxanR Xtra (for BNC)	325 mg / 200 mg (equivalent to 256.23 mg Ibuprofen as Sodium dihydrate) Tablet	Over-the-Counter (OTC) Drug	DR-XY48927

Alaxan: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY37938 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Alaxan"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Alaxan FR (Ibuprofen + Paracetamol)

200 mg / 325 mg Capsule | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY37939

Alaxan FR: Official FDA Philippines registration record

Brand name	Alaxan FR
Generic name	Ibuprofen + Paracetamol
Dosage strength and form	200 mg / 325 mg Capsule
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Non-Steroidal Anti-Inflammatory Drug (NSAID/Analgesic/Antipyretic)
Registration number	DR-XY37939
Manufacturer	PT. Medifarma Laboratories

Alaxan FR: OTC forms registered under the Alaxan brand (capsule availability)

OTC capsule form registered under the Alaxan brand: YES. 3 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Alaxan FR	200 mg / 325 mg Capsule	Over-the-Counter (OTC) Drug	DR-XY37939
Alaxan	200 mg/325 mg Tablet	Over-the-Counter (OTC) Drug	DR-XY37938
AlaxanR Xtra (for BNC)	325 mg / 200 mg (equivalent to 256.23 mg Ibuprofen as Sodium dihydrate) Tablet	Over-the-Counter (OTC) Drug	DR-XY48927

Alaxan FR: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY37939 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Alaxan"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Biogesic (Paracetamol)

250 mg / 5 mL Syrup (Orange Flavor) | Household Remedy (HR) | FDA Reg. No. DR-XY45639

Biogesic: Official FDA Philippines registration record

Brand name	Biogesic
Generic name	Paracetamol
Dosage strength and form	250 mg / 5 mL Syrup (Orange Flavor)
Classification	Household Remedy (HR)
Pharmacologic category	Analgesic / Antipyretic (Anilide)
Registration number	DR-XY45639
Manufacturer	Amherst Laboratories, Inc.

Biogesic: OTC forms registered under the Biogesic brand (capsule availability)

OTC capsule form registered under the Biogesic brand: NO. 7 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Biogesic	100 mg / mL Suspension (Oral Drops) [Orange Sugar Free]	Household Remedy (HR)	DRHR-1167
Biogesic	120 mg/5 mL Syrup (Melon Flavor)	Household Remedy (HR)	DRHR-1843
Biogesic	120 mg/5 mL Syrup (Orange Flavor)	Household Remedy (HR)	DRHR-1748
Biogesic	250 mg / 5 mL Syrup (Orange Flavor)	Household Remedy (HR)	DR-XY45639
Biogesic	250 mg/5mL Syrup (Melon Flavor) Syrup	Over-the-Counter (OTC) Drug	DR-XY46366
Biogesic 325	325 mg Tablet	Over-the-Counter (OTC) Drug	DRP-8763
Biogesic	500 mg Tablet	Over-the-Counter (OTC) Drug	DR-153

Biogesic: Official leaflet (Package Insert / Patient Information Leaflet)

An official leaflet for this registration is published on the FDA Verification Portal: PIL_01 (https://drive.google.com/file/d/17IR_yvQ0Nx1FJcxvlVe97VmM_lFSJcmz/view?usp=drivesdk). Its text (OCR) and the original leaflet pages follow.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Biogesic"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Biogesic: Official leaflet text (OCR transcription)

Machine transcription of the official FDA leaflet for Reg. No. DR-XY45639. OCR may contain errors; the original leaflet pages are attached after this section.

Biogesic leaflet PIL_01

FORMULATION
Each mL syrup (oral drops) contains: Paracetamol…….....…….100 mg
Each 5 mL syrup contains: Paracetamol…….....….120 mg or 250 mg
PRODUCT DESCRIPTION
Paracetamol (Biogesic®) 100 mg/mL Syrup (Oral Drops), 120 mg/5 mL Syrup, and 250 mg/5 mL Syrup (Orange Flavor)
• An orange-colored, clear syrup with an orange odor and taste
Paracetamol (Biogesic®) 120 mg/5 mL Syrup, and 250 mg/5 mL Syrup (Melon Flavor)
• An orange-colored, clear syrup with a melon odor and taste
These products use the TasteRite® technology of PediaTech®. The TasteRite® technology is a unique tastemasking system developed specifically for liquid dosage forms.
This technology significantly reduces the bitterness of medicine so that children taste the flavor and not the medicine.
In addition, these products use Tooth Friendly technology TFTech® to significantly reduce or eliminate enamel loss that leads to tooth decay.
WHAT IS IN THE MEDICINE?
This product contains paracetamol, a fever reducer and pain reliever.
STRENGTH OF THE MEDICINES
See Formulations
WHAT IS THE MEDICINE USED FOR?
HOW MUCH AND HOW OFTEN SHOULD YOU USE THIS MEDICINE?
General Recommendations:
• Infants and children should not exceed 5 doses in each 24-hour period.
• Leave at least 4 hours between doses.
• Do not use continuously for more than 5 days at a time unless directed by a doctor.
• Do not use more than the recommended dose.
• Avoid potential liver damage by giving lower doses of paracetamol in patients classified under the lower body weight group.
This medicine should be taken orally every 4 to 6 hours, as needed for pain and/or fever, or, as directed by a doctor.
Dosing in Children: 10 mg/kg body weight per dose given every 4 to 6 hours, as needed; or, you can follow the dose in the dosing table below:
WHEN SHOULD YOU NOT TAKE THIS MEDICINE?
• If you are allergic to paracetamol or any ingredient of the product
• If you are taking other medicines that contain paracetamol
UNDESIRABLE EFFECTS
Paracetamol, when taken within the recommended dose and duration of treatment, has low incidence of side effects.
Blood and lymphatic system disorders: Anemia (decreased number of healthy red blood cells), changes in the number of white blood cells and platelets such as agranulocytosis (decrease in the number of granulocytes), leukopenia (decrease in the number of leukocytes), methemoglobinemia (presence of methemoglobin in the blood), neutropenia (decrease in the number of neutrophils), pancytopenia (deficiency of all cellular elements in the blood), thrombocytopenia (decrease in the number of platelets); thrombocytopenic purpura (small purple-red marks in the skin or other tissue caused by low number of platelets)
Immune system disorders: Allergic reactions (some are life-threatening) which may cause difficulty swallowing, unexplained wheezing, shortness of breath or difficulty in breathing, skin rash, hives, or angioedema (swelling of face, tongue, lips, or throat); proneness to infections, swelling around the eyes
Metabolism and nutrition disorders: Hyperchloremia (increased chlorine ion in the blood), hyperuricemia (increased uric acid in the blood), loss of appetite, metabolic acidosis (too much acid in the body), sudden weight loss
Nervous system disorders: Headache
Vascular disorders: Unusual bleeding or bruising
Respiratory, thoracic, and mediastinal disorders: Respiratory distress (noisy, congested breathing), runny nose, laryngeal edema (swelling of the voice box)
Gastrointestinal disorders: Mouth ulcers, nausea, sore throat, vomiting
Skin and subcutaneous tissue disorder: Fixed eruption (an allergic reaction described as well-defined, round or oval patches of redness and swelling in skin that recurs in the same site each time a particular drug is taken), itchy rash, severe skin reactions (i.e., Stevens-Johnson syndrome, toxic epidermal necrolysis, and acute generalized exanthematous pustulosis) which may include symptoms such as skin redness, blisters, rash, itching, and peeling of the upper surface of the skin
Hepatobiliary disorders: Chronic hepatic necrosis (death of liver tissues) in prolonged use, liver dysfunction, liver injury, yellowing of eyes and skin
Renal and urinary disorders: Kidney toxicity, papillary necrosis in prolonged use
General disorders and administration site conditions: Fatigue or unusual tiredness
Investigations: Decreased serum bicarbonate, serum calcium, serum sodium; increases in serum alkaline

phosphatase, serum glucose, and transaminase levels
Do not use with any other medicine containing paracetamol (prescription or nonprescription). If you are not sure whether a medicine contains paracetamol, ask a doctor.
Prolonged use of paracetamol with other painkillers [e.g., drugs called nonsteroidal anti-inflammatory drugs (NSAIDs), aspirin] is not recommended since it may increase the risk of toxic effects to the kidneys and eventually of kidney failure.
Chronic intake of alcohol may increase the risk of paracetamol-induced liver toxicity. Acute intake of alcohol may diminish the ability of the body to metabolize large doses of paracetamol. It is advised to consult a doctor before intake of paracetamol.
WHAT OTHER MEDICINE OR FOOD SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?
•
Reg. IPOPHIL P300000040493
8-864522-1 8-UNILAB-1
100 mg/mL Syrup (Oral Drops)
120 mg/5 mL Syrup
250 mg/5 mL Syrup
ANALGESIC / ANTIPYRETIC (Anilide)
The absorption of paracetamol may be increased by metoclopramide or domperidone. Cholestyramine, narcotic pain relievers, and some antidepressants that slow down gastric emptying reduce the absorption of paracetamol. If you are taking these medicines, talk to your doctor first before taking paracetamol.
Medicines which stimulate the enzymes responsible for the metabolic activation of paracetamol such as medicines for convulsion (e.g., phenobarbital), depression (e.g., tricyclic antidepressants), and oral contraceptives may reduce the efficacy of paracetamol and increase susceptibility to the harmful effects to the liver.
Phenothiazines (medicines for serious mental and emotional disorder) may increase the risk of severe hypothermia (abnormally low body temperature)
Paracetamol increases exposure to chloramphenicol. This interaction should be considered when both medicines are given together in malnourished patients.
Probenecid may affect the elimination of paracetamol.
There may be an increased risk of bleeding when using paracetamol and warfarin (a blood thinning medicine) at the same time. Paracetamol, especially its prolonged regular use, increases the blood thinning effect of warfarin and other coumarins. If you are taking warfarin, ask the doctor first before taking paracetamol.
MEDICINE USES
100 mg/mL Syrup (Oral Drops)
•
• For fever reduction.
120 mg/5 mL Syrup and
250 mg/5 mL Syrup
• For the relief of minor aches and pains due to the common cold, flu, headache, teething, sore throat, earache, post-immunization reactions, and toothache
WHEN SHOULD YOU CONSULT YOUR DOCTOR?
Ask a doctor before use if:
• The patient has kidney or liver problems (including alcoholic liver disease)
• The patient is taking other pain killers or any other medicines (See What Other Medicine or Food Should Be Avoided While Taking This Medicine?)
• The patient has G6PD
• The cause of the patient’s illness is not clear or it is accompanied by a rash, breathing difficulties, diarrhea or excessive tiredness or lethargy
• The patient is severely malnourished, anorexic, have a low body mass index (underweight), or excessively drinks alcohol
Stop use and ask a doctor if:
• If symptoms of allergic reaction occur (such as skin redness, blisters, or rash), stop use and seek medical help right away (See Care that Should be Taken when Taking this Medicine).
• Symptoms worsen or do not improve
• New symptoms occur
• Any undesirable effect occurs
• Redness or swelling is present
• Fever gets worse or lasts more than 3 days
• Pain gets worse or lasts more than 5 days in children (<12 years old) and 10 days in adults
• Patient had unexpected bruising or bleeding, and had more infections than usual
• Consult a doctor promptly if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting.
ADVERSE DRUG REACTION REPORTING STATEMENT
For suspected adverse drug reaction, seek medical attention immediately and report to the FDA at www.fda.gov.ph AND Unilab at +632-8-UNILAB-1 (+632-8-864522-1) for Metro Manila or toll-free +1-800-10-UNILAB-1 for provinces, or e-mail productsafety@unilab.com.ph. By reporting undesirable effects, you can help provide more information on the safety of this medicine.

AVAILABILITY
Paracetamol (Biogesic®) 100 mg/mL Syrup Orange (DRHR-1747)
- Pediatech HDPE Plastic Bottle with Child Resistant Plastic Cap x 15 mL (Box of 1's)
Paracetamol (Biogesic®) 120 mg/5 mL Syrup Orange (DRHR-1748)
- Pediatech HDPE Plastic Bottle with Child Resistant Plastic Cap x 60 mL (Box of 1's)
Paracetamol (Biogesic®) 120 mg/5 mL Syrup Melon (DRHR-1843)
Paracetamol (Biogesic®) 250 mg/5 mL Syrup Orange (DR-XY45639)
Paracetamol (Biogesic®) 250 mg/5 mL Syrup Melon (DR-XY46366)
Manufactured by AMHERST LABORATORIES, INC.
UNILAB Pharma Campus, Barangay Mamplasan
Biñan, Laguna, Philippines
for UNILAB, Inc.
No. 66 United Street, Mandaluyong City
Metro Manila, Philippines
DATE OF FIRST AUTHORIZATION
Paracetamol (Biogesic®) - Orange Flavor: November 2016
Paracetamol (Biogesic®) – Melon Flavor: June 2018
DATE OF REVISION OF PATIENT INFORMATION LEAFLET: May 2023 (CCDS 17-047)
For the relief of minor aches and pains due to the common cold, flu,
sore throat, teething, headache, earache, and post-immunization
reactions
Tell your doctor if you are taking other medicines (especially other paracetamol-containing medicines), including vitamins, supplements, and herbal medicines.
WHAT SHOULD YOU DO IF YOU MISS A DOSE?
If you miss a dose, take the next dose if still needed for pain and/or fever and the subsequent doses every 4 to 6 hours thereafter.
Do not double the dose.
HOW SHOULD YOU KEEP THIS MEDICINE?
Store at temperatures not exceeding 30oC.
Keep the product out of sight and reach of children.
For oral drops: Do not refrigerate.
SIGNS AND SYMPTOMS OF OVERDOSE
Overdosage of paracetamol usually involves four phases with the following signs and symptoms:
I.Reduced or loss of appetite, nausea, vomiting, malaise (a general feeling of being unwell), and excessive sweating;
II. Right upper abdominal pain or tenderness, liver enlargement which may be characterized by abdominal discomfort of "feeling full", elevated bilirubin and liver enzyme concentrations, prolongation of prothrombin time, and occasionally decreased urine output;
III. Reduced or loss of appetite, nausea, vomiting, and malaise recur (usually 3 to 5 days after initial symptom onset); signs of liver failure (e.g., jaundice, hypoglycemia, coagulopathy, encephalopathy), and possibly kidney failure and cardiomyopathy (disorder of the heart muscle) may develop;
IV. Recovery or progression to fatal complete liver failure
Metabolic acidosis (excessive acid in the body), encephalopathy (abnormal brain function), coma, and death may also occur. Clinical and laboratory evidence of hepatic toxicity may not be apparent 48 to 72 hours post-ingestion, and reach a maximum after 4 to 6 days.
WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN THE RECOMMENDED DOSAGE?
If you have taken more than the recommended dosage, consult a doctor or contact a poison control center right away, even if you seem well, as there is a risk of delayed serious liver damage. Quick medical attention is important for adults as well as for children even if you do not notice any signs or symptoms.
CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE
• Liver Warning: This product contains paracetamol. Severe liver damage may occur if:
o A child takes more than 5 doses in 24 hours, which is the maximum daily amount;
o An adult or child 12 years old and above takes more than 4 g of paracetamol in 24 hours;
o Taken with other medicines containing paracetamol (or acetaminophen);
• Allergy alert: Paracetamol may cause allergic reactions. Symptoms may include:
o Skin redness
o Blisters
o Skin rash or hives
o Swelling of the face, tongue or throat
o Difficulty swallowing
o Unexplained wheezing or shortness of breath
• Ask a doctor before use if patient has liver or kidney disease.
• Use with caution in patients with glucose-6-phosphate deficiency (G6PD), a hereditary condition which can lead to decreased blood cell count and possible hemolytic anemia (anemia resulting from destruction of red blood cells).
• Do not take more than the recommended dose.
• Do not use after the expiry date on the label.
Recommended Paracetamol Dose in Children
Based on 10 mg/kg body weight per dose

Age Group 100 mg/mL Syrup
(Oral Drops) 120 mg/5 mL Syrup 250 mg/5 mL Syrup
Infants:
0 to 3 months 0.3 to 0.6 mL --- ---
7 months to
under 1 year 1 mL --- ---
Children:
(1 teaspoonful)
2.5 mL
(1/2 teaspoonful)
(1/2 to 1 teaspoonful)
(1 to 11/2
teaspoonsful)
ARTWORK INFORMATION: P300000040493 IN BIOGESIC-PEDIA MELON LINE INSERT
NEW Reg. No. DR-XY46366 Pantone/s:
Old FG/SKU Code PMS No. 180598-01
Basis for
Revision:
New FG/SKU Code KG/PLG No. N/A
Existing Barcode Dimension CCR No.:
New Barcode Existing PM Code N/A
Method of Proofing Performed (Division) Global Vision Inspection (*Provide GV Report to QA) Manual Proofing
Reference Checklist (Division):
CPR/CPN/Approval of country PMS Knifeguide or PL Guide Global Vision Report
Superseded Artwork FDA Approved artwork PM code and FG code CCDS
Evaluation Checklist (QA): Global Vision Report (if applicable)
CPR/Approval of country Dimension (PMS & KG/PLG) Barcode (SAP) Color and Pantone
Photo-eye(tubes/Sachet) Approved CCR Space for LEM CCDS
Unilab logo and seal(CBG Guideline) Code bars (ISFG) Manufacturing Clause FDA Approved AW/Notification
Marketing Division: Name, Signature, Date QA-SMS-PM: Disposition:
APPROVED
N/A Maria Lourdes Genuino, MD DISAPPROVED
N/A N/A Remarks:
Noted By:
Kathyrene Visco
140 x 196 mm (±1mm) -
spread TMP-CCR-0108
EXISTING rev no.
180598
N/A
Black
Change in Brand Name, Update in
Packaging Text
Div. Pharmacist
In-house
Artist/Agency Med. Director Evaluated
Brand By:
Personnel
Div. Head/ Prod.
Dir.

WHEN SHOULD YOU CONSULT YOUR DOCTOR?

Ask a doctor before use if:
- The patient has kidney or liver problems (including alcoholic liver disease)
- The patient is taking other pain killers or any other medicines (See ***What Other Medicine or Food Should Be Avoided While Taking This Medicine?***)
- The patient has G6PD
- The cause of the patient's illness is not clear or it is accompanied by a rash, breathing difficulties, diarrhea or excessive tiredness or lethargy
- The patient is severely malnourished, anorexic, have a low body mass index (underweight), or excessively drinks alcohol

Stop use and ask a doctor if:
- If symptoms of allergic reaction occur (such as skin redness, blisters, or rash), stop use and seek medical help right away (See ***Care that Should be Taken when Taking this Medicine).***
- Symptoms worsen or do not improve
- New symptoms occur
- Any undesirable effect occurs
- Redness or swelling is present
- Fever gets worse or lasts more than 3 days
- Pain gets worse or lasts more than 5 days in children (<12 years old) and 10 days in adults
- Patient had unexpected bruising or bleeding, and had more infections than usual
- Consult a doctor promptly if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting.

ADVERSE DRUG REACTION REPORTING STATEMENT

For suspected adverse drug reaction, seek medical attention immediately and report to the FDA at www.fda.gov.ph AND Unilab at +632-8-**UNILAB**-1 (+632-8-**864522**-1) for Metro Manila or toll-free +1-800-10-UNILAB-1 for provinces, or e-mail productsafety@unilab.com.ph. By reporting undesirable effects, you can help provide more information on the safety of this medicine.

AVAILABILITY

Paracetamol (Biogesic®) 100 mg/mL Syrup Orange (DRHR-1747)
- Pediatech HDPE Plastic Bottle with Child Resistant Plastic Cap x 15 mL (Box of 1's)

Paracetamol (Biogesic®) 120 mg/5 mL Syrup Orange (DRHR-1748)
- Pediatech HDPE Plastic Bottle with Child Resistant Plastic Cap x 60 mL (Box of 1's)

Paracetamol (Biogesic®) 120 mg/5 mL Syrup Melon (DRHR-1843)
- Pediatech HDPE Plastic Bottle with Child Resistant Plastic Cap x 60 mL (Box of 1's)

Paracetamol (Biogesic®) 250 mg/5 mL Syrup Orange (DR-XY45639)
- Pediatech HDPE Plastic Bottle with Child Resistant Plastic Cap x 60 mL (Box of 1's)

Paracetamol (Biogesic®) 250 mg/5 mL Syrup Melon (DR-XY46366)
- Pediatech HDPE Plastic Bottle with Child Resistant Plastic Cap x 60 mL (Box of 1's)

Manufactured by AMHERST LABORATORIES, INC.
UNILAB Pharma Campus, Barangay Mamplasan
Biñan, Laguna, Philippines
for **UNILAB, Inc.**
No. 66 United Street, Mandaluyong City
Metro Manila, Philippines

DATE OF FIRST AUTHORIZATION

Paracetamol (Biogesic®) - Orange Flavor: November 2016
Paracetamol (Biogesic®) – Melon Flavor: June 2018

DATE OF REVISION OF PATIENT INFORMATION LEAFLET: May 2023 (CCDS 17-047)

UNILAB Consumer Care
8-UNILAB-1
8-864522-1
Provincial Toll Free: 1-800-10-8645221
Email: info@unilab.com.ph
Website: www.unilab.com.ph

UNILAB
Trusted Quality Healthcare

Reg. IPOPHIL

P300000040493

UNILAB

PARACETAMOL BIOGESIC®

100 mg/mL Syrup (Oral Drops)
120 mg/5 mL Syrup
250 mg/5 mL Syrup

ANALGESIC / ANTIPYRETIC (Anilide)

FORMULATION

Each mL syrup (oral drops) contains: Paracetamol.....100 mg
Each 5 mL syrup contains: Paracetamol.....120 mg or 250 mg

PRODUCT DESCRIPTION

Paracetamol (Biogesic®) 100 mg/mL Syrup (Oral Drops), 120 mg/5 mL Syrup, and 250 mg/5 mL Syrup (Orange Flavor)
- An orange-colored, clear syrup with an orange odor and taste

Paracetamol (Biogesic®) 120 mg/5 mL Syrup, and 250 mg/5 mL Syrup (Melon Flavor)
- An orange-colored, clear syrup with a melon odor and taste

These products use the ***TasteRite®*** technology of ***PediaTech®***. The ***TasteRite®*** technology is a unique tastemasking system developed specifically for liquid dosage forms.
This technology significantly reduces the bitterness of medicine so that children taste the flavor and not the medicine.

In addition, these products use Tooth Friendly technology ***TFTech®*** to significantly reduce or eliminate enamel loss that leads to tooth decay.

WHAT IS IN THE MEDICINE?

This product contains paracetamol, a fever reducer and pain reliever.

STRENGTH OF THE MEDICINES

See Formulations

WHAT IS THE MEDICINE USED FOR?

MEDICINE	USES
100 mg/mL Syrup (Oral Drops)	• For the relief of minor aches and pains due to the common cold, flu, sore throat, teething, headache, earache, and post-immunization reactions • For fever reduction.
120 mg/5 mL Syrup and 250 mg/5 mL Syrup	• For the relief of minor aches and pains due to the common cold, flu, headache, teething, sore throat, earache, post-immunization reactions, and toothache • For fever reduction.

HOW MUCH AND HOW OFTEN SHOULD YOU USE THIS MEDICINE?

General Recommendations:
- Infants and children should not exceed 5 doses in each 24-hour period.
- Leave at least 4 hours between doses.
- Do not use continuously for more than 5 days at a time unless directed by a doctor.
- Do not use more than the recommended dose.
- Avoid potential liver damage by giving lower doses of paracetamol in patients classified under the lower body weight group.

This medicine should be taken orally every 4 to 6 hours, as needed for pain and/or fever, or, as directed by a doctor.

Dosing in Children: 10 mg/kg body weight per dose given every 4 to 6 hours, as needed; or, you can follow the dose in the dosing table below:

140 mm

XXXX

Age Group	Recommended Paracetamol Dose in Children Based on 10 mg/kg body weight per dose		
	100 mg/mL Syrup (Oral Drops)	120 mg/5 mL Syrup	250 mg/5 mL Syrup
Infants:			
0 to 3 months	0.3 to 0.6 mL	---	---
4 to 6 months	0.6 to 1 mL	---	---
7 months to under 1 year	1 mL	---	---
Children:			
1 to 2 years	1 to 1.2 mL	5 mL (1 teaspoonful)	2.5 mL (1/2 teaspoonful)
3 to 6 years	---	5 to 10 mL (1 to 2 teaspoonsful)	2.5 to 5 mL (1/2 to 1 teaspoonful)
7 to 12 years	---	10 to 15 mL (2 to 3 teaspoonsful)	5 to 7.5 mL (1 to 11/2 teaspoonsful)

WHEN SHOULD YOU NOT TAKE THIS MEDICINE?
- If you are allergic to paracetamol or any ingredient of the product
- If you are taking other medicines that contain paracetamol

UNDESIRABLE EFFECTS

Paracetamol, when taken within the recommended dose and duration of treatment, has low incidence of side effects.

Blood and lymphatic system disorders: Anemia (decreased number of healthy red blood cells), changes in the number of white blood cells and platelets such as agranulocytosis (decrease in the number of granulocytes), leukopenia (decrease in the number of leukocytes), methemoglobinemia (presence of methemoglobin in the blood), neutropenia (decrease in the number of neutrophils), pancytopenia (deficiency of all cellular elements in the blood), thrombocytopenia (decrease in the number of platelets); thrombocytopenic purpura (small purple-red marks in the skin or other tissue caused by low number of platelets)
Immune system disorders: Allergic reactions (some are life-threatening) which may cause difficulty swallowing, unexplained wheezing, shortness of breath or difficulty in breathing, skin rash, hives, or angioedema (swelling of face, tongue, lips, or throat); proneness to infections, swelling around the eyes
Metabolism and nutrition disorders: Hyperchloremia (increased chlorine ion in the blood), hyperuricemia (increased uric acid in the blood), loss of appetite, metabolic acidosis (too much acid in the body), sudden weight loss
Nervous system disorders: Headache
Vascular disorders: Unusual bleeding or bruising
Respiratory, thoracic, and mediastinal disorders: Respiratory distress (noisy, congested breathing), runny nose, laryngeal edema (swelling of the voice box)
Gastrointestinal disorders: Mouth ulcers, nausea, sore throat, vomiting
Skin and subcutaneous tissue disorder: Fixed eruption (an allergic reaction described as well-defined, round or oval patches of redness and swelling in skin that recurs in the same site each time a particular drug is taken), itchy rash, severe skin reactions (i.e., Stevens-Johnson syndrome, toxic epidermal necrolysis, and acute generalized exanthematous pustulosis) which may include symptoms such as skin redness, blisters, rash, itching, and peeling of the upper surface of the skin
Hepatobiliary disorders: Chronic hepatic necrosis (death of liver tissues) in prolonged use, liver dysfunction, liver injury, yellowing of eyes and skin
Renal and urinary disorders: Kidney toxicity, papillary necrosis in prolonged use
General disorders and administration site conditions: Fatigue or unusual tiredness
Investigations: Decreased serum bicarbonate, serum calcium, serum sodium; increases in serum alkaline phosphatase, serum glucose, and transaminase levels

WHAT OTHER MEDICINE OR FOOD SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?
- Do not use with any other medicine containing paracetamol (prescription or nonprescription). If you are not sure whether a medicine contains paracetamol, ask a doctor.
- Prolonged use of paracetamol with other painkillers [e.g., drugs called nonsteroidal anti-inflammatory drugs (NSAIDs), aspirin] is not recommended since it may increase the risk of toxic effects to the kidneys and eventually of kidney failure.
- Chronic intake of alcohol may increase the risk of paracetamol-induced liver toxicity. Acute intake of alcohol may diminish the ability of the body to metabolize large doses of paracetamol. It is advised to consult a doctor before intake of paracetamol.
- The absorption of paracetamol may be increased by metoclopramide or domperidone. Cholestyramine, narcotic pain relievers, and some antidepressants that slow down gastric emptying reduce the absorption of paracetamol. If you are taking these medicines, talk to your doctor first before taking paracetamol.
- Medicines which stimulate the enzymes responsible for the metabolic activation of paracetamol such as medicines for convulsion (e.g., phenobarbital), depression (e.g., tricyclic antidepressants), and oral contraceptives may reduce the efficacy of paracetamol and increase susceptibility to the harmful effects to the liver.
- Phenothiazines (medicines for serious mental and emotional disorder) may increase the risk of severe hypothermia (abnormally low body temperature)
- Paracetamol increases exposure to chloramphenicol. This interaction should be considered when both medicines are given together in malnourished patients.
- Probenecid may affect the elimination of paracetamol.
- There may be an increased risk of bleeding when using paracetamol and warfarin (a blood thinning medicine) at the same time. Paracetamol, especially its prolonged regular use, increases the blood thinning effect of warfarin and other coumarins. If you are taking warfarin, ask the doctor first before taking paracetamol.

Tell your doctor if you are taking other medicines (especially other paracetamol-containing medicines), including vitamins, supplements, and herbal medicines.

WHAT SHOULD YOU DO IF YOU MISS A DOSE?

If you miss a dose, take the next dose if still needed for pain and/or fever and the subsequent doses every 4 to 6 hours thereafter.

Do not double the dose.

HOW SHOULD YOU KEEP THIS MEDICINE?

Store at temperatures not exceeding 30°C.
Keep the product out of sight and reach of children.
For oral drops: **Do not refrigerate.**

SIGNS AND SYMPTOMS OF OVERDOSE

Overdosage of paracetamol usually involves four phases with the following signs and symptoms:
I.Reduced or loss of appetite, nausea, vomiting, malaise (a general feeling of being unwell), and excessive sweating;
II. Right upper abdominal pain or tenderness, liver enlargement which may be characterized by abdominal discomfort of "feeling full", elevated bilirubin and liver enzyme concentrations, prolongation of prothrombin time, and occasionally decreased urine output;
III. Reduced or loss of appetite, nausea, vomiting, and malaise recur (usually 3 to 5 days after initial symptom onset); signs of liver failure (e.g., jaundice, hypoglycemia, coagulopathy, encephalopathy), and possibly kidney failure and cardiomyopathy (disorder of the heart muscle) may develop;
IV. Recovery or progression to fatal complete liver failure

Metabolic acidosis (excessive acid in the body), encephalopathy (abnormal brain function), coma, and death may also occur. Clinical and laboratory evidence of hepatic toxicity may not be apparent 48 to 72 hours post-ingestion, and reach a maximum after 4 to 6 days.

WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN THE RECOMMENDED DOSAGE?

If you have taken more than the recommended dosage, consult a doctor or contact a poison control center right away, even if you seem well, as there is a risk of delayed serious liver damage. Quick medical attention is important for adults as well as for children even if you do not notice any signs or symptoms.

CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE

- **Liver Warning:** This product contains paracetamol. Severe liver damage may occur if:
  - o A child takes more than 5 doses in 24 hours, which is the maximum daily amount;
  - o An adult or child 12 years old and above takes more than 4 g of paracetamol in 24 hours;
  - o Taken with other medicines containing paracetamol (or acetaminophen);
- **Allergy alert:** Paracetamol may cause allergic reactions. Symptoms may include:
  - o Skin redness
  - o Blisters
  - o Skin rash or hives
  - o Swelling of the face, tongue or throat
  - o Difficulty swallowing
  - o Unexplained wheezing or shortness of breath
- Ask a doctor before use if patient has liver or kidney disease.
- Use with caution in patients with glucose-6-phosphate deficiency (G6PD), a hereditary condition which can lead to decreased blood cell count and possible hemolytic anemia (anemia resulting from destruction of red blood cells).
- Do not take more than the recommended dose.
- Do not use after the expiry date on the label.

APPROVED
By Kathyrene Y. Visco, RPh at 3:25 pm, Jun 13, 2023

APPROVED
By Maria Lourdes G. Genuino,MD at 4:10 pm, Jun 13, 2023

196 mm

ARTWORK INFORMATION: P300000040493 IN BIOGESIC-PEDIA MELON LINE INSERT

■ **NEW** ☐ **EXISTING** rev no. ____

Reg. No. DR-XY46366
Pantone/s: Black
Old FG/SKU Code 180598
PMS No. 180598-01
Basis for Revision: Change in Brand Name, Update in Packaging Text
New FG/SKU Code N/A
KG/PLG No. N/A
Existing Barcode N/A
Dimension 140 x 196 mm (±1mm) - spread
CCR No.: TMP-CCR-0108
New Barcode N/A
Existing PM Code N/A

Method of Proofing Performed (Division) ☐ Global Vision Inspection (*Provide GV Report to QA) ■ Manual Proofing

Reference Checklist (Division):
■ CPR/CPN/Approval of country ■ PMS ☐ Knifeguide or PL Guide ☐ Global Vision Report
☐ Superseded Artwork ☐ FDA Approved artwork ■ PM code and FG code ■ CCDS

Evaluation Checklist (QA):
☑ CPR/Approval of country ☑ PM code and FG code ☐ Global Vision Report (if applicable)
☐ Photo-eye(tubes/Sachet) ☑ Dimension (PMS & KG/PLG) ☐ Barcode (SAP) ☑ Color and Pantone
☑ Unilab logo and seal(CBG Guideline) ☐ Approved CCR ☐ Space for LEM ☐ CCDS
☑ Code bars (ISFG) ☑ Manufacturing Clause ☐ FDA Approved AW/Notification

Marketing Division: *Name, Signature, Date*
In-house Artist/Agency N/A
Brand Personnel N/A
Med. Director Maria Lourdes Genuino, MD
Div. Head/ Prod. Dir. N/A
Div. Pharmacist Kathyrene Visco

QA-SMS-PM:
Evaluated By:
Noted By:

Disposition:
☐ APPROVED
☐ DISAPPROVED
Remarks:

U-F-SQM-044/01

APPROVED
UL-CQA&C

APPROVED
By Jonna Marie Sy at 6:40 pm, Jun 17, 2023

APPROVED
By Amalia B. Magadia at 9:54 am, Jun 19, 2023

Biogesic (Paracetamol)

250 mg/5mL Syrup (Melon Flavor) Syrup | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY46366

Biogesic: Official FDA Philippines registration record

Brand name	Biogesic
Generic name	Paracetamol
Dosage strength and form	250 mg/5mL Syrup (Melon Flavor) Syrup
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Analgesic/ Antipyretic (Anilide)
Registration number	DR-XY46366
Manufacturer	Amherst Laboratories, Inc.

Biogesic: OTC forms registered under the Biogesic brand (capsule availability)

OTC capsule form registered under the Biogesic brand: NO. 7 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Biogesic	100 mg / mL Suspension (Oral Drops) [Orange Sugar Free]	Household Remedy (HR)	DRHR-1167
Biogesic	120 mg/5 mL Syrup (Melon Flavor)	Household Remedy (HR)	DRHR-1843
Biogesic	120 mg/5 mL Syrup (Orange Flavor)	Household Remedy (HR)	DRHR-1748
Biogesic	250 mg / 5 mL Syrup (Orange Flavor)	Household Remedy (HR)	DR-XY45639
Biogesic	250 mg/5mL Syrup (Melon Flavor) Syrup	Over-the-Counter (OTC) Drug	DR-XY46366
Biogesic 325	325 mg Tablet	Over-the-Counter (OTC) Drug	DRP-8763
Biogesic	500 mg Tablet	Over-the-Counter (OTC) Drug	DR-153

Biogesic: Official leaflet (Package Insert / Patient Information Leaflet)

An official leaflet for this registration is published on the FDA Verification Portal: PIL_01 (https://drive.google.com/file/d/1SeQvM-ElkZQqgWRE1CvGOXVCOiEQeguM/view?usp=drivesdk). Its text (OCR) and the original leaflet pages follow.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Biogesic"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Biogesic: Official leaflet text (OCR transcription)

Machine transcription of the official FDA leaflet for Reg. No. DR-XY46366. OCR may contain errors; the original leaflet pages are attached after this section.

Biogesic leaflet PIL_01

FORMULATION
Each mL syrup (oral drops) contains: Paracetamol…….....…….100 mg
Each 5 mL syrup contains: Paracetamol…….....….120 mg or 250 mg
PRODUCT DESCRIPTION
Paracetamol (Biogesic®) 100 mg/mL Syrup (Oral Drops), 120 mg/5 mL Syrup, and 250 mg/5 mL Syrup (Orange Flavor)
• An orange-colored, clear syrup with an orange odor and taste
Paracetamol (Biogesic®) 120 mg/5 mL Syrup, and 250 mg/5 mL Syrup (Melon Flavor)
• An orange-colored, clear syrup with a melon odor and taste
These products use the TasteRite® technology of PediaTech®. The TasteRite® technology is a unique tastemasking system developed specifically for liquid dosage forms.
This technology significantly reduces the bitterness of medicine so that children taste the flavor and not the medicine.
In addition, these products use Tooth Friendly technology TFTech® to significantly reduce or eliminate enamel loss that leads to tooth decay.
WHAT IS IN THE MEDICINE?
This product contains paracetamol, a fever reducer and pain reliever.
STRENGTH OF THE MEDICINES
See Formulations
WHAT IS THE MEDICINE USED FOR?
HOW MUCH AND HOW OFTEN SHOULD YOU USE THIS MEDICINE?
General Recommendations:
• Infants and children should not exceed 5 doses in each 24-hour period.
• Leave at least 4 hours between doses.
• Do not use continuously for more than 5 days at a time unless directed by a doctor.
• Do not use more than the recommended dose.
• Avoid potential liver damage by giving lower doses of paracetamol in patients classified under the lower body weight group.
This medicine should be taken orally every 4 to 6 hours, as needed for pain and/or fever, or, as directed by a doctor.
Dosing in Children: 10 mg/kg body weight per dose given every 4 to 6 hours, as needed; or, you can follow the dose in the dosing table below:
WHEN SHOULD YOU NOT TAKE THIS MEDICINE?
• If you are allergic to paracetamol or any ingredient of the product
• If you are taking other medicines that contain paracetamol
UNDESIRABLE EFFECTS
Paracetamol, when taken within the recommended dose and duration of treatment, has low incidence of side effects.
Blood and lymphatic system disorders: Anemia (decreased number of healthy red blood cells), changes in the number of white blood cells and platelets such as agranulocytosis (decrease in the number of granulocytes), leukopenia (decrease in the number of leukocytes), methemoglobinemia (presence of methemoglobin in the blood), neutropenia (decrease in the number of neutrophils), pancytopenia (deficiency of all cellular elements in the blood), thrombocytopenia (decrease in the number of platelets); thrombocytopenic purpura (small purple-red marks in the skin or other tissue caused by low number of platelets)
Immune system disorders: Allergic reactions (some are life-threatening) which may cause difficulty swallowing, unexplained wheezing, shortness of breath or difficulty in breathing, skin rash, hives, or angioedema (swelling of face, tongue, lips, or throat); proneness to infections, swelling around the eyes
Metabolism and nutrition disorders: Hyperchloremia (increased chlorine ion in the blood), hyperuricemia (increased uric acid in the blood), loss of appetite, metabolic acidosis (too much acid in the body), sudden weight loss
Nervous system disorders: Headache
Vascular disorders: Unusual bleeding or bruising
Respiratory, thoracic, and mediastinal disorders: Respiratory distress (noisy, congested breathing), runny nose, laryngeal edema (swelling of the voice box)
Gastrointestinal disorders: Mouth ulcers, nausea, sore throat, vomiting
Skin and subcutaneous tissue disorder: Fixed eruption (an allergic reaction described as well-defined, round or oval patches of redness and swelling in skin that recurs in the same site each time a particular drug is taken), itchy rash, severe skin reactions (i.e., Stevens-Johnson syndrome, toxic epidermal necrolysis, and acute generalized exanthematous pustulosis) which may include symptoms such as skin redness, blisters, rash, itching, and peeling of the upper surface of the skin
Hepatobiliary disorders: Chronic hepatic necrosis (death of liver tissues) in prolonged use, liver dysfunction, liver injury, yellowing of eyes and skin
Renal and urinary disorders: Kidney toxicity, papillary necrosis in prolonged use
General disorders and administration site conditions: Fatigue or unusual tiredness
Investigations: Decreased serum bicarbonate, serum calcium, serum sodium; increases in serum alkaline

phosphatase, serum glucose, and transaminase levels
Do not use with any other medicine containing paracetamol (prescription or nonprescription). If you are not surewhether a medicine contains paracetamol, ask a doctor.
Prolonged use of paracetamol with other painkillers [e.g., drugs called nonsteroidal anti-inflammatory drugs (NSAIDs), aspirin] is not recommended since it may increase the risk of toxic effects to the kidneys and eventually of kidney failure.
Chronic intake of alcohol may increase the risk of paracetamol-induced liver toxicity. Acute intake of alcohol may diminish the ability of the body to metabolize large doses of paracetamol. It is advised to consult a doctor before intake of paracetamol.
WHAT OTHER MEDICINE OR FOOD SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?
•
Reg. IPOPHIL P300000040493
8-864522-1 8-UNILAB-1
100 mg/mL Syrup (Oral Drops)
120 mg/5 mL Syrup
250 mg/5 mL Syrup
ANALGESIC / ANTIPYRETIC (Anilide)
The absorption of paracetamol may be increased by metoclopramide or domperidone. Cholestyramine, narcotic pain relievers, and some antidepressants that slow down gastric emptying reduce the absorption of paracetamol. If you are taking these medicines, talk to your doctor first before taking paracetamol.
Medicines which stimulate the enzymes responsible for the metabolic activation of paracetamol such as medicines for convulsion (e.g., phenobarbital), depression (e.g., tricyclic antidepressants), and oral contraceptives may reduce the efficacy of paracetamol and increase susceptibility to the harmful effects to the liver.
Phenothiazines (medicines for serious mental and emotional disorder) may increase the risk of severe hypothermia (abnormally low body temperature)
Paracetamol increases exposure to chloramphenicol. This interaction should be considered when both medicines are given together in malnourished patients.
Probenecid may affect the elimination of paracetamol.
There may be an increased risk of bleeding when using paracetamol and warfarin (a blood thinning medicine) at the same time. Paracetamol, especially its prolonged regular use, increases the blood thinning effect of warfarin and other coumarins. If you are taking warfarin, ask the doctor first before taking paracetamol.
MEDICINE USES
100 mg/mL Syrup (Oral Drops)
•
• For fever reduction.
120 mg/5 mL Syrup and
250 mg/5 mL Syrup
• For the relief of minor aches and pains due to the common cold, flu, headache, teething, sore throat, earache, post-immunization reactions, and toothache
WHEN SHOULD YOU CONSULT YOUR DOCTOR?
Ask a doctor before use if:
• The patient has kidney or liver problems (including alcoholic liver disease)
• The patient is taking other pain killers or any other medicines (See What Other Medicine or Food Should Be Avoided While Taking This Medicine?)
• The patient has G6PD
• The cause of the patient’s illness is not clear or it is accompanied by a rash, breathing difficulties, diarrhea or excessive tiredness or lethargy
• The patient is severely malnourished, anorexic, have a low body mass index (underweight), or excessively drinks alcohol
Stop use and ask a doctor if:
• If symptoms of allergic reaction occur (such as skin redness, blisters, or rash), stop use and seek medical help right away (See Care that Should be Taken when Taking this Medicine).
• Symptoms worsen or do not improve
• New symptoms occur
• Any undesirable effect occurs
• Redness or swelling is present
• Fever gets worse or lasts more than 3 days
• Pain gets worse or lasts more than 5 days in children (<12 years old) and 10 days in adults
• Patient had unexpected bruising or bleeding, and had more infections than usual
• Consult a doctor promptly if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting.
ADVERSE DRUG REACTION REPORTING STATEMENT
For suspected adverse drug reaction, seek medical attention immediately and report to the FDA at www.fda.gov.ph AND Unilab at+632-8-UNILAB-1 (+632-8-864522-1) for Metro Manila or toll-free +1-800-10-UNILAB-1 for provinces, or e-mail productsafety@unilab.com.ph. By reporting undesirable effects, you can help provide more information on the safety of this medicine.

AVAILABILITY
Paracetamol (Biogesic®) 100 mg/mL Syrup Orange (DRHR-1747)
- Pediatech HDPE Plastic Botttle with Child Resistant Plastic Cap x 15 mL (Box of 1's)
Paracetamol (Biogesic®) 120 mg/5 mL Syrup Orange (DRHR-1748)
- Pediatech HDPE Plastic Botttle with Child Resistant Plastic Cap x 60 mL (Box of 1's)
Paracetamol (Biogesic®) 120 mg/5 mL Syrup Melon (DRHR-1843)
- Pediatech HDPE Plastic Bottle with Child Resistant Plastic Cap x 60 mL (Box of 1's)
Paracetamol (Biogesic®) 250 mg/5 mL Syrup Orange (DR-XY45639)
Paracetamol (Biogesic®) 250 mg/5 mL Syrup Melon (DR-XY46366)
Manufactured by AMHERST LABORATORIES, INC.
UNILAB Pharma Campus, Barangay Mamplasan
Biñan, Laguna, Philippines
for UNILAB, Inc.
No. 66 United Street, Mandaluyong City
Metro Manila, Philippines
DATE OF FIRST AUTHORIZATION
Paracetamol (Biogesic®) - Orange Flavor: November 2016
Paracetamol (Biogesic®) – Melon Flavor: June 2018
DATE OF REVISION OF PATIENT INFORMATION LEAFLET: May 2023 (CCDS 17-047)
For the relief of minor aches and pains due to the common cold, flu,
sore throat, teething, headache, earache, and post-immunization
reactions
Tell your doctor if you are taking other medicines (especially other paracetamol-containing medicines), including vitamins, supplements, and herbal medicines.
WHAT SHOULD YOU DO IF YOU MISS A DOSE?
If you miss a dose, take the next dose if still needed for pain and/or fever and the subsequent doses every 4 to 6 hours thereafter.
Do not double the dose.
HOW SHOULD YOU KEEP THIS MEDICINE?
Store at temperatures not exceeding 30oC.
Keep the product out of sight and reach of children.
For oral drops: Do not refrigerate.
SIGNS AND SYMPTOMS OF OVERDOSE
Overdosage of paracetamol usually involves four phases with the following signs and symptoms:
I.Reduced or loss of appetite, nausea, vomiting, malaise (a general feeling of being unwell), and excessive sweating;
II. Right upper abdominal pain or tenderness, liver enlargement which may be characterized by abdominal discomfort of "feeling full", elevated bilirubin and liver enzyme concentrations, prolongation of prothrombin time, and occasionally decreased urine output;
III. Reduced or loss of appetite, nausea, vomiting, and malaise recur (usually 3 to 5 days after initial symptom onset); signs of liver failure (e.g., jaundice, hypoglycemia, coagulopathy, encephalopathy), and possibly kidney failure and cardiomyopathy (disorder of the heart muscle) may develop;
IV. Recovery or progression to fatal complete liver failure
Metabolic acidosis (excessive acid in the body), encephalopathy (abnormal brain function), coma, and death may also occur. Clinical and laboratory evidence of hepatic toxicity may not be apparent 48 to 72 hours post-ingestion, and reach a maximum after 4 to 6 days.
WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN THE RECOMMENDED DOSAGE?
If you have taken more than the recommended dosage, consult a doctor or contact a poison control center right away, even if you seem well, as there is a risk of delayed serious liver damage. Quick medical attention is important for adults as well as for children even if you do not notice any signs or symptoms.
CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE
• Liver Warning: This product contains paracetamol. Severe liver damage may occur if:
o A child takes more than 5 doses in 24 hours, which is the maximum daily amount;
o An adult or child 12 years old and above takes more than 4 g of paracetamol in 24 hours;
o Taken with other medicines containing paracetamol (or acetaminophen);
• Allergy alert: Paracetamol may cause allergic reactions. Symptoms may include:
o Skin redness
o Blisters
o Skin rash or hives
o Swelling of the face, tongue or throat
o Difficulty swallowing
o Unexplained wheezing or shortness of breath
• Ask a doctor before use if patient has liver or kidney disease.
• Use with caution in patients with glucose-6-phosphate deficiency (G6PD), a hereditary condition which can lead to decreased blood cell count and possible hemolytic anemia (anemia resulting from destruction of red blood cells).
• Do not take more than the recommended dose.
• Do not use after the expiry date on the label.
Recommended Paracetamol Dose in Children

Based on 10 mg/kg body weight per dose
Age Group 100 mg/mL Syrup
(Oral Drops) 120 mg/5 mL Syrup 250 mg/5 mL Syrup
Infants:
0 to 3 months 0.3 to 0.6 mL --- ---
7 months to
under 1 year 1 mL --- ---
Children:
(1 teaspoonful)
2.5 mL
(1/2 teaspoonful)
(1/2 to 1 teaspoonful)
(1 to 11/2
teaspoonsful)
ARTWORK INFORMATION: P300000040493 IN BIOGESIC-PEDIA MELON LINE INSERT
NEW Reg. No. DR-XY46366 Pantone/s:
Old FG/SKU Code PMS No. 180598-01
Basis for
Revision:
New FG/SKU Code KG/PLG No. N/A
Existing Barcode Dimension CCR No.:
New Barcode Existing PM Code N/A
Method of Proofing Performed (Division) Global Vision Inspection (*Provide GV Report to QA) Manual Proofing
Reference Checklist (Division):
CPR/CPN/Approval of country PMS Knifeguide or PL Guide Global Vision Report
Superseded Artwork FDA Approved artwork PM code and FG code CCDS
Evaluation Checklist (QA): Global Vision Report (if applicable)
CPR/Approval of country Dimension (PMS & KG/PLG) Barcode (SAP) Color and Pantone
Photo-eye(tubes/Sachet) Approved CCR Space for LEM CCDS
Unilab logo and seal(CBG Guideline) Code bars (ISFG) Manufacturing Clause FDA Approved AW/Notification
Marketing Division: Name, Signature, Date QA-SMS-PM: Disposition:
APPROVED
N/A Maria Lourdes Genuino, MD DISAPPROVED
N/A N/A Remarks:
Noted By:
Kathyrene Visco
140 x 196 mm (±1mm) -
spread TMP-CCR-0108
EXISTING rev no.
180598
N/A
Black
Change in Brand Name, Update in
Packaging Text
Div. Pharmacist
In-house
Artist/Agency Med. Director Evaluated
Brand By:
Personnel
Div. Head/ Prod.
Dir.

WHEN SHOULD YOU CONSULT YOUR DOCTOR?
Ask a doctor before use if:
- The patient has kidney or liver problems (including alcoholic liver disease)
- The patient is taking other pain killers or any other medicines (See ***What Other Medicine or Food Should Be Avoided While Taking This Medicine?***)
- The patient has G6PD
- The cause of the patient's illness is not clear or it is accompanied by a rash, breathing difficulties, diarrhea or excessive tiredness or lethargy
- The patient is severely malnourished, anorexic, have a low body mass index (underweight), or excessively drinks alcohol

Stop use and ask a doctor if:
- If symptoms of allergic reaction occur (such as skin redness, blisters, or rash), stop use and seek medical help right away (See ***Care that Should be Taken when Taking this Medicine).***
- Symptoms worsen or do not improve
- New symptoms occur
- Any undesirable effect occurs
- Redness or swelling is present
- Fever gets worse or lasts more than 3 days
- Pain gets worse or lasts more than 5 days in children (<12 years old) and 10 days in adults
- Patient had unexpected bruising or bleeding, and had more infections than usual
- Consult a doctor promptly if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting.

ADVERSE DRUG REACTION REPORTING STATEMENT
For suspected adverse drug reaction, seek medical attention immediately and report to the FDA at www.fda.gov.ph AND Unilab at+632-8-**UNILAB**-1 (+632-8-**864522**-1) for Metro Manila or toll-free +1-800-10-UNILAB-1 for provinces, or e-mail productsafety@unilab.com.ph. By reporting undesirable effects, you can help provide more information on the safety of this medicine.

AVAILABILITY
Paracetamol (Biogesic®) 100 mg/mL Syrup Orange (DRHR-1747)
- Pediatech HDPE Plastic Botttle with Child Resistant Plastic Cap x 15 mL (Box of 1's)

Paracetamol (Biogesic®) 120 mg/5 mL Syrup Orange (DRHR-1748)
- Pediatech HDPE Plastic Botttle with Child Resistant Plastic Cap x 60 mL (Box of 1's)

Paracetamol (Biogesic®) 120 mg/5 mL Syrup Melon (DRHR-1843)
- Pediatech HDPE Plastic Bottle with Child Resistant Plastic Cap x 60 mL (Box of 1's)

Paracetamol (Biogesic®) 250 mg/5 mL Syrup Orange (DR-XY45639)
- Pediatech HDPE Plastic Botttle with Child Resistant Plastic Cap x 60 mL (Box of 1's)

Paracetamol (Biogesic®) 250 mg/5 mL Syrup Melon (DR-XY46366)
- Pediatech HDPE Plastic Bottle with Child Resistant Plastic Cap x 60 mL (Box of 1's)

Manufactured by AMHERST LABORATORIES, INC.
UNILAB Pharma Campus, Barangay Mamplasan
Biñan, Laguna, Philippines
for **UNILAB, Inc.**
No. 66 United Street, Mandaluyong City
Metro Manila, Philippines

DATE OF FIRST AUTHORIZATION
Paracetamol (Biogesic®) - Orange Flavor: November 2016
Paracetamol (Biogesic®) – Melon Flavor: June 2018

DATE OF REVISION OF PATIENT INFORMATION LEAFLET: May 2023 (CCDS 17-047)

UNILAB Consumer Care
8-UNILAB-1
8-864522-1
Provincial Toll Free: 1-800-10-8645221
Email: info@unilab.com.ph
Website: www.unilab.com.ph

UNILAB
Trusted Quality Healthcare

Reg. IPOPHIL

P300000040493

UNILAB

PARACETAMOL BIOGESIC®

100 mg/mL Syrup (Oral Drops)
120 mg/5 mL Syrup
250 mg/5 mL Syrup

ANALGESIC / ANTIPYRETIC (Anilide)

FORMULATION
Each mL syrup (oral drops) contains: Paracetamol.....100 mg
Each 5 mL syrup contains: Paracetamol.....120 mg or 250 mg

PRODUCT DESCRIPTION
Paracetamol (Biogesic®) 100 mg/mL Syrup (Oral Drops), 120 mg/5 mL Syrup, and 250 mg/5 mL Syrup (Orange Flavor)
- An orange-colored, clear syrup with an orange odor and taste

Paracetamol (Biogesic®) 120 mg/5 mL Syrup, and 250 mg/5 mL Syrup (Melon Flavor)
- An orange-colored, clear syrup with a melon odor and taste

These products use the ***TasteRite®*** technology of ***PediaTech®***. The ***TasteRite®*** technology is a unique tastemasking system developed specifically for liquid dosage forms.
This technology significantly reduces the bitterness of medicine so that children taste the flavor and not the medicine.

In addition, these products use Tooth Friendly technology ***TFTech®*** to significantly reduce or eliminate enamel loss that leads to tooth decay.

WHAT IS IN THE MEDICINE?
This product contains paracetamol, a fever reducer and pain reliever.

STRENGTH OF THE MEDICINES
See Formulations

WHAT IS THE MEDICINE USED FOR?

MEDICINE	USES
100 mg/mL Syrup (Oral Drops)	• For the relief of minor aches and pains due to the common cold, flu, sore throat, teething, headache, earache, and post-immunization reactions • For fever reduction.
120 mg/5 mL Syrup and 250 mg/5 mL Syrup	• For the relief of minor aches and pains due to the common cold, flu, headache, teething, sore throat, earache, post-immunization reactions, and toothache • For fever reduction.

HOW MUCH AND HOW OFTEN SHOULD YOU USE THIS MEDICINE?

General Recommendations:
- Infants and children should not exceed 5 doses in each 24-hour period.
- Leave at least 4 hours between doses.
- Do not use continuously for more than 5 days at a time unless directed by a doctor.
- Do not use more than the recommended dose.
- Avoid potential liver damage by giving lower doses of paracetamol in patients classified under the lower body weight group.

This medicine should be taken orally every 4 to 6 hours, as needed for pain and/or fever, or, as directed by a doctor.

Dosing in Children: 10 mg/kg body weight per dose given every 4 to 6 hours, as needed; or, you can follow the dose in the dosing table below:

Age Group	Recommended Paracetamol Dose in Children Based on 10 mg/kg body weight per dose		
	100 mg/mL Syrup (Oral Drops)	**120 mg/5 mL Syrup**	**250 mg/5 mL Syrup**
Infants:			
0 to 3 months	0.3 to 0.6 mL	---	---
4 to 6 months	0.6 to 1 mL	---	---
7 months to under 1 year	1 mL	---	---
Children:			
1 to 2 years	1 to 1.2 mL	5 mL (1 teaspoonful)	2.5 mL (1/2 teaspoonful)
3 to 6 years	---	5 to 10 mL (1 to 2 teaspoonsful)	2.5 to 5 mL (1/2 to 1 teaspoonful)
7 to 12 years	---	10 to 15 mL (2 to 3 teaspoonsful)	5 to 7.5 mL (1 to 11/2 teaspoonsful)

WHEN SHOULD YOU NOT TAKE THIS MEDICINE?
- If you are allergic to paracetamol or any ingredient of the product
- If you are taking other medicines that contain paracetamol

UNDESIRABLE EFFECTS
Paracetamol, when taken within the recommended dose and duration of treatment, has low incidence of side effects.

Blood and lymphatic system disorders: Anemia (decreased number of healthy red blood cells), changes in the number of white blood cells and platelets such as agranulocytosis (decrease in the number of granulocytes), leukopenia (decrease in the number of leukocytes), methemoglobinemia (presence of methemoglobin in the blood), neutropenia (decrease in the number of neutrophils), pancytopenia (deficiency of all cellular elements in the blood), thrombocytopenia (decrease in the number of platelets); thrombocytopenic purpura (small purple-red marks in the skin or other tissue caused by low number of platelets)
Immune system disorders: Allergic reactions (some are life-threatening) which may cause difficulty swallowing, unexplained wheezing, shortness of breath or difficulty in breathing, skin rash, hives, or angioedema (swelling of face, tongue, lips, or throat); proneness to infections, swelling around the eyes
Metabolism and nutrition disorders: Hyperchloremia (increased chlorine ion in the blood), hyperuricemia (increased uric acid in the blood), loss of appetite, metabolic acidosis (too much acid in the body), sudden weight loss
Nervous system disorders: Headache
Vascular disorders: Unusual bleeding or bruising
Respiratory, thoracic, and mediastinal disorders: Respiratory distress (noisy, congested breathing), runny nose, laryngeal edema (swelling of the voice box)
Gastrointestinal disorders: Mouth ulcers, nausea, sore throat, vomiting
Skin and subcutaneous tissue disorder: Fixed eruption (an allergic reaction described as well-defined, round or oval patches of redness and swelling in skin that recurs in the same site each time a particular drug is taken), itchy rash, severe skin reactions (i.e., Stevens-Johnson syndrome, toxic epidermal necrolysis, and acute generalized exanthematous pustulosis) which may include symptoms such as skin redness, blisters, rash, itching, and peeling of the upper surface of the skin
Hepatobiliary disorders: Chronic hepatic necrosis (death of liver tissues) in prolonged use, liver dysfunction, liver injury, yellowing of eyes and skin
Renal and urinary disorders: Kidney toxicity, papillary necrosis in prolonged use
General disorders and administration site conditions: Fatigue or unusual tiredness
Investigations: Decreased serum bicarbonate, serum calcium, serum sodium; increases in serum alkaline phosphatase, serum glucose, and transaminase levels

WHAT OTHER MEDICINE OR FOOD SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?
- Do not use with any other medicine containing paracetamol (prescription or nonprescription). If you are not surewhether a medicine contains paracetamol, ask a doctor.
- Prolonged use of paracetamol with other painkillers [e.g., drugs called nonsteroidal anti-inflammatory drugs (NSAIDs), aspirin] is not recommended since it may increase the risk of toxic effects to the kidneys and eventually of kidney failure.
- Chronic intake of alcohol may increase the risk of paracetamol-induced liver toxicity. Acute intake of alcohol may diminish the ability of the body to metabolize large doses of paracetamol. It is advised to consult a doctor before intake of paracetamol.
- The absorption of paracetamol may be increased by metoclopramide or domperidone. Cholestyramine, narcotic pain relievers, and some antidepressants that slow down gastric emptying reduce the absorption of paracetamol. If you are taking these medicines, talk to your doctor first before taking paracetamol.
- Medicines which stimulate the enzymes responsible for the metabolic activation of paracetamol such as medicines for convulsion (e.g., phenobarbital), depression (e.g., tricyclic antidepressants), and oral contraceptives may reduce the efficacy of paracetamol and increase susceptibility to the harmful effects to the liver.
- Phenothiazines (medicines for serious mental and emotional disorder) may increase the risk of severe hypothermia (abnormally low body temperature)
- Paracetamol increases exposure to chloramphenicol. This interaction should be considered when both medicines are given together in malnourished patients.
- Probenecid may affect the elimination of paracetamol.
- There may be an increased risk of bleeding when using paracetamol and warfarin (a blood thinning medicine) at the same time. Paracetamol, especially its prolonged regular use, increases the blood thinning effect of warfarin and other coumarins. If you are taking warfarin, ask the doctor first before taking paracetamol.

Tell your doctor if you are taking other medicines (especially other paracetamol-containing medicines), including vitamins, supplements, and herbal medicines.

WHAT SHOULD YOU DO IF YOU MISS A DOSE?
If you miss a dose, take the next dose if still needed for pain and/or fever and the subsequent doses every 4 to 6 hours thereafter.

Do not double the dose.

HOW SHOULD YOU KEEP THIS MEDICINE?
Store at temperatures not exceeding 30°C.
Keep the product out of sight and reach of children.
For oral drops: **Do not refrigerate.**

SIGNS AND SYMPTOMS OF OVERDOSE
Overdosage of paracetamol usually involves four phases with the following signs and symptoms:
I.Reduced or loss of appetite, nausea, vomiting, malaise (a general feeling of being unwell), and excessive sweating;
II. Right upper abdominal pain or tenderness, liver enlargement which may be characterized by abdominal discomfort of "feeling full", elevated bilirubin and liver enzyme concentrations, prolongation of prothrombin time, and occasionally decreased urine output;
III. Reduced or loss of appetite, nausea, vomiting, and malaise recur (usually 3 to 5 days after initial symptom onset); signs of liver failure (e.g., jaundice, hypoglycemia, coagulopathy, encephalopathy), and possibly kidney failure and cardiomyopathy (disorder of the heart muscle) may develop;
IV. Recovery or progression to fatal complete liver failure

Metabolic acidosis (excessive acid in the body), encephalopathy (abnormal brain function), coma, and death may also occur. Clinical and laboratory evidence of hepatic toxicity may not be apparent 48 to 72 hours post-ingestion, and reach a maximum after 4 to 6 days.

WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN THE RECOMMENDED DOSAGE?
If you have taken more than the recommended dosage, consult a doctor or contact a poison control center right away, even if you seem well, as there is a risk of delayed serious liver damage. Quick medical attention is important for adults as well as for children even if you do not notice any signs or symptoms.

CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE

- **Liver Warning:** This product contains paracetamol. Severe liver damage may occur if:
  - o A child takes more than 5 doses in 24 hours, which is the maximum daily amount;
  - o An adult or child 12 years old and above takes more than 4 g of paracetamol in 24 hours;
  - o Taken with other medicines containing paracetamol (or acetaminophen);
- **Allergy alert:** Paracetamol may cause allergic reactions. Symptoms may include:
  - o Skin redness
  - o Blisters
  - o Skin rash or hives
  - o Swelling of the face, tongue or throat
  - o Difficulty swallowing
  - o Unexplained wheezing or shortness of breath
- Ask a doctor before use if patient has liver or kidney disease.
- Use with caution in patients with glucose-6-phosphate deficiency (G6PD), a hereditary condition which can lead to decreased blood cell count and possible hemolytic anemia (anemia resulting from destruction of red blood cells).
- Do not take more than the recommended dose.
- Do not use after the expiry date on the label.

140 mm

XXXX

196 mm

ARTWORK INFORMATION:	**P300000040493 IN BIOGESIC-PEDIA MELON LINE INSERT**				
■ **NEW** ☐ **EXISTING** rev no.		Reg. No.	DR-XY46366	Pantone/s:	Black
Old FG/SKU Code	180598	PMS No.	180598-01	Basis for Revision:	Change in Brand Name, Update in Packaging Text
New FG/SKU Code	N/A	KG/PLG No.	N/A		
Existing Barcode	N/A	Dimension	140 x 196 mm (±1mm) - spread	CCR No.:	TMP-CCR-0108
New Barcode	N/A	Existing PM Code	N/A		

Method of Proofing Performed (Division) ☐ Global Vision Inspection (*Provide GV Report to QA) ■ Manual Proofing

Reference Checklist (Division):
■ CPR/CPN/Approval of country ■ PMS ☐ Knifeguide or PL Guide ☐ Global Vision Report
☐ Superseded Artwork ☐ FDA Approved artwork ■ PM code and FG code ■ CCDS

Evaluation Checklist (QA):
☐ PM code and FG code ☐ Global Vision Report (if applicable)
☐ CPR/Approval of country ☐ Dimension (PMS & KG/PLG) ☐ Barcode (SAP) ☐ Color and Pantone
☐ Photo-eye(tubes/Sachet) ☐ Approved CCR ☐ Space for LEM ☐ CCDS
☐ Unilab logo and seal(CBG Guideline) ☐ Code bars (ISFG) ☐ Manufacturing Clause ☐ FDA Approved AW/Notification

Marketing Division: *Name, Signature, Date*
In-house Artist/Agency: N/A
Brand Personnel: N/A
Med. Director: Maria Lourdes Genuino, MD
Div. Head/ Prod. Dir.: N/A
Div. Pharmacist: Kathyrene Visco

QA-SMS-PM:
Evaluated By:
Noted By:

Disposition:
☐ APPROVED
☐ DISAPPROVED
Remarks:

U-F-SQM-044/01

Biogesic (Paracetamol)

500 mg Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-153

Biogesic: Official FDA Philippines registration record

Brand name	Biogesic
Generic name	Paracetamol
Dosage strength and form	500 mg Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Analgesic/Antipyretic
Registration number	DR-153
Manufacturer	UNILAB, Inc.

Biogesic: OTC forms registered under the Biogesic brand (capsule availability)

OTC capsule form registered under the Biogesic brand: NO. 7 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Biogesic	100 mg / mL Suspension (Oral Drops) [Orange Sugar Free]	Household Remedy (HR)	DRHR-1167
Biogesic	120 mg/5 mL Syrup (Melon Flavor)	Household Remedy (HR)	DRHR-1843
Biogesic	120 mg/5 mL Syrup (Orange Flavor)	Household Remedy (HR)	DRHR-1748
Biogesic	250 mg / 5 mL Syrup (Orange Flavor)	Household Remedy (HR)	DR-XY45639
Biogesic	250 mg/5mL Syrup (Melon Flavor) Syrup	Over-the-Counter (OTC) Drug	DR-XY46366
Biogesic 325	325 mg Tablet	Over-the-Counter (OTC) Drug	DRP-8763
Biogesic	500 mg Tablet	Over-the-Counter (OTC) Drug	DR-153

Biogesic: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-153 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Biogesic"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Biogesic 325 (Paracetamol)

325 mg Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DRP-8763

Biogesic 325: Official FDA Philippines registration record

Brand name	Biogesic 325
Generic name	Paracetamol
Dosage strength and form	325 mg Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Analgesic / Antipyretic
Registration number	DRP-8763
Manufacturer	PT Medifarma Laboratories

Biogesic 325: OTC forms registered under the Biogesic brand (capsule availability)

OTC capsule form registered under the Biogesic brand: NO. 7 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Biogesic	100 mg / mL Suspension (Oral Drops) [Orange Sugar Free]	Household Remedy (HR)	DRHR-1167
Biogesic	120 mg/5 mL Syrup (Melon Flavor)	Household Remedy (HR)	DRHR-1843
Biogesic	120 mg/5 mL Syrup (Orange Flavor)	Household Remedy (HR)	DRHR-1748
Biogesic	250 mg / 5 mL Syrup (Orange Flavor)	Household Remedy (HR)	DR-XY45639
Biogesic	250 mg/5mL Syrup (Melon Flavor) Syrup	Over-the-Counter (OTC) Drug	DR-XY46366
Biogesic 325	325 mg Tablet	Over-the-Counter (OTC) Drug	DRP-8763
Biogesic	500 mg Tablet	Over-the-Counter (OTC) Drug	DR-153

Biogesic 325: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRP-8763 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Biogesic"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Calpol (Paracetamol)

250 mg / 5 mL Suspension | Household Remedy (HR) | FDA Reg. No. DRHR-1964

Calpol: Official FDA Philippines registration record

Brand name	Calpol
Generic name	Paracetamol
Dosage strength and form	250 mg / 5 mL Suspension
Classification	Household Remedy (HR)
Pharmacologic category	Analgesic / Antipyretic
Registration number	DRHR-1964
Manufacturer	PT. Sterling Products Indonesia

Calpol: OTC forms registered under the Calpol brand (capsule availability)

OTC capsule form registered under the Calpol brand: NO. 5 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Calpol (Strawberry Flavor)	120 mg/ 5 mL Suspension	Household Remedy (HR)	DRHR-495
Calpol	120 mg/5 mL Suspension (Strawberry Flavor)	Over-the-Counter (OTC) Drug	DRHR-2132
Calpol	120mg/5mL Suspension (Strawberry Flavor)	Household Remedy (HR)	DRHR-2131
Calpol	250 mg / 5 mL Suspension	Household Remedy (HR)	DRHR-1964
Calpol	250mg/5ml Suspension (Orange Flavor)	Household Remedy (HR)	DRHR-913

Calpol: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-1964 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Calpol"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Calpol (Paracetamol)

120mg/5mL Suspension (Strawberry Flavor) | Household Remedy (HR) | FDA Reg. No. DRHR-2131

Calpol: Official FDA Philippines registration record

Brand name	Calpol
Generic name	Paracetamol
Dosage strength and form	120mg/5mL Suspension (Strawberry Flavor)
Classification	Household Remedy (HR)
Pharmacologic category	Analgesic/ Antipyretic
Registration number	DRHR-2131
Manufacturer	PT. Glaxo Wellcome Indonesia

Calpol: OTC forms registered under the Calpol brand (capsule availability)

OTC capsule form registered under the Calpol brand: NO. 5 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Calpol (Strawberry Flavor)	120 mg/ 5 mL Suspension	Household Remedy (HR)	DRHR-495
Calpol	120 mg/5 mL Suspension (Strawberry Flavor)	Over-the-Counter (OTC) Drug	DRHR-2132
Calpol	120mg/5mL Suspension (Strawberry Flavor)	Household Remedy (HR)	DRHR-2131
Calpol	250 mg / 5 mL Suspension	Household Remedy (HR)	DRHR-1964
Calpol	250mg/5ml Suspension (Orange Flavor)	Household Remedy (HR)	DRHR-913

Calpol: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-2131 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Calpol"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Dolfenal (Mefenamic Acid)

250 mg Film-Coated Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DRP-8716

Dolfenal: Official FDA Philippines registration record

Brand name	Dolfenal
Generic name	Mefenamic Acid
Dosage strength and form	250 mg Film-Coated Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Anti-Inflammatory and Anti-Rheumatic Products, Non-Steroids (Fenamates)
Registration number	DRP-8716
Manufacturer	Amherst Laboratories, Inc.

Dolfenal: OTC forms registered under the Dolfenal brand (capsule availability)

OTC capsule form registered under the Dolfenal brand: NO. 1 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Dolfenal	250 mg Film-Coated Tablet	Over-the-Counter (OTC) Drug	DRP-8716

Dolfenal: Official leaflet (Package Insert / Patient Information Leaflet)

An official leaflet for this registration is published on the FDA Verification Portal: PIL_01 (https://drive.google.com/file/d/12vv-Tt2jsdzDQh04-L4gA2MvHc0HsaTz/view?usp=drivesdk). Its text (OCR) and the original leaflet pages follow.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Dolfenal"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Dolfenal: Official leaflet text (OCR transcription)

Machine transcription of the official FDA leaflet for Reg. No. DRP-8716. OCR may contain errors; the original leaflet pages are attached after this section.

Dolfenal leaflet PIL_01

westmont
pharmcceulicals
ARTWORK SPECIFICATION
PRODUCT
DOLFENAL 250 mg
TABLET
ITIEM OE
COMPONENT/ PACK
INSERT
SIZE
BLACK
COLORS
ARTWORK DEVELOPMENT STATUS
DATE CREATED
2/09/06
DATE REVISED
11/26/19
SERIES OF
REVISIONS
CONTACT PERSON
G. VILLEGAS
Mefenamic Acid
GENERIC NAME
DR-XY2384
DIMENSION
196 × 280
G.VILLEGAS
CREATED BY
REVISED BY
G. VILLEGAS
OF PRINTS
UNILAB
Mefenamic Acid
itching, nausea, rashes, ringing in the ears (tinnitus), and vomiting.
bleeding/perforation, headache, heartburn, increased bleeding time,
function and increase blood levels of these medicines (e.g., digoxin)
Cardiac Glycosides: NSAIDs may worsen heart failure, reduce kidney
WHEN SHOULD YOU CONSULT YOUR DOCTOR?
Dolfenal
Less common undesirable effects (<1% of patients) include:
Ciclosporin and Tacrolimus: Increased risk of kidney damage
Ask a doctor before use if you are:
• Having problems or serious side effects from taking pain relievers or
Anti-inflammatory and Anti-Rheumatic Products,
Non-Steroids (Feñamates)
250 mg FILM-COATED TABLET
infection in the bloodstream or body tissues (sepsis), black or blue marks
Blood: Abnormal enlargement of the lymph nodes, anemia, bacterial
or bruises (ecchymosis), decreased hematocrit, low platelet count, low
white blood cell count (i.e., leukopenia, neutropenia, agranulocytosis,
Lithium and Methotrexate: Mefenamic acid increases the amount of
lithium and methotrexate in the blood, thus increasing the risk of toxicity
Medicines Affecting Liver Enzymes: Medicines which hinder the
fever reducers
• Taking aspirin or other NSAIDs
• Under a doctor's care for any serious condition
• Trying to become pregnant; NSAIDs may impair fertility in women. This
• Taking other medicines
FORMULATION
Each film-coated tablet contains:

eosinophilia, pancytopenia), red or purple discoloration of the skin (purpura)
enzymes responsible for the metabolism of mefenamic acid may alter its safety and efficacy
effect is reversible upon stopping the medicine.
Mefenamic Acid..
250 mg
Immune System: Allergic reactions including asthma. breathing difficulty.
Medicines that Lower Blood Sugar (for Diabetes): Mefenamic acid
Ask a doctor before use if:
Immune System: Allergic reactions including asthma, breathing difficulty,
• Stomach bleeding warning applies to you
Mefenamic acid (Dolfenal®) 250 mg Tablet: Pink, capsule-shaped
PRODUCT DESCRIPTION
film-coated tablet
or skin reactions [e.g., rapid swelling beneath the skin (angioedema), widespread scaling of the skin (exfoliative dermatitis), Stevens-Johnson syndrome, toxic epidermal necrolysis, and erythema multiforme]
may increase plasma levels of medicines used to treat diabetes (suifonylureas such as gliclazide, glipizide, glimepiride) by inhibiting their metabolism.
• You have heart problems, previous stroke or might be at risk of these conditions (e.g., high blood pressure, high cholesterol, diabetes, or if you are a smoker)
• You have a history of stomach problems, such as heartburn, Crohn's
This product contains mefenamic acid which belongs to a group of
WHAT IS IN THE MEDICINE?
medicines called non-steroidal anti-inflammatory drugs (NSAIDs). Like other NSAIDs, mefenamic acid works by changing the body's chemical response to pain, swelling and fever, resulting in relief of symptoms of inflammation (e.g., swelling, redness) and relief of pain.
sodium in blood, weight changes
Nervous System: Anxiety, coma, dream abnormalities, drowsiness, hallucination, inflammation of the meninges (meningitis), inflammation of
Metabolic: Appetite changes, fluid retention, high blood sugar, low
Psychiatric: Confusion, depression, nervousness
Mifepristone: Mefenamic acid should not be used for 8 to 12 days after mifepristone administration because it can reduce the effect of mifepristone.
Probenecid: May reduce the metabolism and excretion of mefenamic acid, thus increasing the risk of mefenamic acid toxicity.
in the lining of the rectum and colon)
• You suffer from systemic lupus erythematosus (SLE) or other
disease (inflammation of the digestive system) or ulcerative colitis (ulcers
• You have asthma
auto-immune diseases
• You have liver or kidney problems
WHAT IS THIS MEDICINE USED FOR?
• For the relief of mild to moderate pain including headache, dental pain,
STRENGTH OF THE MEDICINE
See Formulation.
(heavy menstrual period) due to dysfunctional causes
the optic nerve (optic neuritis), insomnia, loss of strength, numbness or impairment, inflammation or infection of the membrane lining the eyelids
tingling of the skin, seizures, sleepiness, tremors, vertigo
Special Senses: Blurred vision, ear pain, eye irritation/pain, hearing
(conjunctivitis), reversible loss of color vision, visual disturbances
Cardiovascular: Abnormal heart rhythms, congestive heart failure, heart
Zidovudine: An increased risk of blood problems such as bleeding into a joint (hemarthrosis) and bruising (hematoma) may result when given with
Quinolone Antibiotics: A possible increased risk of convulsions may occur with mefenamic acid
mefenamic acid
Tell your doctor if you are taking other medicines, including vitamins,
• You experience any of the following signs of stomach bleeding:
• New symptoms occur
Stop use and ask a doctor if:

• An allergic reaction occurs
o Feel faint
o Vomit blood
o Have bloody or black stools
o Have stomach pain that does not get better
• For the relief of pain associated with musculoskeletal and joint disorders including osteoarthritis and rheumatoid arthritis
attack, high blood pressure, inflammation of the walls of blood vessels, loss of consciousness, low blood pressure, palpitations, rapid heart rate, stroke
WHAT SHOULD YOU DO IF YOU MISS A DOSE?
supplements, and herbal medicines.
• Symptoms do not get better
• Headache is persistent
Like other NSAIDs, the lowest effective dose should be used for the shortest possible time. After observing the response to initial therapy with this product, the dose and frequency should be adjusted to suit individual patient needs.
HOW MUCH AND HOW OFTEN SHOULD YOU USE THIS MEDICINE?
the mucous linings in the mouth, inflammation of the pancreas, inflammation of the stomach, inflammation of the tongue, inflammation or
Lungs: Asthma, breathing difficulty, respiratory depression, pneumonia
Stomach/Gut: Anorexia, belching, dry mouth, dyspepsia, inflammation of irritation of the esophagus, passing black or bloody stools, passing stools
If a dose is missed, take the next dose if still needed for pain and the
HOW SHOULD YOU KEEP THIS MEDICINE?
Keep the product out of sight and reach of children.
subsequent dose every 8 hours thereafter.
Do not double the dose.
If you are pregnant or breastfeeding, ask a doctor before use. It is especially important not to use mefenamic acid during the last three because it may cause problems in the unborn child or complications during delivery.
ADVERSE DRUG REACTION REPORTING STATEMENT
This medicine is given orally and may be taken with food or milk if stomach upset occurs.
with excess fat, rectal bleeding, vomiting with blood, worsen inflammation of the intestine or colon
Store at temperatures not exceeding 30°C.
SIGNS AND SYMPTOMS OF OVERDOSE
and report to the FDA at www.fda.gov.ph AND Unilab at +632-8-UNILAB-1
For suspected adverse drug reaction, seek medical attention immediately (+632-8-864522-1) for Metro Manila or toll-free +1-800-10-UNILAB-1 for
Usual Recommended Dose in Adults and Adolescents older than 14 years:
Liver: Abnormal liver function tests, abnormal yellowing of the skin (jaundice), inflammation of the liver (hepatitis), liver damage, liver failure
epigastric pain. Gastrointestinal bleeding can occur. Hypertension, acute
Symptoms following acute overdose of NSAIDs, such as mefenamic acid, undesirable effects, you can help provide more information on the safety of this medicine.
provinces, or e-mail productsafety@unilab.com.ph. By reporting
Skin: Hair loss, hives, itching, perspiration, photosensitivity, swelling of
kipry osy ay e y odsse esr sy ecur.
1 to 2 tablets every 8 hours while symptoms persist ("period pains" and
For the relief of dysmenorrhea and premenstrual syndrome (PMS):
other associated symptoms), or, as directed by a doctor.
Kidneys: Abnormal kidney function, decreased urine output, excess serum proteins in the urine, increased urine output, inflammation of the
the face or voice box
Anaphylactoid reactions have been reported with recommended doses of NSAIDs, and may occur following an overdose.
WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN THE
100s.
AVAILABILITY:
• For menstrual cramps (period pain), mefenamic acid is usually taken

during each period as soon as the cramps begin and continued for a
tubules (interstitial nephritis), kidney failure, pain when urinating, blood in
kidneys or urinary bladder, inflammation of the spaces between the kidney
RECOMMENDED DOSAGE?
If more than the recommended dosage has been given, consult a doctor
DRP-8716
few days until the pain goes away.
the urine
or contact a Poison Control Center right away.
Date of Revision: March 2020
• For heavy menstrual periods, mefenamic acid is taken from the
advice.
beginning of the period and continued according to your doctor's
Body as a Whole: Death, fatigue, fever, general ill feeling, infection,
organ failure
CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE
• Allergy Alert: Mefenamic acid may cause a severe allergic reaction
Date of First Authorization: May 25, 2007
• Do not take for more than 7 days unless directed by a doctor.
which may include:
Reg. IPOPHIL
Investigations: False-positive for bile in the urine
o Hives (elevated, whitish or reddish patches on the skin with
WHEN SHOULD YOU NOT TAKE THIS MEDICINE?
severe itching or pricking sensations);
• If you are allergic to mefenamic acid, aspirin or any other NSAID, or to
WHAT OTHER MEDICINE OR FOOD SHOULD BE AVOIDED WHILE
o Shock (anaphylactic), a hypersensitivity reaction resulting to
other ingredients in the product
TAKING THIS MEDICINE?
generalized skin lesions and itchiness, followed by vascular
angioedema (rapid swelling that occurs in the tissue just below the
• If you are currently taking aspirin or other NSAIDs
• If you have bronchospasm (constriction of air passages of the lungs),
surface of the skin), nasal polyps or allergic-type reactions after taking
aspirin or other NSAIDs
• If you have had or are suffering from stomach ulcers, bleeding or other
stomach problems such as inflammatory bowel disease
• If you had diarrhea with this medicine in the past. Diarrhea may recur if
Aspirin and Other NSAIDs: Mefenamic acid should not be given
together with aspirin, salicylates or other NSAIDs (e.g., ketorolac,
etoricoxib) because of the increased risk of stomach and intestinal
bleeding.
Corticosteroids, Selective Serotonin-Reuptake Inhibitors (SSRIs),
ibuprofen, naproxen) including COX-2 inhibitors (e.g., celecoxib,
and Anti-platelet Agents: May increase the rísk of stomach bleeding with
• Stomach Bleeding Warning: This product contains an NSAID, which
o Facial swelling
o Skin reddening
o Asthma (wheezing)
o Skin rash or skin blisters
distress
collapse and often accompanied by life-threatening respiratory
Manufactured by
No. 66 United Street, Mandaluyong City
Metro Manila, Philippines
AMHERST LABORATORIES, INC.
UNILAB Pharma Campus
Barangay Mamplasan, Biñan, Laguna
Philippines for UNILAB, Inc.
you take mefenamic acid again.
mefenamic acid
may cause severe stomach bleeding. The chance is higher if you:
• If you have severe heart, liver or kidney disease
o Are age 60 years or older
• High potassium in the blood
• Bleeding in the brain or other bleeding disorders

aminoglycoside toxicity by decreasing the excretion of aminoglycoside
Aminoglycosides: NSAIDs, such as mefenamic acid, increase
o Have had stomach ulcers or bleeding problems
o Take a blood thinning (anticoagulant) or steroid medicine
UNILAB Consumer Care
• Dengue fever, since NSAIDs may increase the risk of bleeding
antibiotics (e.g., amikacin, gentamicin, neomycin, tobramycin).
• During the last 3 months of pregnancy
NSAIDs (aspirin, ibuprofen or others)
8-UNILAB-1
• Breastfeeding women, because of the potential for serious
undesirable effects in breastfed infants
Antacids: The plasma levels of mefenamic acid may be increased when
taken together with antacids such as magnesium hydroxide.
o Have 3 or more alcoholic drinks everyday while using this product
o Take more or for a longer time than directed
8-864522-1
• Right before or after heart surgery
• Children and adolescents younger than 14 years old
UNDESIRABLE EFFECTS
The most frequently reported undesirable effects (1 to 10%) of
these drugs; diuretics may also cause acute reduction in kidney function
of NSAIDs
Antihypertensives and Diuretics: NSAIDS may reduce the effect of
• The risk of heart attack or stroke may increase if you use more than or
• Mefenamic acid should be used with caution in elderly patients suffering
longer than directed.
Provincial Toll Free: 1-800-10-8645221
Website: www.unilab.com.ph
Email: info@unilab.com.ph
Trusted Quality Healthgat.
UNILAB
mefenamic acid include abdominal pain, abnormal kidney function,
Blood Thinning Medicines (Anticoagulants): NSAIDs may enhance
from dehydration and kidney disease.
anemia, constipation, diarrhea, dizziness, dyspepsia, edema, elevated
the effects of these agents (e.g., warfarin). An increase in the International
liver enzymes (e.g., alanine transaminase or aspartate transaminase),
• Do not use after the expiry date on the label.
flatulence,
gastrointestinal
ulcers
(gastric/duodenal),
gross
bleeding, may be observed when using paracetamol and warfarin at the
same time.
V
7

westmont pharmaceuticals

ARTWORK SPECIFICATION

PRODUCT	ITEM CODE	COMPONENT/ PACK SIZE	COLORS
DOLFENAL 250 mg TABLET	**P300000030292**	**INSERT**	**BLACK**
CONTACT PERSON	GENERIC NAME	DR # / FR #	DIMENSION
G. VILLEGAS	**Mefenamic Acid**	**DR-XY2384**	**196 X 280**

ARTWORK DEVELOPMENT STATUS

DATE CREATED	DATE REVISED	SERIES OF REVISIONS
2/09/06	**11/26/19**	
CREATED BY **G.VILLEGAS**	REVISED BY **G. VILLEGAS**	# OF PRINTS

UNILAB

Mefenamic Acid Dolfenal®

250 mg FILM-COATED TABLET

Anti-inflammatory and Anti-Rheumatic Products, Non-Steroids (Fenamates)

FORMULATION
Each film-coated tablet contains:
Mefenamic Acid…… 250 mg

PRODUCT DESCRIPTION
Mefenamic acid (Dolfenal®) 250 mg Tablet: Pink, capsule-shaped film-coated tablet

WHAT IS IN THE MEDICINE?
This product contains mefenamic acid which belongs to a group of medicines called non-steroidal anti-inflammatory drugs (NSAIDs). Like other NSAIDs, mefenamic acid works by changing the body's chemical response to pain, swelling and fever, resulting in relief of symptoms of inflammation (e.g., swelling, redness) and relief of pain.

STRENGTH OF THE MEDICINE
See Formulation.

WHAT IS THIS MEDICINE USED FOR?
• For the relief of mild to moderate pain including headache, dental pain, pain after childbirth or surgery, primary dysmenorrhea, and menorrhagia (heavy menstrual period) due to dysfunctional causes

• For the relief of pain associated with musculoskeletal and joint disorders including osteoarthritis and rheumatoid arthritis

HOW MUCH AND HOW OFTEN SHOULD YOU USE THIS MEDICINE?
Like other NSAIDs, the lowest effective dose should be used for the shortest possible time. After observing the response to initial therapy with this product, the dose and frequency should be adjusted to suit individual patient needs.

This medicine is given orally and may be taken with food or milk if stomach upset occurs.

Usual Recommended Dose in Adults and Adolescents older than 14 years:
1 to 2 tablets every 8 hours, as needed, or, as directed by a doctor.

For the relief of dysmenorrhea and premenstrual syndrome (PMS):
1 to 2 tablets every 8 hours while symptoms persist ("period pains" and other associated symptoms), or, as directed by a doctor.

- For menstrual cramps (period pain), mefenamic acid is usually taken during each period as soon as the cramps begin and continued for a few days until the pain goes away.
- For heavy menstrual periods, mefenamic acid is taken from the beginning of the period and continued according to your doctor's advice.
- Do not take for more than 7 days unless directed by a doctor.

WHEN SHOULD YOU NOT TAKE THIS MEDICINE?
• If you are allergic to mefenamic acid, aspirin or any other NSAID, or to other ingredients in the product
• If you are currently taking aspirin or other NSAIDs
• If you have bronchospasm (constriction of air passages of the lungs), angioedema (rapid swelling that occurs in the tissue just below the surface of the skin), nasal polyps or allergic-type reactions after taking aspirin or other NSAIDs
• If you have had or are suffering from stomach ulcers, bleeding or other stomach problems such as inflammatory bowel disease
• If you had diarrhea with this medicine in the past. Diarrhea may recur if you take mefenamic acid again.
• If you have severe heart, liver or kidney disease
• Bleeding in the brain or other bleeding disorders
• High potassium in the blood
• Dengue fever, since NSAIDs may increase the risk of bleeding
• During the last 3 months of pregnancy
• Breastfeeding women, because of the potential for serious undesirable effects in breastfed infants
• Children and adolescents younger than 14 years old
• Right before or after heart surgery

UNDESIRABLE EFFECTS
The most frequently reported undesirable effects (1 to 10%) of mefenamic acid include abdominal pain, abnormal kidney function, anemia, constipation, diarrhea, dizziness, dyspepsia, edema, elevated liver enzymes (e.g., alanine transaminase or aspartate transaminase), flatulence, gastrointestinal ulcers (gastric/duodenal), gross bleeding/perforation, headache, heartburn, increased bleeding time, itching, nausea, rashes, ringing in the ears (tinnitus), and vomiting.

Less common undesirable effects (<1% of patients) include:

Blood: Abnormal enlargement of the lymph nodes, anemia, bacterial infection in the bloodstream or body tissues (sepsis), black or blue marks or bruises (ecchymosis), decreased hematocrit, low platelet count, low white blood cell count (i.e., leukopenia, neutropenia, agranulocytosis, eosinophilia, pancytopenia), red or purple discoloration of the skin (purpura)

Immune System: Allergic reactions including asthma, breathing difficulty, or skin reactions [e.g., rapid swelling beneath the skin (angioedema), widespread scaling of the skin (exfoliative dermatitis), Stevens-Johnson syndrome, toxic epidermal necrolysis, and erythema multiforme]

Metabolic: Appetite changes, fluid retention, high blood sugar, low sodium in blood, weight changes

Psychiatric: Confusion, depression, nervousness

Nervous System: Anxiety, coma, dream abnormalities, drowsiness, hallucination, inflammation of the meninges (meningitis), inflammation of the optic nerve (optic neuritis), insomnia, loss of strength, numbness or tingling of the skin, seizures, sleepiness, tremors, vertigo

Special Senses: Blurred vision, ear pain, eye irritation/pain, hearing impairment, inflammation or infection of the membrane lining the eyelids (conjunctivitis), reversible loss of color vision, visual disturbances

Cardiovascular: Abnormal heart rhythms, congestive heart failure, heart attack, high blood pressure, inflammation of the walls of blood vessels, loss of consciousness, low blood pressure, palpitations, rapid heart rate, stroke

Lungs: Asthma, breathing difficulty, respiratory depression, pneumonia

Stomach/Gut: Anorexia, belching, dry mouth, dyspepsia, inflammation of the mucous linings in the mouth, inflammation of the pancreas, inflammation of the stomach, inflammation of the tongue, inflammation or irritation of the esophagus, passing black or bloody stools, passing stools with excess fat, rectal bleeding, vomiting with blood, worsen inflammation of the intestine or colon

Liver: Abnormal liver function tests, abnormal yellowing of the skin (jaundice), inflammation of the liver (hepatitis), liver damage, liver failure

Skin: Hair loss, hives, itching, perspiration, photosensitivity, swelling of the face or voice box

Kidneys: Abnormal kidney function, decreased urine output, excess serum proteins in the urine, increased urine output, inflammation of the kidneys or urinary bladder, inflammation of the spaces between the kidney tubules (interstitial nephritis), kidney failure, pain when urinating, blood in the urine

Body as a Whole: Death, fatigue, fever, general ill feeling, infection, organ failure

Investigations: False-positive for bile in the urine

WHAT OTHER MEDICINE OR FOOD SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?
Aspirin and Other NSAIDs: Mefenamic acid should not be given together with aspirin, salicylates or other NSAIDs (e.g., ketorolac, ibuprofen, naproxen) including COX-2 inhibitors (e.g., celecoxib, etoricoxib) because of the increased risk of stomach and intestinal bleeding.

Corticosteroids, Selective Serotonin-Reuptake Inhibitors (SSRIs), and Anti-platelet Agents: May increase the risk of stomach bleeding with mefenamic acid

Aminoglycosides: NSAIDs, such as mefenamic acid, increase aminoglycoside toxicity by decreasing the excretion of aminoglycoside antibiotics (e.g., amikacin, gentamicin, neomycin, tobramycin).

Antacids: The plasma levels of mefenamic acid may be increased when taken together with antacids such as magnesium hydroxide.

Antihypertensives and Diuretics: NSAIDS may reduce the effect of these drugs; diuretics may also cause acute reduction in kidney function of NSAIDs

Blood Thinning Medicines (Anticoagulants): NSAIDs may enhance the effects of these agents (e.g., warfarin). An increase in the International Normalized Ratio (INR), which may serve as a sign of increased risk for bleeding, may be observed when using paracetamol and warfarin at the same time.

Cardiac Glycosides: NSAIDs may worsen heart failure, reduce kidney function and increase blood levels of these medicines (e.g., digoxin)

Ciclosporin and Tacrolimus: Increased risk of kidney damage

Lithium and Methotrexate: Mefenamic acid increases the amount of lithium and methotrexate in the blood, thus increasing the risk of toxicity

Medicines Affecting Liver Enzymes: Medicines which hinder the enzymes responsible for the metabolism of mefenamic acid may alter its safety and efficacy

Medicines that Lower Blood Sugar (for Diabetes): Mefenamic acid may increase plasma levels of medicines used to treat diabetes (sulfonylureas such as gliclazide, glipizide, glimepiride) by inhibiting their metabolism.

Mifepristone: Mefenamic acid should not be used for 8 to 12 days after mifepristone administration because it can reduce the effect of mifepristone.

Probenecid: May reduce the metabolism and excretion of mefenamic acid, thus increasing the risk of mefenamic acid toxicity.

Quinolone Antibiotics: A possible increased risk of convulsions may occur with mefenamic acid

Zidovudine: An increased risk of blood problems such as bleeding into a joint (hemarthrosis) and bruising (hematoma) may result when given with mefenamic acid

Tell your doctor if you are taking other medicines, including vitamins, supplements, and herbal medicines.

WHAT SHOULD YOU DO IF YOU MISS A DOSE?
If a dose is missed, take the next dose if still needed for pain and the subsequent dose every 8 hours thereafter.

Do not double the dose.

HOW SHOULD YOU KEEP THIS MEDICINE?
Keep the product out of sight and reach of children.
Store at temperatures not exceeding 30°C.

SIGNS AND SYMPTOMS OF OVERDOSE
Symptoms following acute overdose of NSAIDs, such as mefenamic acid, are usually limited to lethargy (fatigue), drowsiness, nausea, vomiting and epigastric pain. Gastrointestinal bleeding can occur. Hypertension, acute kidney failure, respiratory depression and coma may also occur. Anaphylactoid reactions have been reported with recommended doses of NSAIDs, and may occur following an overdose.

WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN THE RECOMMENDED DOSAGE?
If more than the recommended dosage has been given, consult a doctor or contact a Poison Control Center right away.

CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE
• **Allergy Alert:** Mefenamic acid may cause a severe allergic reaction which may include:
  - o Hives (elevated, whitish or reddish patches on the skin with severe itching or pricking sensations);
  - o Shock (anaphylactic), a hypersensitivity reaction resulting to generalized skin lesions and itchiness, followed by vascular collapse and often accompanied by life-threatening respiratory distress
  - o Facial swelling
  - o Skin reddening
  - o Asthma (wheezing)
  - o Skin rash or skin blisters

• **Stomach Bleeding Warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
  - o Are age 60 years or older
  - o Have had stomach ulcers or bleeding problems
  - o Take a blood thinning (anticoagulant) or steroid medicine
  - o Take other medicines containing prescription or nonprescription NSAIDs (aspirin, ibuprofen or others)
  - o Have 3 or more alcoholic drinks everyday while using this product
  - o Take more or for a longer time than directed

• The risk of heart attack or stroke may increase if you use more than or longer than directed.

• Mefenamic acid should be used with caution in elderly patients suffering from dehydration and kidney disease.

• Do not use after the expiry date on the label.

WHEN SHOULD YOU CONSULT YOUR DOCTOR?

Ask a doctor before use if you are:
• Having problems or serious side effects from taking pain relievers or fever reducers
• Taking aspirin or other NSAIDs
• Taking other medicines
• Under a doctor's care for any serious condition
• Trying to become pregnant; NSAIDs may impair fertility in women. This effect is reversible upon stopping the medicine.

Ask a doctor before use if:
• Stomach bleeding warning applies to you
• You have heart problems, previous stroke or might be at risk of these conditions (e.g., high blood pressure, high cholesterol, diabetes, or if you are a smoker)
• You have a history of stomach problems, such as heartburn, Crohn's disease (inflammation of the digestive system) or ulcerative colitis (ulcers in the lining of the rectum and colon)
• You have liver or kidney problems
• You have asthma
• You suffer from systemic lupus erythematosus (SLE) or other auto-immune diseases

Stop use and ask a doctor if:
• You experience any of the following signs of stomach bleeding:
  - o Feel faint
  - o Vomit blood
  - o Have bloody or black stools
  - o Have stomach pain that does not get better
• An allergic reaction occurs
• New symptoms occur
• Symptoms do not get better
• Headache is persistent

If you are pregnant or breastfeeding, ask a doctor before use. It is especially important not to use mefenamic acid during the last three months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

ADVERSE DRUG REACTION REPORTING STATEMENT
For suspected adverse drug reaction, seek medical attention immediately and report to the FDA at www.fda.gov.ph AND Unilab at +632-8-**UNILAB**-1 (+632-8-**864522**-1) for Metro Manila or toll-free +1-800-10-UNILAB-1 for provinces, or e-mail productsafety@unilab.com.ph. By reporting undesirable effects, you can help provide more information on the safety of this medicine.

AVAILABILITY:
Mefenamic Acid (Dolfenal®) 250 mg film-coated tablet in flex foil, box of 100s.

DRP-8716

Date of Revision: March 2020

Date of First Authorization: May 25, 2007

Reg. IPOPHIL

Manufactured by
AMHERST LABORATORIES, INC.
UNILAB Pharma Campus
Barangay Mamplasan, Biñan, Laguna
Philippines for **UNILAB, Inc.**
No. 66 United Street, Mandaluyong City
Metro Manila, Philippines

UNILAB Consumer Care
8-UNILAB-1
8-864522-1
Provincial Toll Free: 1-800-10-8645221
Email: info@unilab.com.ph
Website: www.unilab.com.ph

UNILAB
Trusted Quality Healthcare

P300000030292

280 mm

196 mm

PANTONE BLACK C

Efficascent (Menthol + Methyl Salicylate + Camphor)

140 mg/100 mg/35 mg per gram Ointment | Household Remedy (HR) | FDA Reg. No. DRHR-2201

Efficascent: Official FDA Philippines registration record

Brand name	Efficascent
Generic name	Menthol + Methyl Salicylate + Camphor
Dosage strength and form	140 mg/100 mg/35 mg per gram Ointment
Classification	Household Remedy (HR)
Pharmacologic category	Counterirritant
Registration number	DRHR-2201
Manufacturer	International Pharmaceuticals, Inc.

Efficascent: OTC forms registered under the Efficascent brand (capsule availability)

OTC capsule form registered under the Efficascent brand: NO. 20 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Efficascent Relaxscent Balm	120 mg / 50 mg / 10 mg per gram Balm	Household Remedy (HR)	DRHR-2544
Efficascent Oil	13.846 mL / 3.846 g / 0.423 g per 100 mL Liniment	Household Remedy (HR)	DRHR-247
Efficascent Ointment	14 g / 10 g / 3.5 g per 100 g Ointment	Household Remedy (HR)	DRHR-1270
Efficascent	140 mg/100 mg/35 mg per gram Ointment	Household Remedy (HR)	DRHR-2201
Efficascent Oil	168.73 mg / 38.46 mg / 4.23 mg per mL Liniment	Household Remedy (HR)	DRHR-2119
Efficascent Oil	168.73 mg/38.46 mg/4.23 mg per mL Liniment	Household Remedy (HR)	DRHR-2272
Efficascent Oil – Gold	19.80 g/13.20 g/4.40 g per 100 mL Topical Solution	Household Remedy (HR)	DRHR-1787
Efficascent Boost Pain Relief Massage Roll-On	191.60 mg / 67.06 mg / 38.32 mg per mL Lotion	Household Remedy (HR)	DRHR-2169
Efficascent Boost Pain Relief Massage Roll-on	191.60 mg/67.06 mg/38.32 mg per mL Lotion	Household Remedy (HR)	DRHR-2014
Efficascent Oil – Extra Strength	21 g/ 7.5 g/ 6 g per 100 mL Liniment	Household Remedy (HR)	DRHR-24
Efficascent Oil - Extra Strength	225 mg / 75 mg / 60 mg per mL Liniment	Household Remedy (HR)	DRHR-2120
Efficascent Oil – Extra Strength	225 mg / 75 mg / 60 mg per mL Liniment	Household Remedy (HR)	DRHR-2271
Efficascent Oil - Extreme	30 g / 7.5 g / 6 g per 100 mL Liniment	Household Remedy (HR)	DRHR-1323
Efficascent Oil – Extreme	300 mg / 75 mg / 60 mg per mL Liniment	Household Remedy (HR)	DRHR-2121

Brand name	Dosage strength and form	Classification	Reg. No.
Efficascent Relaxscent Oil	45.3 g / 4.36 g / 4.36 g per 100 mL Liniment	Household Remedy (HR)	DRHR-1910
Efficascent Relaxscent Oil	453.44 mg / 43.6 mg /43.6 mg per mL Liniment	Household Remedy (HR)	DRHR-2270
Efficascent Relaxscent Oil – Calm	454.5 mg / 52.4 mg / 43.7 mg per mL Liniment	Household Remedy (HR)	DRHR-1961
Efficascent Relaxscent Oil – Calm	454.5 mg / 52.4 mg / 43.7 mg per mL Liniment (Lavender Scent)	Household Remedy (HR)	DRHR-2042
Efficascent Relaxscent Oil – Refresh	454.5 mg / 52.4 mg / 43.7 mg per mL Liniment (Lemon Scent)	Household Remedy (HR)	DRHR-2089
Efficascent Relaxscent Oil – Intense	482.4 mg / 52.6 mg /43.9 mg per mL Liniment	Household Remedy (HR)	DRHR-2143

Efficascent: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-2201 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Efficascent"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Efficascent Oil (Methyl Salicylate + Camphor + Menthol)

168.73 mg / 38.46 mg / 4.23 mg per mL Liniment | Household Remedy (HR) | FDA Reg. No. DRHR-2119

Efficascent Oil: Official FDA Philippines registration record

Brand name	Efficascent Oil
Generic name	Methyl Salicylate + Camphor + Menthol
Dosage strength and form	168.73 mg / 38.46 mg / 4.23 mg per mL Liniment
Classification	Household Remedy (HR)
Pharmacologic category	Counterirritant
Registration number	DRHR-2119
Manufacturer	International Pharmaceuticals, Inc.

Efficascent Oil: OTC forms registered under the Efficascent brand (capsule availability)

OTC capsule form registered under the Efficascent brand: NO. 20 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Efficascent Relaxscent Balm	120 mg / 50 mg / 10 mg per gram Balm	Household Remedy (HR)	DRHR-2544
Efficascent Oil	13.846 mL / 3.846 g / 0.423 g per 100 mL Liniment	Household Remedy (HR)	DRHR-247
Efficascent Ointment	14 g / 10 g / 3.5 g per 100 g Ointment	Household Remedy (HR)	DRHR-1270
Efficascent	140 mg/100 mg/35 mg per gram Ointment	Household Remedy (HR)	DRHR-2201
Efficascent Oil	168.73 mg / 38.46 mg / 4.23 mg per mL Liniment	Household Remedy (HR)	DRHR-2119
Efficascent Oil	168.73 mg/38.46 mg/4.23 mg per mL Liniment	Household Remedy (HR)	DRHR-2272
Efficascent Oil – Gold	19.80 g/13.20 g/4.40 g per 100 mL Topical Solution	Household Remedy (HR)	DRHR-1787
Efficascent Boost Pain Relief Massage Roll-On	191.60 mg / 67.06 mg / 38.32 mg per mL Lotion	Household Remedy (HR)	DRHR-2169
Efficascent Boost Pain Relief Massage Roll-on	191.60 mg/67.06 mg/38.32 mg per mL Lotion	Household Remedy (HR)	DRHR-2014
Efficascent Oil – Extra Strength	21 g/ 7.5 g/ 6 g per 100 mL Liniment	Household Remedy (HR)	DRHR-24
Efficascent Oil - Extra Strength	225 mg / 75 mg / 60 mg per mL Liniment	Household Remedy (HR)	DRHR-2120
Efficascent Oil – Extra Strength	225 mg / 75 mg / 60 mg per mL Liniment	Household Remedy (HR)	DRHR-2271
Efficascent Oil - Extreme	30 g / 7.5 g / 6 g per 100 mL Liniment	Household Remedy (HR)	DRHR-1323
Efficascent Oil – Extreme	300 mg / 75 mg / 60 mg per mL Liniment	Household Remedy (HR)	DRHR-2121

Brand name	Dosage strength and form	Classification	Reg. No.
Efficascent Relaxscent Oil	45.3 g / 4.36 g / 4.36 g per 100 mL Liniment	Household Remedy (HR)	DRHR-1910
Efficascent Relaxscent Oil	453.44 mg / 43.6 mg /43.6 mg per mL Liniment	Household Remedy (HR)	DRHR-2270
Efficascent Relaxscent Oil – Calm	454.5 mg / 52.4 mg / 43.7 mg per mL Liniment	Household Remedy (HR)	DRHR-1961
Efficascent Relaxscent Oil – Calm	454.5 mg / 52.4 mg / 43.7 mg per mL Liniment (Lavender Scent)	Household Remedy (HR)	DRHR-2042
Efficascent Relaxscent Oil – Refresh	454.5 mg / 52.4 mg / 43.7 mg per mL Liniment (Lemon Scent)	Household Remedy (HR)	DRHR-2089
Efficascent Relaxscent Oil – Intense	482.4 mg / 52.6 mg /43.9 mg per mL Liniment	Household Remedy (HR)	DRHR-2143

Efficascent Oil: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-2119 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Efficascent"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Flanax (Naproxen Sodium)

220 mg Film-Coated Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY46695

Flanax: Official FDA Philippines registration record

Brand name	Flanax
Generic name	Naproxen Sodium
Dosage strength and form	220 mg Film-Coated Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Non-Steroidal Anti-inflammatory Drug
Registration number	DR-XY46695
Manufacturer	Interphil Laboratories, Inc.

Flanax: OTC forms registered under the Flanax brand (capsule availability)

OTC capsule form registered under the Flanax brand: NO. 7 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
FlanaxHot	102 mg/ 54.35 mg/ 13.65 mg per gram Topical Cream	Household Remedy (HR)	DRHR-2228
FlanaxACE HOT	2 g/1g/800 mg / 13 mg per 100 g Transdermal Patch	Household Remedy (HR)	DRHR-2413
Flanax	220 mg Film-Coated Tablet	Over-the-Counter (OTC) Drug	DR-XY46695
Flanax	275 mg Film-Coated Tablet	Over-the-Counter (OTC) Drug	DR-XY46696
FlanaxACE	4.98/28/1.48/900 mg per 100 g Transdermal Patch	Household Remedy (HR)	DRHR-2412
FlanaxCool	40 mg/g (4% w/w) Topical Gel	Household Remedy (HR)	DRHR-2227
FlanaxPlus	5 mg/ 102 mg/ 54.3 mg/ 13.7 mg per gram Topical Gel	Over-the-Counter (OTC) Drug	DR-XY48481

Flanax: Official leaflet (Package Insert / Patient Information Leaflet)

An official leaflet for this registration is published on the FDA Verification Portal: PIL_01 (https://drive.google.com/file/d/1z8gqd63aGKBVn9Wemtbns5Ws39L94f7f/view?usp=drivesdk). Its text (OCR) and the original leaflet pages follow.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Flanax"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Flanax: Official leaflet text (OCR transcription)

Machine transcription of the official FDA leaflet for Reg. No. DR-XY46695. OCR may contain errors; the original leaflet pages are attached after this section.

Flanax leaflet PIL_01

barcode
Naproxen Sodium
Flanax
220 mg
TAISHO
Non-Steroidal Anti-inflammatory Drug
1.
Description of the Product
Naproxen Sodium (Flanax®) is a blue, opaque film-coated, modified oval-shaped tablet.
2.
What is in the Medicine?
This product contains Naproxen Sodium which belongs to a group of medicines called non-steroidal anti-inflammatory drugs (also known as NSAIDs). Like other NSAIDs, Naproxen Sodium works by changing the swelling, redness) and relief of pain and/or fever.
3.
Strength of the Medicine
One tablet contains 220 mg of the active ingredient, Naproxen sodium (or the sodium salt of naproxen) – made from a combination of Naproxen 200 mg and 20 mg sodium (of low content, equating to a slice of wheat bread).
4.
What is this Medicine used for?
Naproxen sodium (Flanax®) can be used to relieve headache, toothache, muscular ache and backache, pain of cold, and fever.
5.
How much and how often should you use this Medicine?
Each dose should be taken with water on full stomach after eating. Adults: 1 tablet every 12 hours while symptoms persist. Elderly (>65): 1 tablet per day and no more than 2 tablets per day. Take Naproxen only if with pain and stop when pain is gone.
6.
When should you not take this Medicine?
Stomach and intestinal pain should not be treated with Naproxen Sodium (Flanax®).
Naproxen Sodium (Flanax®) should not be taken by patients with: a) history of disease of the stomach or of the intestines (ulcer), b) blood coagulation disorders or are receiving drug therapy that interferes with blood coagulation; c) kidney impairment; d) allergy to Naproxen; e) allergy to aspirin or other anti-inflammatory and pain relieving drugs, f) history of Cardiovascular Arrest, Myocardial Infarction, Coronary Artery Bypass Graft, reactions in patients allergic to aspirin and to other anti-inflammatory or pain relieving drugs.
7.
Care that should be taken when taking this Medicine?
A doctor should be consulted if the pain persists or gets worse. A doctor or a pharmacist should be consulted if: a) any other drugs are taken on a regular basis; b) heartburn, upset stomach or stomach pain occurs; c) even Naproxen Sodium (Flanax®) should not be taken for more than 10 consecutive days, unless directed by a doctor. Do not exceed the indicated dosage and immediately consult a doctor in the case of an overdose.
8.
Undesirable Effects of this Medicine
This medicine can cause varying degrees of inconvenient effects in certain people. The most frequent adverse events are - abdominal discomfort, stomach discomfort and heartburn, headache, nausea, vomiting, peripheral edema, allergy, ringing in the ears, skin rash, urticaria, pruritus, and dizziness.
In case of unusual reactions, stop treatment immediately and consult your doctor.
9.
What other medicine or food should be avoided while using this Medicine?
When taking Naproxen, ask your doctor before using other medications for fever, swelling or flu-like
Diclofenac, etc)
Avos na the
risk for gastrointestinal bleeding if taken together with Naproxen
Avoid taking alcohol since this can also increase the risk for gastrointestinal bleeding
Client: TAISHO
ARTWORK SPECIFICATION
Job Description: Flanax 220mg Insert
Agent:
File Name:
Artist:

JOEY
TAISHO_Flanax 220mg Insert_120622
Others:...
ARTWORK DEVELOPMENT
Revision Number:
Item Code:
Revised/Prepared By:
Date Started: Nov. 02, 2022
Revision Date:
Revisions Summary:
Spread Size:
Folded Size/
No. of Outs:
Fla
Dimension:
139.7mm x 215.9mm
CJ
GOODREAM
CREATIVES
PRINTERS CORPORATION
Co-administration with anticoagulants or blood thinning agents such as warfarin or dicumarol or antiplatelet drugs may increase the risk of gastrointestinal bleeding
For patients on corticosteroids, taking Naproxen or any other NSAID increases the chance of
Enalapril, Lisinopril, etc.) and beta-blockers (Propranolol, Carvedilol, Metroprolol, etc.) and Angiotensin Receptor Blockers (Losartan, Telmisartan)
Naproxen may reduce the effect of some diuretics like thiazides and furosemide
Taking Naproxen with Lithium may increase the levels of Lithium in the body which may increase the risk of toxicity
Taking Cholestyramine with Naproxen may affect the absorption of Naproxen
Taking Methotrexate with Naproxen may increase the risk for toxicity for methotrexate
Combining antidepressants such as selective serotonin reuptake inhibitors (SSRI) like Fluoxetine, Citalopram, Paroxetine, etc with Naproxen can increase the risk of GIT bleeding
10.
What should you do if you miss a dose?
If you skip or miss a dose, take the next dose particularly if there is still pain and inflammation and continue with your regular dosing schedule of every 8 to 12 hours. Do not take a double dose to make up for a missed one.
11.
Signs and Symptoms of Overdose
Drowsiness, heart burn, indigestion, nausea or vomiting
12.
What to do when you have taken more than the recommended dosage?
Should a patient ingest more than a large quantity of Naproxen Sodium (Flanax) accidentally or purposefully, seek medical attention immediately. The stomach may be emptied, and usual supportive measures employed.
13.
How should you keep this Medicine?
Naproxen Sodium (Flanax®) should be kept out of reach of children.
Store at temperatures not exceeding 30°C.
Do not use Naproxen Sodium (Flanax®) after the expiry date shown on the carton or blister strip.
14.
When should you consult your doctor?
For further information, consult your doctor or pharmacist who has detailed scientific information.
15.
What are the commercially available forms of packaging?
Naproxen Sodium (Flanax®) 220 mg Film-Coated Tablet - Alu/PVC Blister Pack x 10's (Box of 100's and 500's)
16.
ADR Reporting Statement
For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph.
Seek medical attention immediately at the first sign of any adverse drug reaction.
17.
Registration Number
DR-XY46695
18.
Date of First Authorization
11 September 2019
19.
Date of Revision of Patient Information Leaflet

October 2022
20.
Name and Address of Manufacturer, Marketing Authorization Holder, and Distributor
Manufactured by
Interphil Laboratories, Inc.
Canlubang Industrial Estate, Bo. Pittland,
Cabuyao, Laguna
For
Taisho Pharmaceuticals (Philippines), Inc.
8/F Clipp Center Bldg., 11th Ave. cor. 39th St.,
Bonifacio Global City, Taguig City
TAISHO
Distributed by
Zuellig Pharma Corp.
Km. 14 West Service Road Superhighway
Corner Edison Avenue, Brgy. Sun Valley, Parañaque City
11/2022
DIECUT LEGEND
PLEASE Check proof carefully
for errors and omissions.
Cut
Approval on this artwork
Perforation
DIECUT
Scoring
GLUE AREA
constitutes acceptance of full
COLORS
responsibility for all errors,
omissions and legal and ethical
BLACK
compliance in this document.
GOODREAM
PRINTERS
CORP. will not accept liability
for errors overlooked at this
at Size:
stage of proofing. Any changes
from your previously approved
copy will be charged extra
OF No.
according to both time and
THE DISCLOSURE AND CONTENTS OF THIS DRAWING ARE CONFIDENTIAL AND
materials.

barcode

Naproxen Sodium
Flanax® 220 mg

TAISHO

Non-Steroidal Anti-inflammatory Drug

1. Description of the Product
Naproxen Sodium (Flanax®) is a blue, opaque film-coated, modified oval-shaped tablet.

2. What is in the Medicine?
This product contains Naproxen Sodium which belongs to a group of medicines called non-steroidal anti-inflammatory drugs (also known as NSAIDs). Like other NSAIDs, Naproxen Sodium works by changing the body's chemical response to pain, swelling and fever resulting in relief of symptoms of inflammation (e.g., swelling, redness) and relief of pain and/or fever.

3. Strength of the Medicine
One tablet contains 220 mg of the active ingredient, Naproxen sodium (or the sodium salt of naproxen) – made from a combination of Naproxen 200 mg and 20 mg sodium (of low content, equating to a slice of wheat bread).

4. What is this Medicine used for?
Naproxen sodium (Flanax®) can be used to relieve headache, toothache, muscular ache and backache, pain of menstrual cramps (i.e., period pain), minor pain of arthritis, minor aches and pains associated with the common cold, and fever.

5. How much and how often should you use this Medicine?
Each dose should be taken with water on full stomach after eating. Adults: 1 tablet every 12 hours while symptoms persist. Elderly (>65): 1 tablet per day and no more than 2 tablets per day. Take Naproxen only if with pain and stop when pain is gone.
Do not give this drug to children under 12 years old, except under the advice and supervision of a doctor.

6. When should you not take this Medicine?
Stomach and intestinal pain should not be treated with Naproxen Sodium (Flanax®).
Naproxen Sodium (Flanax®) should not be taken by patients with: a) history of disease of the stomach or of the intestines (ulcer), b) blood coagulation disorders or are receiving drug therapy that interferes with blood coagulation; c) kidney impairment; d) allergy to Naproxen; e) allergy to aspirin or other anti-inflammatory and pain relieving drugs, f) history of Cardiovascular Arrest, Myocardial Infarction, Coronary Artery Bypass Graft, uncontrolled hypertension, Congestive Heart Failure II-IV. Naproxen Sodium (Flanax®) can cause allergic reactions in patients allergic to aspirin and to other anti-inflammatory or pain relieving drugs.

7. Care that should be taken when taking this Medicine?
A doctor should be consulted if the pain persists or gets worse. A doctor or a pharmacist should be consulted if: a) any other drugs are taken on a regular basis; b) heartburn, upset stomach or stomach pain occurs; c) even mild symptoms persist; d) you suffer from another illness or any unusual symptoms occur, e) you are allergic.
Naproxen Sodium (Flanax®) should not be taken for more than 10 consecutive days, unless directed by a doctor.
Do not exceed the indicated dosage and immediately consult a doctor in the case of an overdose.

8. Undesirable Effects of this Medicine
This medicine can cause varying degrees of inconvenient effects in certain people. The most frequent adverse events are – abdominal discomfort, stomach discomfort and heartburn, headache, nausea, vomiting, peripheral edema, allergy, ringing in the ears, skin rash, urticaria, pruritus, and dizziness.
In case of unusual reactions, stop treatment immediately and consult your doctor.

9. What other medicine or food should be avoided while using this Medicine?
- When taking Naproxen, ask your doctor before using other medications for fever, swelling or flu-like symptoms because they may contain chemicals similar to Naproxen (such as Ibuprofen, Aspirin, Diclofenac, etc)
- Avoid taking other NSAIDs such as Aspirin, Ibuprofen, Diclofenac or Ketoprofen as these may increase the risk for gastrointestinal bleeding if taken together with Naproxen
- Avoid taking alcohol since this can also increase the risk for gastrointestinal bleeding
- Co-administration with anticoagulants or blood thinning agents such as warfarin or dicumarol or antiplatelet drugs may increase the risk of gastrointestinal bleeding
- For patients on corticosteroids, taking Naproxen or any other NSAID increases the chance of gastrointestinal bleeding. This may be due to steroids reducing gastric mucus production.
- Naproxen may reduce or blunt the effect of some antihypertensive drugs such as ACE Inhibitors (Captopril, Enalapril, Lisinopril, etc.) and beta-blockers (Propranolol, Carvedilol, Metroprolol, etc.) and Angiotensin Receptor Blockers (Losartan, Telmisartan)
- Naproxen may reduce the effect of some diuretics like thiazides and furosemide
- Taking Naproxen with Lithium may increase the levels of Lithium in the body which may increase the risk of toxicity
- Taking Cholestyramine with Naproxen may affect the absorption of Naproxen
- Taking Methotrexate with Naproxen may increase the risk for toxicity for methotrexate
- Combining antidepressants such as selective serotonin reuptake inhibitors (SSRI) like Fluoxetine, Citalopram, Paroxetine, etc with Naproxen can increase the risk of GIT bleeding

10. What should you do if you miss a dose?
If you skip or miss a dose, take the next dose particularly if there is still pain and inflammation and continue with your regular dosing schedule of every 8 to 12 hours. Do not take a double dose to make up for a missed one.

11. Signs and Symptoms of Overdose
Drowsiness, heart burn, indigestion, nausea or vomiting

12. What to do when you have taken more than the recommended dosage?
Should a patient ingest more than a large quantity of Naproxen Sodium (Flanax) accidentally or purposefully, **seek medical attention immediately**. The stomach may be emptied, and usual supportive measures employed.

13. How should you keep this Medicine?
Naproxen Sodium (Flanax®) should be kept out of reach of children.
Store at temperatures not exceeding 30°C.
Do not use Naproxen Sodium (Flanax®) after the expiry date shown on the carton or blister strip.

14. When should you consult your doctor?
For further information, consult your doctor or pharmacist who has detailed scientific information.

15. What are the commercially available forms of packaging?
Naproxen Sodium (Flanax®) 220 mg Film-Coated Tablet - Alu/PVC Blister Pack x 10's (Box of 100's and 500's)

16. ADR Reporting Statement
For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph.
Seek medical attention immediately at the first sign of any adverse drug reaction.

17. Registration Number
DR-XY46695

18. Date of First Authorization
11 September 2019

19. Date of Revision of Patient Information Leaflet
October 2022

20. Name and Address of Manufacturer, Marketing Authorization Holder, and Distributor

TAISHO

Manufactured by
Interphil Laboratories, Inc.
Canlubang Industrial Estate, Bo. Pittland,
Cabuyao, Laguna
For
Taisho Pharmaceuticals (Philippines), Inc.
8/F Clipp Center Bldg., 11th Ave. cor. 39th St.,
Bonifacio Global City, Taguig City
Distributed by
Zuellig Pharma Corp.
Km. 14 West Service Road Superhighway
Corner Edison Avenue, Brgy. Sun Valley, Parañaque City

1
11/2022

Client: **TAISHO**	**ARTWORK SPECIFICATION**	**DIECUT LEGEND**
Agent: _____	Job Description: **Flanax 220mg Insert**	DIECUT; Cut; Perforation; Scoring; GLUE AREA
Artist: **JOEY**	File Name: **TAISHO_Flanax 220mg Insert_120622**	**COLORS**
ARTWORK DEVELOPMENT	Others:.....	BLACK
Revision Number:	Item Code: _____	
Revised/Prepared By:	Date Started: **Nov. 02, 2022** / Revision Date:	
Revisions Summary:	Spread Size: **139.7mm x 215.9mm** / Folded Size/ Dimension: / No. of Outs: / Flat Size:	
	GRP GOODREAM PRINTERS CORPORATION **CREATIVES** / CJOF No. **XXXX**	

THE DISCLOSURE AND CONTENTS OF THIS DRAWING ARE CONFIDENTIAL AND MAY NOT BE COPIED OR REPRODUCED IN ANY MANNER WITHOUT PERMISSION .

PLEASE Check proof carefully for errors and omissions. Approval on this artwork constitutes acceptance of full responsibility for all errors, omissions and legal and ethical compliance in this document. **GOODREAM PRINTERS CORP.** will not accept liability for errors overlooked at this stage of proofing. Any changes from your previously approved copy will be charged extra according to both time and materials.

Flanax (Naproxen Sodium)

275 mg Film-Coated Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY46696

Flanax: Official FDA Philippines registration record

Brand name	Flanax
Generic name	Naproxen Sodium
Dosage strength and form	275 mg Film-Coated Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Non-Steroidal Anti-Inflammatory Drug
Registration number	DR-XY46696
Manufacturer	PT. Taisho Pharmaceutical Indonesia Tbk.

Flanax: OTC forms registered under the Flanax brand (capsule availability)

OTC capsule form registered under the Flanax brand: NO. 7 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
FlanaxHot	102 mg/ 54.35 mg/ 13.65 mg per gram Topical Cream	Household Remedy (HR)	DRHR-2228
FlanaxACE HOT	2 g/1g/800 mg / 13 mg per 100 g Transdermal Patch	Household Remedy (HR)	DRHR-2413
Flanax	220 mg Film-Coated Tablet	Over-the-Counter (OTC) Drug	DR-XY46695
Flanax	275 mg Film-Coated Tablet	Over-the-Counter (OTC) Drug	DR-XY46696
FlanaxACE	4.98/28/1.48/900 mg per 100 g Transdermal Patch	Household Remedy (HR)	DRHR-2412
FlanaxCool	40 mg/g (4% w/w) Topical Gel	Household Remedy (HR)	DRHR-2227
FlanaxPlus	5 mg/ 102 mg/ 54.3 mg/ 13.7 mg per gram Topical Gel	Over-the-Counter (OTC) Drug	DR-XY48481

Flanax: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY46696 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Flanax"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

FlanaxACE (Menthol + Tocopherol Acetate (Vitamin E) + Glycol Salicylate + dl-Camphor)

4.98/28/1.48/900 mg per 100 g Transdermal Patch | Household Remedy (HR) | FDA Reg. No. DRHR-2412

FlanaxACE: Official FDA Philippines registration record

Brand name	FlanaxACE
Generic name	Menthol + Tocopherol Acetate (Vitamin E) + Glycol Salicylate + dl-Camphor
Dosage strength and form	4.98/28/1.48/900 mg per 100 g Transdermal Patch
Classification	Household Remedy (HR)
Pharmacologic category	Topical Analgesic
Registration number	DRHR-2412
Manufacturer	Tokuhon Corporation (Miyashiro Factory)

FlanaxACE: OTC forms registered under the Flanax brand (capsule availability)

OTC capsule form registered under the Flanax brand: NO. 7 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
FlanaxHot	102 mg/ 54.35 mg/ 13.65 mg per gram Topical Cream	Household Remedy (HR)	DRHR-2228
FlanaxACE HOT	2 g/1g/800 mg / 13 mg per 100 g Transdermal Patch	Household Remedy (HR)	DRHR-2413
Flanax	220 mg Film-Coated Tablet	Over-the-Counter (OTC) Drug	DR-XY46695
Flanax	275 mg Film-Coated Tablet	Over-the-Counter (OTC) Drug	DR-XY46696
FlanaxACE	4.98/28/1.48/900 mg per 100 g Transdermal Patch	Household Remedy (HR)	DRHR-2412
FlanaxCool	40 mg/g (4% w/w) Topical Gel	Household Remedy (HR)	DRHR-2227
FlanaxPlus	5 mg/ 102 mg/ 54.3 mg/ 13.7 mg per gram Topical Gel	Over-the-Counter (OTC) Drug	DR-XY48481

FlanaxACE: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-2412 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Flanax"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Katinko (Isopropyl Alcohol)

70% (v/v) Solution | Household Remedy (HR) | FDA Reg. No. DRHR-2243

Katinko: Official FDA Philippines registration record

Brand name	Katinko
Generic name	Isopropyl Alcohol
Dosage strength and form	70% (v/v) Solution
Classification	Household Remedy (HR)
Pharmacologic category	Antiseptic/Disinfectant
Registration number	DRHR-2243
Manufacturer	Greenstone Pharmaceutical, Inc.

Katinko: OTC forms registered under the Katinko brand (capsule availability)

OTC capsule form registered under the Katinko brand: NO. 48 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Katinko Lavender Massage Oil	1 g/1 g per 100 mL Oil	Household Remedy (HR)	DRHR-2528
Katinko Massage Oil	1 g/1 g per 100 mL Oil	Household Remedy (HR)	DRHR-2529
Katinko Calming	10 mg / 5 mg per mL Liniment (Lavender Scent)	Over-the-Counter (OTC) Drug	DRHR-2350
Katinko Peppermint	10.5g/1g/0.5g per 100g Ointment Stick	Household Remedy (HR)	DRHR-2235
Katinko Peppermint Oil	11.5 g/2 g/2 g/1 g per mL Liniment	Household Remedy (HR)	DRHR-2233
Katinko	110mg/100mg/50mg per gram Ointment (Ginger & Patchouli Scent)	Household Remedy (HR)	DRHR-2536
Katinko	140 mg / 70 mg 10 mg per g Gel	Household Remedy (HR)	DRHR-2364
Katinko	140 mg/40 mg per 2 mL Inhaler	Household Remedy (HR)	DRHR-1955
Katinko Ginger Massage	2 g/2 g per 100 mL Oil	Household Remedy (HR)	DRHR-2526
Katinko Calming	2 g/2 g/2 g per 100 g Ointment (Lavender Scent)	Household Remedy (HR)	DRHR-2209
Katinko	2 g/2 g/2 g per 100 g Ointment Stick (Ginger and Patchouli Scent)	Household Remedy (HR)	DRHR-2244
Katinko	2 g/2 g/2 g per 100 mL Liniment (Ginger and Patchouli Oil Scent)	Household Remedy (HR)	DRHR-2216
Katinko Peppermint Massage Oil	2 g/500 mg per 100 mL Oil	Household Remedy (HR)	DRHR-2530
Katinko (Ginger & Patchouli Oil Scent)	2.0g / 2.0g /2.0g per 100 mL Liniment	Household Remedy (HR)	DRHR-2148

Brand name	Dosage strength and form	Classification	Reg. No.
Katinko	20 mg/110 mg/50 mg per gram Ointment	Household Remedy (HR)	DRHR-2525
Katinko White Extra Strength	200 mg/100 mg/50 mg/30 mg per gram Ointment	Household Remedy (HR)	DRHR-2466
Katinko Peppermint	200 mg/150 mg per gram Ointment	Household Remedy (HR)	DRHR-2547
Katinko Sports Max Relief	200 mg/200 mg per 1 g Cream	Household Remedy (HR)	DRHR-3464
Katinko White Extra Strength	200mg/100mg/50mg/30 mg per g ointment	Household Remedy (HR)	DRHR-2474
Katinko (Ginger & Patchouli Scent)	20mg 20mg / 20mg per 1g Ointment	Household Remedy (HR)	DRHR-2321
Katinko Peppermint	250 mg/250 mg/100 mg per mL Nasal Inhaler	Household Remedy (HR)	DRHR-2475
Katinko Calming	250 mg/250 mg/100 mg/mL per Inhaler Inhaler	Household Remedy (HR)	DRHR-2453
Katinko Fragrant Oil	27 mg/ 25 mg/ 6 mg per 100 mL Liniment Oil	Household Remedy (HR)	DRHR-1516
Katinko Fragrant Oil	27 mg/ 25 mg/ 6 mg per 100 mL Liniment Oil	Household Remedy (HR)	HRP-724
Katinko Calming	2g/2g/2g per 100g Ointment Stick	Household Remedy (HR)	DRHR-2208
Katinko Oil	30 mg/15 mg/15 mg/12 mg per 100 mL Liniment	Household Remedy (HR)	DRHR-2037
Katinko Oil	30 mg/15 mg/15 mg/12 mg per 100 mL Liniment	Household Remedy (HR)	HRP-723
Katinko Oil	30 mg/15 mg/15 mg/12 mg per 100 mL Liniment Oil	Household Remedy (HR)	DRHR-1519
Katinko Gingery	302.5 mg/302.5 mg per mL Inhaler	Household Remedy (HR)	DRHR-2486
Katinko Ginger & Patchouli	302.5 mg/302.5 mg/20 mg per mL Liniment/Inhaler (Lemongrass/Ginger/Patchouli Scent)	Household Remedy (HR)	DRHR-2395
KATINKO	42.7 mg per 100 mg Ointment Stick	Household Remedy (HR)	DRHR-1935
Katinko	42.7 mg per 100 mg Ointment Stick	Household Remedy (HR)	HRP-722
Katinko Double	460mg/460mg per mL Inhaler	Household Remedy (HR)	DRHR-2548
Katinko Calming	50mg/50mg/50mg per gram Ointment (Lavender Scent)	Household Remedy (HR)	DRHR-2537
Katinko Sulfura	60 mg/40 mg/25 mg per gram Topical Cream	Household Remedy (HR)	DRHR-2550
Katinko Alco	70 mL/100 mL (70% v/v) Solution	Household Remedy (HR)	DRHR-2338

Brand name	Dosage strength and form	Classification	Reg. No.
Katinko	70% (v/v) Solution	Household Remedy (HR)	DRHR-2243
Katinko Sports	8 mg 7.5 mg/2.6 mg Spray	Household Remedy (HR)	DRHR-2300
Katinko Soft White	80 mg/75 mg/31 mg/26 mg per gram Ointment	Household Remedy	DRHR-1725
Katinko White	80 mg/75 mg/31 mg/26 mg per gram Ointment	Household Remedy (HR)	HRP-655
Katinko	80mcg/75mcg/26mcg per gram Ointment	Household Remedy (HR)	DRHR-1580
Katinko	8mg/7.5mg/2.6mg per 100mg Ointment Stick	Household Remedy (HR)	DRHR-1151
Katinko	9.6 mg/7.5 mg/2.6 mg per 100 mL Oil Liniment	Household Remedy (HR)	DRHR-1326
Katinko	Formulation (each mL contains): Inhaler (1 mL): 250 mg Camphor Eucalyptus Oil (as 1-8, Cineole) 250 mg Peppermint Oil (as Menthol) 100 mg Liniment (1 mL): Peppermint Oil (as Menthol) 11.5 g Eucalyptus Oil (as 1-8, Cineole) 28 2 g Methyl Salicylate 1 g Camphor Inhaler/Liniment (Peppermint)	Household Remedy (HR)	DRHR-2463
Katinko	Formulation (each mL contains): Inhaler (1 mL): Camphor Menthol Lavender Oil (as Linalool) 250 mg 250 mg 100 mg Liniment (1 mL): Methyl Salicylate 200 mg Lavender Oil (as Linalool) 200 mg 50 mg Camphor 50 mg Menthol Inhaler/Liniment (Lavender)	Household Remedy (HR)	DRHR-2462
Katinko Peppermint	Formulation: Each 100 g contains: Peppermint Oil 10.5 g Methyl Salicylate 1 g Camphor 0.5 g Ointment	Household Remedy (HR)	DRHR-2234
Katinko Peppermint Oil	Formulation: Each 100 mL contains: Peppermint Oil 11.5 g Eucalyptus Oil 2 g Methyl Salicylate 2 g Camphor 1 g Liniment (Ginger and Patchouli Oil Scent)	Household Remedy (HR)	DRHR-2236
Katinko Duo Relief	Formulation; Liniment: Each mL contains Methyl Salicylate 300 mg Menthol 150 mg Eucalyptus Oil 100 mg Camphor 100 mg Inhaler: Each mL contains Methyl Salicylate 450 mg Menthol 450 mg Liniment/Inhaler	Household Remedy (HR)	DRHR-2361

Katinko: Official leaflet (Package Insert / Patient Information Leaflet)

An official leaflet for this registration is published on the FDA Verification Portal: PIL_01 (https://drive.google.com/file/d/1skS0LJYvaGrC4VKSUpx6Hp2gIXyKA2aA/view?usp=drivesdk). Its text (OCR) and the original leaflet pages follow.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Katinko"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Katinko: Official leaflet text (OCR transcription)


Machine transcription of the official FDA leaflet for Reg. No. DRHR-2243. OCR may contain errors; the original leaflet pages are attached after this section.

Katinko leaflet PIL_01

PROPOSED CHANGE
500 mL
DRHR-2243
ISOPROPYL
ALCOHOL
Alco
KATINKO
70% (v/v) Solution
with
Moisturizer
ANTISEPTIC/DISINFECTANT
PROUDLY MADE IN THE PHILIPPINES
Proposed Brand Name: Alco Katinko
-Yellow, Green, and Red Background
*Philippine Packaging
BN:
MD:
ED:
Manufactured by:
GREENSTONE PHARMACEUTICAL, INC. Anabu Hills Industrial Estate,
Anabu 1-C, Imus, Cavite, Philippines 4103
Contact Us:
(+632) 8922-4794 / (+632) 3433-6014
(+63917) KATINKO / (+63917) 528-4656
sales@katinko.com
care@katinko.com
feedback.katinko@gmail.com
CUSTOMER CARE
KATINKO
Scan this
Patient
Information
Leaflet (PIL)
FORMULATION
Each 100 mL contains:
(Isopropyl Alcohol, USP)
INDICATION: For disinfection, sterilization, relief of prickly heat, insect bites, hygienic purposes, after shave, and general antiseptic. For use in the hospital or sick room as a rubdown or for general massage purposes.
DIRECTION: Apply on hands and rub on skin.
PRECAUTION:Can cause skin rash, itching, dryness and redness.
WARNING: FOR EXTERNAL USE ONLY. Flammable. Keep out of reach of children. Do not swallow or use near eyes, ears, and mouth. For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph. Seek medical attention immediately at the first sign of any adverse drug reaction.
STORAGE CONDITION: Store at temperatures not exceeding 30°C. Do not freeze.
70 mL
250 mL
DRHR-2243
ISOPROPYL
ALCOHOL
Alco
KATINKO

70% (v/v) Solution
with
Moisturizer
ANTISEPTIC/DISINFECTANT
PROUDLY MADE IN THE PHILIPPINES
*Philippine Packaging
Manufactured by:
GREENSTONE PHARMACEUTICAL, INC. Anabu Hills Industrial Estate,
Anabu 1-C, Imus, Cavite,
Philippines 4103
BN:
MD:
ED:
Contact Us:
(+632) 8922-4794 / (+632) 3433-6014
(+63917) KATINKO / (+63917) 528-4656
sales@katinko.com
care@katinko.com
feedback.katinko@gmail.com
CUSTOMER CARE Scan this for
Patient
Informatio
Leaflet (PIL)
KATINKO
FORMULATION
Each 100 mL contains:
(Isopropyl Alcohol, USP)
INDICATION: For disinfection, sterilization, relief of prickly heat, insect bites, hygienic purposes, after shave, and general antiseptic. For use in the hospital or sick room as a rubdown or for general massage purposes.
DIRECTION: Apply on hands and rub on skin.
PRECAUTION:Can cause skin rash, itching, dryness and redness.
WARNING:FOR EXTERNAL USE ONLY.Flammable. Keep out of reach of children. Do not swallow or use near eyes, ears, and mouth. For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph. Seek medical attention immediately at the first sign of any adverse drug reaction.
STORAGE CONDITION: Store at temperatures not exceeding 30°C. Do not freeze.
70 mL

PROPOSED CHANGE



Proposed Brand Name: Alco Katinko
-Yellow, Green, and Red Background

FORMULATION

Each 100 mL contains:

(Isopropyl Alcohol, USP) 70 mL

INDICATION: For disinfection, sterilization, relief of prickly heat, insect bites, hygienic purposes, after shave, and general antiseptic. For use in the hospital or sick room as a rubdown or for general massage purposes.

DIRECTION: Apply on hands and rub on skin.

PRECAUTION: Can cause skin rash, itching, dryness and redness.

WARNING: FOR EXTERNAL USE ONLY. Flammable. Keep out of reach of children. Do not swallow or use near eyes, ears, and mouth. For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph. Seek medical attention immediately at the first sign of any adverse drug reaction.

STORAGE CONDITION: Store at temperatures not exceeding 30°C. Do not freeze.

Scan this Patient Information Leaflet (PIL)

CUSTOMER CARE

Contact Us:
(+632) 8922-4794 / (+632) 3433-6014
(+63917) KATINKO / (+63917) 528-4656
sales@katinko.com
care@katinko.com
feedback.katinko@gmail.com

Manufactured by:

GREENSTONE

GREENSTONE PHARMACEUTICAL, INC.
Anabu Hills Industrial Estate,
Anabu 1-C, Imus, Cavite,
Philippines 4103

BN:
MD:
ED:

*Philippine Packaging

4 806038 000557

250 mL

DRHR-2243

ISOPROPYL ALCOHOL

Alco KATINKO®

70% (v/v) Solution

with Moisturizer

ANTISEPTIC/DISINFECTANT

PROUDLY MADE IN THE PHILIPPINES

FORMULATION

Each 100 mL contains:

(Isopropyl Alcohol, USP) 70 mL

INDICATION: For disinfection, sterilization, relief of prickly heat, insect bites, hygienic purposes, after shave, and general antiseptic. For use in the hospital or sick room as a rubdown or for general massage purposes.

DIRECTION: Apply on hands and rub on skin.

PRECAUTION: Can cause skin rash, itching, dryness and redness.

WARNING: FOR EXTERNAL USE ONLY. Flammable. Keep out of reach of children. Do not swallow or use near eyes, ears, and mouth. For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph. Seek medical attention immediately at the first sign of any adverse drug reaction.

STORAGE CONDITION: Store at temperatures not exceeding 30°C. Do not freeze.

Scan this for Patient Informatio Leaflet (PIL)

CUSTOMER CARE

Contact Us:
(+632) 8922-4794 / (+632) 3433-6014
(+63917) KATINKO / (+63917) 528-4656
sales@katinko.com
care@katinko.com
feedback.katinko@gmail.com

Manufactured by:

GREENSTONE PHARMACEUTICAL, INC.
Anabu Hills Industrial Estate,
Anabu 1-C, Imus, Cavite,
Philippines 4103

BN:
MD:
ED:

*Philippine Packaging

4 806038 000540

KATINKO (Menthol)

42.7 mg per 100 mg Ointment Stick | Household Remedy (HR) | FDA Reg. No. DRHR-1935

KATINKO: Official FDA Philippines registration record

Brand name	KATINKO
Generic name	Menthol
Dosage strength and form	42.7 mg per 100 mg Ointment Stick
Classification	Household Remedy (HR)
Pharmacologic category	Counterirritant
Registration number	DRHR-1935
Manufacturer	Greenstone Pharmaceutical, Inc.

KATINKO: OTC forms registered under the Katinko brand (capsule availability)

OTC capsule form registered under the Katinko brand: NO. 48 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Katinko Lavender Massage Oil	1 g/1 g per 100 mL Oil	Household Remedy (HR)	DRHR-2528
Katinko Massage Oil	1 g/1 g per 100 mL Oil	Household Remedy (HR)	DRHR-2529
Katinko Calming	10 mg / 5 mg per mL Liniment (Lavender Scent)	Over-the-Counter (OTC) Drug	DRHR-2350
Katinko Peppermint	10.5g/1g/0.5g per 100g Ointment Stick	Household Remedy (HR)	DRHR-2235
Katinko Peppermint Oil	11.5 g/2 g/2 g/1 g per mL Liniment	Household Remedy (HR)	DRHR-2233
Katinko	110mg/100mg/50mg per gram Ointment (Ginger & Patchouli Scent)	Household Remedy (HR)	DRHR-2536
Katinko	140 mg / 70 mg 10 mg per g Gel	Household Remedy (HR)	DRHR-2364
Katinko	140 mg/40 mg per 2 mL Inhaler	Household Remedy (HR)	DRHR-1955
Katinko Ginger Massage	2 g/2 g per 100 mL Oil	Household Remedy (HR)	DRHR-2526
Katinko Calming	2 g/2 g/2 g per 100 g Ointment (Lavender Scent)	Household Remedy (HR)	DRHR-2209
Katinko	2 g/2 g/2 g per 100 g Ointment Stick (Ginger and Patchouli Scent)	Household Remedy (HR)	DRHR-2244
Katinko	2 g/2 g/2 g per 100 mL Liniment (Ginger and Patchouli Oil Scent)	Household Remedy (HR)	DRHR-2216
Katinko Peppermint Massage Oil	2 g/500 mg per 100 mL Oil	Household Remedy (HR)	DRHR-2530
Katinko (Ginger & Patchouli Oil Scent)	2.0g / 2.0g /2.0g per 100 mL Liniment	Household Remedy (HR)	DRHR-2148

Brand name	Dosage strength and form	Classification	Reg. No.
Katinko	20 mg/110 mg/50 mg per gram Ointment	Household Remedy (HR)	DRHR-2525
Katinko White Extra Strength	200 mg/100 mg/50 mg/30 mg per gram Ointment	Household Remedy (HR)	DRHR-2466
Katinko Peppermint	200 mg/150 mg per gram Ointment	Household Remedy (HR)	DRHR-2547
Katinko Sports Max Relief	200 mg/200 mg per 1 g Cream	Household Remedy (HR)	DRHR-3464
Katinko White Extra Strength	200mg/100mg/50mg/30 mg per g ointment	Household Remedy (HR)	DRHR-2474
Katinko (Ginger & Patchouli Scent)	20mg 20mg / 20mg per 1g Ointment	Household Remedy (HR)	DRHR-2321
Katinko Peppermint	250 mg/250 mg/100 mg per mL Nasal Inhaler	Household Remedy (HR)	DRHR-2475
Katinko Calming	250 mg/250 mg/100 mg/mL per Inhaler Inhaler	Household Remedy (HR)	DRHR-2453
Katinko Fragrant Oil	27 mg/ 25 mg/ 6 mg per 100 mL Liniment Oil	Household Remedy (HR)	DRHR-1516
Katinko Fragrant Oil	27 mg/ 25 mg/ 6 mg per 100 mL Liniment Oil	Household Remedy (HR)	HRP-724
Katinko Calming	2g/2g/2g per 100g Ointment Stick	Household Remedy (HR)	DRHR-2208
Katinko Oil	30 mg/15 mg/15 mg/12 mg per 100 mL Liniment	Household Remedy (HR)	DRHR-2037
Katinko Oil	30 mg/15 mg/15 mg/12 mg per 100 mL Liniment	Household Remedy (HR)	HRP-723
Katinko Oil	30 mg/15 mg/15 mg/12 mg per 100 mL Liniment Oil	Household Remedy (HR)	DRHR-1519
Katinko Gingery	302.5 mg/302.5 mg per mL Inhaler	Household Remedy (HR)	DRHR-2486
Katinko Ginger & Patchouli	302.5 mg/302.5 mg/20 mg per mL Liniment/Inhaler (Lemongrass/Ginger/Patchouli Scent)	Household Remedy (HR)	DRHR-2395
KATINKO	42.7 mg per 100 mg Ointment Stick	Household Remedy (HR)	DRHR-1935
Katinko	42.7 mg per 100 mg Ointment Stick	Household Remedy (HR)	HRP-722
Katinko Double	460mg/460mg per mL Inhaler	Household Remedy (HR)	DRHR-2548
Katinko Calming	50mg/50mg/50mg per gram Ointment (Lavender Scent)	Household Remedy (HR)	DRHR-2537
Katinko Sulfura	60 mg/40 mg/25 mg per gram Topical Cream	Household Remedy (HR)	DRHR-2550
Katinko Alco	70 mL/100 mL (70% v/v) Solution	Household Remedy (HR)	DRHR-2338

Brand name	Dosage strength and form	Classification	Reg. No.
Katinko	70% (v/v) Solution	Household Remedy (HR)	DRHR-2243
Katinko Sports	8 mg 7.5 mg/2.6 mg Spray	Household Remedy (HR)	DRHR-2300
Katinko Soft White	80 mg/75 mg/31 mg/26 mg per gram Ointment	Household Remedy	DRHR-1725
Katinko White	80 mg/75 mg/31 mg/26 mg per gram Ointment	Household Remedy (HR)	HRP-655
Katinko	80mcg/75mcg/26mcg per gram Ointment	Household Remedy (HR)	DRHR-1580
Katinko	8mg/7.5mg/2.6mg per 100mg Ointment Stick	Household Remedy (HR)	DRHR-1151
Katinko	9.6 mg/7.5 mg/2.6 mg per 100 mL Oil Liniment	Household Remedy (HR)	DRHR-1326
Katinko	Formulation (each mL contains): Inhaler (1 mL): 250 mg Camphor Eucalyptus Oil (as 1-8, Cineole) 250 mg Peppermint Oil (as Menthol) 100 mg Liniment (1 mL): Peppermint Oil (as Menthol) 11.5 g Eucalyptus Oil (as 1-8, Cineole) 28 2 g Methyl Salicylate 1 g Camphor Inhaler/Liniment (Peppermint)	Household Remedy (HR)	DRHR-2463
Katinko	Formulation (each mL contains): Inhaler (1 mL): Camphor Menthol Lavender Oil (as Linalool) 250 mg 250 mg 100 mg Liniment (1 mL): Methyl Salicylate 200 mg Lavender Oil (as Linalool) 200 mg 50 mg Camphor 50 mg Menthol Inhaler/Liniment (Lavender)	Household Remedy (HR)	DRHR-2462
Katinko Peppermint	Formulation: Each 100 g contains: Peppermint Oil 10.5 g Methyl Salicylate 1 g Camphor 0.5 g Ointment	Household Remedy (HR)	DRHR-2234
Katinko Peppermint Oil	Formulation: Each 100 mL contains: Peppermint Oil 11.5 g Eucalyptus Oil 2 g Methyl Salicylate 2 g Camphor 1 g Liniment (Ginger and Patchouli Oil Scent)	Household Remedy (HR)	DRHR-2236
Katinko Duo Relief	Formulation; Liniment: Each mL contains Methyl Salicylate 300 mg Menthol 150 mg Eucalyptus Oil 100 mg Camphor 100 mg Inhaler: Each mL contains Methyl Salicylate 450 mg Menthol 450 mg Liniment/Inhaler	Household Remedy (HR)	DRHR-2361

KATINKO: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-1935 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Katinko"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Katinko (Methyl Salicylate+Menthol+Camphor)

8mg/7.5mg/2.6mg per 100mg Ointment Stick | Household Remedy (HR) | FDA Reg. No. DRHR-1151

Katinko: Official FDA Philippines registration record

Brand name	Katinko
Generic name	Methyl Salicylate+Menthol+Camphor
Dosage strength and form	8mg/7.5mg/2.6mg per 100mg Ointment Stick
Classification	Household Remedy (HR)
Pharmacologic category	Counterirritant
Registration number	DRHR-1151
Manufacturer	Greenstone Pharmaceutical Inc.

Katinko: OTC forms registered under the Katinko brand (capsule availability)

OTC capsule form registered under the Katinko brand: NO. 48 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Katinko Lavender Massage Oil	1 g/1 g per 100 mL Oil	Household Remedy (HR)	DRHR-2528
Katinko Massage Oil	1 g/1 g per 100 mL Oil	Household Remedy (HR)	DRHR-2529
Katinko Calming	10 mg / 5 mg per mL Liniment (Lavender Scent)	Over-the-Counter (OTC) Drug	DRHR-2350
Katinko Peppermint	10.5g/1g/0.5g per 100g Ointment Stick	Household Remedy (HR)	DRHR-2235
Katinko Peppermint Oil	11.5 g/2 g/2 g/1 g per mL Liniment	Household Remedy (HR)	DRHR-2233
Katinko	110mg/100mg/50mg per gram Ointment (Ginger & Patchouli Scent)	Household Remedy (HR)	DRHR-2536
Katinko	140 mg / 70 mg 10 mg per g Gel	Household Remedy (HR)	DRHR-2364
Katinko	140 mg/40 mg per 2 mL Inhaler	Household Remedy (HR)	DRHR-1955
Katinko Ginger Massage	2 g/2 g per 100 mL Oil	Household Remedy (HR)	DRHR-2526
Katinko Calming	2 g/2 g/2 g per 100 g Ointment (Lavender Scent)	Household Remedy (HR)	DRHR-2209
Katinko	2 g/2 g/2 g per 100 g Ointment Stick (Ginger and Patchouli Scent)	Household Remedy (HR)	DRHR-2244
Katinko	2 g/2 g/2 g per 100 mL Liniment (Ginger and Patchouli Oil Scent)	Household Remedy (HR)	DRHR-2216
Katinko Peppermint Massage Oil	2 g/500 mg per 100 mL Oil	Household Remedy (HR)	DRHR-2530
Katinko (Ginger & Patchouli Oil Scent)	2.0g / 2.0g /2.0g per 100 mL Liniment	Household Remedy (HR)	DRHR-2148

Brand name	Dosage strength and form	Classification	Reg. No.
Katinko	20 mg/110 mg/50 mg per gram Ointment	Household Remedy (HR)	DRHR-2525
Katinko White Extra Strength	200 mg/100 mg/50 mg/30 mg per gram Ointment	Household Remedy (HR)	DRHR-2466
Katinko Peppermint	200 mg/150 mg per gram Ointment	Household Remedy (HR)	DRHR-2547
Katinko Sports Max Relief	200 mg/200 mg per 1 g Cream	Household Remedy (HR)	DRHR-3464
Katinko White Extra Strength	200mg/100mg/50mg/30 mg per g ointment	Household Remedy (HR)	DRHR-2474
Katinko (Ginger & Patchouli Scent)	20mg 20mg / 20mg per 1g Ointment	Household Remedy (HR)	DRHR-2321
Katinko Peppermint	250 mg/250 mg/100 mg per mL Nasal Inhaler	Household Remedy (HR)	DRHR-2475
Katinko Calming	250 mg/250 mg/100 mg/mL per Inhaler Inhaler	Household Remedy (HR)	DRHR-2453
Katinko Fragrant Oil	27 mg/ 25 mg/ 6 mg per 100 mL Liniment Oil	Household Remedy (HR)	DRHR-1516
Katinko Fragrant Oil	27 mg/ 25 mg/ 6 mg per 100 mL Liniment Oil	Household Remedy (HR)	HRP-724
Katinko Calming	2g/2g/2g per 100g Ointment Stick	Household Remedy (HR)	DRHR-2208
Katinko Oil	30 mg/15 mg/15 mg/12 mg per 100 mL Liniment	Household Remedy (HR)	DRHR-2037
Katinko Oil	30 mg/15 mg/15 mg/12 mg per 100 mL Liniment	Household Remedy (HR)	HRP-723
Katinko Oil	30 mg/15 mg/15 mg/12 mg per 100 mL Liniment Oil	Household Remedy (HR)	DRHR-1519
Katinko Gingery	302.5 mg/302.5 mg per mL Inhaler	Household Remedy (HR)	DRHR-2486
Katinko Ginger & Patchouli	302.5 mg/302.5 mg/20 mg per mL Liniment/Inhaler (Lemongrass/Ginger/Patchouli Scent)	Household Remedy (HR)	DRHR-2395
KATINKO	42.7 mg per 100 mg Ointment Stick	Household Remedy (HR)	DRHR-1935
Katinko	42.7 mg per 100 mg Ointment Stick	Household Remedy (HR)	HRP-722
Katinko Double	460mg/460mg per mL Inhaler	Household Remedy (HR)	DRHR-2548
Katinko Calming	50mg/50mg/50mg per gram Ointment (Lavender Scent)	Household Remedy (HR)	DRHR-2537
Katinko Sulfura	60 mg/40 mg/25 mg per gram Topical Cream	Household Remedy (HR)	DRHR-2550
Katinko Alco	70 mL/100 mL (70% v/v) Solution	Household Remedy (HR)	DRHR-2338

Brand name	Dosage strength and form	Classification	Reg. No.
Katinko	70% (v/v) Solution	Household Remedy (HR)	DRHR-2243
Katinko Sports	8 mg 7.5 mg/2.6 mg Spray	Household Remedy (HR)	DRHR-2300
Katinko Soft White	80 mg/75 mg/31 mg/26 mg per gram Ointment	Household Remedy	DRHR-1725
Katinko White	80 mg/75 mg/31 mg/26 mg per gram Ointment	Household Remedy (HR)	HRP-655
Katinko	80mcg/75mcg/26mcg per gram Ointment	Household Remedy (HR)	DRHR-1580
Katinko	8mg/7.5mg/2.6mg per 100mg Ointment Stick	Household Remedy (HR)	DRHR-1151
Katinko	9.6 mg/7.5 mg/2.6 mg per 100 mL Oil Liniment	Household Remedy (HR)	DRHR-1326
Katinko	Formulation (each mL contains): Inhaler (1 mL): 250 mg Camphor Eucalyptus Oil (as 1-8, Cineole) 250 mg Peppermint Oil (as Menthol) 100 mg Liniment (1 mL): Peppermint Oil (as Menthol) 11.5 g Eucalyptus Oil (as 1-8, Cineole) 28 2 g Methyl Salicylate 1 g Camphor Inhaler/Liniment (Peppermint)	Household Remedy (HR)	DRHR-2463
Katinko	Formulation (each mL contains): Inhaler (1 mL): Camphor Menthol Lavender Oil (as Linalool) 250 mg 250 mg 100 mg Liniment (1 mL): Methyl Salicylate 200 mg Lavender Oil (as Linalool) 200 mg 50 mg Camphor 50 mg Menthol Inhaler/Liniment (Lavender)	Household Remedy (HR)	DRHR-2462
Katinko Peppermint	Formulation: Each 100 g contains: Peppermint Oil 10.5 g Methyl Salicylate 1 g Camphor 0.5 g Ointment	Household Remedy (HR)	DRHR-2234
Katinko Peppermint Oil	Formulation: Each 100 mL contains: Peppermint Oil 11.5 g Eucalyptus Oil 2 g Methyl Salicylate 2 g Camphor 1 g Liniment (Ginger and Patchouli Oil Scent)	Household Remedy (HR)	DRHR-2236
Katinko Duo Relief	Formulation; Liniment: Each mL contains Methyl Salicylate 300 mg Menthol 150 mg Eucalyptus Oil 100 mg Camphor 100 mg Inhaler: Each mL contains Methyl Salicylate 450 mg Menthol 450 mg Liniment/Inhaler	Household Remedy (HR)	DRHR-2361

Katinko: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-1151 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Katinko"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Medicol (Ibuprofen)

200 mg Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY24958

Medicol: Official FDA Philippines registration record

Brand name	Medicol
Generic name	Ibuprofen
Dosage strength and form	200 mg Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Non-Steroidal Anti-Inflammatory Drug (NSAID)/Analgesic/Antipyretic
Registration number	DR-XY24958
Manufacturer	UNILAB, Inc.

Medicol: OTC forms registered under the Medicol brand (capsule availability)

OTC capsule form registered under the Medicol brand: YES. 6 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Medicol Advance	200 mg Softgel Capsule	Over-the-Counter (OTC)	DR-XY46460
Medicol Advance	200 mg Softgel Capsule	Over-the-Counter (OTC) Drug	DR-XY34641
Medicol	200 mg Tablet	Over-the-Counter (OTC) Drug	DR-XY24958
Medicol Advance 400	400 mg Softgel Capsule	Over-the-Counter (OTC) Drug	DR-XY41760
Medicol Advance 400	400 mg Softgel Capsule	Over-the-Counter Drug (OTC)	DR-XY46461
Medicol I-400	400 mg Tablet	Over-the-Counter (OTC) Drug	DR-XY24952

Medicol: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY24958 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Medicol"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Medicol I-400 (Ibuprofen)

400 mg Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY24952

Medicol I-400: Official FDA Philippines registration record

Brand name	Medicol I-400
Generic name	Ibuprofen
Dosage strength and form	400 mg Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Non-Steroidal Anti-Inflammatory Drug (NSAID)
Registration number	DR-XY24952
Manufacturer	UNILAB, Inc.

Medicol I-400: OTC forms registered under the Medicol brand (capsule availability)

OTC capsule form registered under the Medicol brand: YES. 6 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Medicol Advance	200 mg Softgel Capsule	Over-the-Counter (OTC)	DR-XY46460
Medicol Advance	200 mg Softgel Capsule	Over-the-Counter (OTC) Drug	DR-XY34641
Medicol	200 mg Tablet	Over-the-Counter (OTC) Drug	DR-XY24958
Medicol Advance 400	400 mg Softgel Capsule	Over-the-Counter (OTC) Drug	DR-XY41760
Medicol Advance 400	400 mg Softgel Capsule	Over-the-Counter Drug (OTC)	DR-XY46461
Medicol I-400	400 mg Tablet	Over-the-Counter (OTC) Drug	DR-XY24952

Medicol I-400: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY24952 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Medicol"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Rexidol Forte (Paracetamol + Caffeine)

500 mg/ 65 mg Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY46238

Rexidol Forte: Official FDA Philippines registration record

Brand name	Rexidol Forte
Generic name	Paracetamol + Caffeine
Dosage strength and form	500 mg/ 65 mg Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Analgesic/ Antipyretic
Registration number	DR-XY46238
Manufacturer	PT. Medifarma Laboratories

Rexidol Forte: OTC forms registered under the Rexidol brand (capsule availability)

OTC capsule form registered under the Rexidol brand: NO. 1 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Rexidol Forte	500 mg/ 65 mg Tablet	Over-the-Counter (OTC) Drug	DR-XY46238

Rexidol Forte: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY46238 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Rexidol"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Salonpas (Methyl Salicylate + l-Menthol + Tocopherol Acetate (Vitamin E) + dl-Camphor)

36 mg/ 33 mg/ 12 mg/ 7.1 mg Medicated Patch | Household Remedy (HR) | FDA Reg. No. DRHR-821

Salonpas: Official FDA Philippines registration record

Brand name	Salonpas
Generic name	Methyl Salicylate + l-Menthol + Tocopherol Acetate (Vitamin E) + dl-Camphor
Dosage strength and form	36 mg/ 33 mg/ 12 mg/ 7.1 mg Medicated Patch
Classification	Household Remedy (HR)
Pharmacologic category	Counterirritant/Topical Analgesic
Registration number	DRHR-821
Manufacturer	Hisamitsu Vietnam Pharmaceutical Co., Ltd.

Salonpas: OTC forms registered under the Salonpas brand (capsule availability)

OTC capsule form registered under the Salonpas brand: NO. 2 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Salonpas Jet Spray	100 mg 30 mg Aerosol Spray	Household Remedy (HR)	DRHR-2304
Salonpas	36 mg/ 33 mg/ 12 mg/ 7.1 mg Medicated Patch	Household Remedy (HR)	DRHR-821

Salonpas: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-821 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Salonpas"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Salonpas Jet Spray (Methyl Salicylate + Menthol)

100 mg 30 mg Aerosol Spray | Household Remedy (HR) | FDA Reg. No. DRHR-2304

Salonpas Jet Spray: Official FDA Philippines registration record

Brand name	Salonpas Jet Spray
Generic name	Methyl Salicylate + Menthol
Dosage strength and form	100 mg 30 mg Aerosol Spray
Classification	Household Remedy (HR)
Pharmacologic category	Counterirritant
Registration number	DRHR-2304
Manufacturer	Daizo Corporation (Aerosol Division, Kyoto Plant

Salonpas Jet Spray: OTC forms registered under the Salonpas brand (capsule availability)

OTC capsule form registered under the Salonpas brand: NO. 2 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Salonpas Jet Spray	100 mg 30 mg Aerosol Spray	Household Remedy (HR)	DRHR-2304
Salonpas	36 mg/ 33 mg/ 12 mg/ 7.1 mg Medicated Patch	Household Remedy (HR)	DRHR-821

Salonpas Jet Spray: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-2304 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Salonpas"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Saridon (Paracetamol + Caffeine)

500 mg/50 mg Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY46456

Saridon: Official FDA Philippines registration record

Brand name	Saridon
Generic name	Paracetamol + Caffeine
Dosage strength and form	500 mg/50 mg Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Analgesic/Antipyretic
Registration number	DR-XY46456
Manufacturer	PT. Bayer Indonesia

Saridon: OTC forms registered under the Saridon brand (capsule availability)

OTC capsule form registered under the Saridon brand: NO. 3 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Saridon Triple Action	250 mg/150 mg/50 mg Tablet	Over-the-Counter (OTC) Drug	DR-XY36639
Saridon Max	275 mg Film-Coated Tablet	Over-the-Counter (OTC) Drug	DRP-1872-02
Saridon	500 mg/50 mg Tablet	Over-the-Counter (OTC) Drug	DR-XY46456

Saridon: Official leaflet (Package Insert / Patient Information Leaflet)

An official leaflet for this registration is published on the FDA Verification Portal: PIL_01 (https://drive.google.com/file/d/1SQ35eXDR6ZeNwH3q9pHJkiqPn40ax1m7/view?usp=drivesdk). Its text (OCR) and the original leaflet pages follow.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Saridon"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Saridon: Official leaflet text (OCR transcription)

Machine transcription of the official FDA leaflet for Reg. No. DR-XY46456. OCR may contain errors; the original leaflet pages are attached after this section.

Saridon leaflet PIL_01

NNNNNN
500 mg/50 mg Tablet
Analgesic/Antipyretic
1. Name of the Product
Paracetamol + Caffeine (Saridon®)
2. Description of the Product
The product contains paracetamol and caffeine in tablet form which is classi
ed as antipyretic and analgesic.
3. What is in the Medicine?
Each tablet contains:
Paracetamol…….. 500 mg
Caffeine……. 50 mg
4. Strength of the Medicine
Paracetamol + Caffeine (Saridon®) 500 mg/50 mg Tablet
5. What is the Medicine used for?
• For fast and effective relief of mild to severe headaches.
• For the relief of pain such as headache, toothache, menstrual discomfort, postoperative and rheumatic pain.
• For the relief of pain and fever associated with colds and “
u”.
6. How much and how often should you use this Medicine?
Single dose.
Adults: 1-2 tablets
Adolescents aged 12-16 years: 1 tablet
If necessary, up to three single doses may be taken within 24 hours.
The tablets should be taken with ample water or other
uid.
It should not be taken for more than one week or at doses higher than those recommended, except as directed by a doctor.
Patients with hepatic impairment
Paracetamol is contraindicated in patients with severe hepatic insu
ciency (Child-Pugh>9). Paracetamol should be used with caution in
patients with mild to moderate hepatic insu
ciency (Child-Pugh <9).
Patients with renal impairment
Paracetamol should be used with particular caution in patients with severe renal failure. Dose adjustment may be necessary in patients
with impaired renal function.
7. When should you not take this Medicine?
It should not be used if any of the following conditions is present:
• Allergy to phenylbutazone-containing products,
• Known hypersensitivity to paracetamol, acetylsalicylic acid or proven allergy to caffeine,
• Inherited de
ciency of glucose-6-phosphate dehydrogenase (evidenced by hemolytic anemia),
• Acute hepatic porphyria
Special caution should be taken (lowering of the dosage and/or spacing out of the doses) in the following conditions:
• Impaired hepatic function (in cases of chronic alcohol abuse or hepatitis for example),
• Impaired renal function,
• Gilbert’s syndrome (
uctuating benign jaundice secondary to glucuronyl-transferase de
ciency),
• Hematopoetic dysfunction
It should not be given to infants or children under 12 years of age.
It is recommended that it should not be used during pregnancy, particularly in the
rst trimester and in the
nal six weeks.
It should not be used by nursing women.
8. Care that should be taken when taking this Medicine?
• Should not be taken regularly for a prolonged period, except as directed by a doctor.
• Prolonged intake of paracetamol-containing analgesics resulting in high cumulative doses may in rare instances
lead to analgesic nephropathy and largely irreversible renal insu
ciency.

• Very rare cases of serious skin reactions have been reported. In the event of skin reddening, rash, blisters or peeling, discontinue paracetamol use and seek medical attention immediately.
• Continual use of analgesics for the treatment of headache may itself lead to chronic headaches.
• Special caution for patients with asthma, chronic rhinitis or chronic urticaria, particularly those hypersensitive to other anti-in
ammatory agents.
• There have been isolated reports of asthma attacks and anaphylactic reaction associated with paracetamolcontaining preparations in susceptible individuals.
• Should not be used after the expiry date shown in the pack.
• Pregnancy and Lactation: Category C.
º Available data from epidemiological studies on the therapeutic use of paracetamol and possible undesirable effects regarding pregnancy or fetal/neonatal development continue to support a positive bene
t/risk balance. In reproduction studies of oral
paracetamol, no evidence of malformations or fetotoxicity was found, and prospective data regarding paracetamol overdose in pregnancy showed no increase in risk of malformations. Paracetamol may be used throughout pregnancy, but as a precautionary measure should only be taken under a doctor's advice. It is not recommended that paracetamol be used over a prolonged time period, in higher dosages or in combination with other drugs during pregnancy, as the safe use under such conditions has not been con
rmed. • Fertility
º There is limited evidence that drugs which inhibit cyclooxygenase/ prostaglandin synthesis may cause impairment of female fertility
by an effect on ovulation, which is reversible upon withdrawal of treatment. Because paracetamol is believed to inhibit prostaglandin synthesis it is possible that it may impair fertility, although this has not been demonstrated.
9. Undesirable Effects of this Medicine
• Allergic reactions such as skin rash and urticaria – occasional
• Hypersensitivity reactions with symptoms such as pruritus, erythema, urticaria, angiodema, dyspnea or asthma – very rare
• Isolated reports of anaphylactoid reactions or anaphylactic shocks
• Isolated cases of thrombocytopenia, leucopenia, agranulocytosis and pancytopenia associated with paracetamol have been reported.
10. What other medicine or food should be avoided while taking this Medicine?
Hypnotics, Anti-epileptic drugs (phenobarbital, phenytoin, carbamazepine), Rifampicin, Propanthelin, Metoclopropamide, Chloramphenicol, Zidovudine, Barbiturates, Antihistaminics, Benzodiazepines, Oral contraceptives, Cimetidine, Disul
ram, Theophylline,
Ephedrine, Smoking, Alcohol.
11. What should you do if you miss a dose?
No available information.
12. Signs and Symptoms of Overdose
Following overdose, symptoms of paracetamol poisoning often appear within 24-48 hours, but may be delayed beyond this time.
Symptoms: Poisoning is more likely to occur in the presence of alcohol. Liver damage (hepatocellular necrosis) and impairment of liver
function may occur and may progress to hepatic coma. Clinical indications of liver damage may not manifest until 2-4 days following overdose. Hypokalemia and metabolic acidosis (including lactic acidosis) may also develop in the setting of acute and/or chronic overdose.
There have been isolated reports of acute renal failure following paracetamol overdose.
13. What to do when you have taken more than the recommended dosage?
Management of overdose: The cytotoxic effects can be lessened by intravenous administration of sulfhydryl compounds such as cysteamine or N-acetylcysteine, if possible within eight hours of ingestion of the overdose.
14. How should you keep this Medicine?
Store at temperatures not exceeding 30oC.
15. When should you consult your doctor?
If symptoms persist with the administration of the product, consult your doctor.
Marketing Authorization Holder:
Bayer Philippines, Inc.
8th Floor Science Hub Tower 1, Campus Avenue Corner Turin St.,
McKinley Hill Cyberpark, Pinagsama, Taguig City
Manufacturer:
P.T. Bayer Indonesia
Jalan Raya Jakarta Bogor KM 32
Cisalak, Kec. Sukmajaya, Kota Depok,
Jawa Barat, Indonesia
At the
rst sign of any adverse drug reaction, patient must seek medical
attention immediately.
Report any suspected adverse drug reaction or product complaint to
MedInfoCH-PH@bayer.com or drugsafety.philippines@bayer.com or to the
Philippines FDA at adr@fda.gov.ph.
Inquiries can also be directed to:

Bayer Philippines, Inc.
Taguig City, Philippines
E-mail:MedInfoCH-PH@bayer.com
drugsafety.philippines@bayer.com
Registration No.: DR-XY46456
Date of First Authorization:
15 November 2018
Date of Revision of Patient Information Lea
et:
November 2022 (Based from CCDS version 3 dated 22 May 2020)
FEVER & GENERAL PAIN
TBD
Pharmacode
140
190
70

140

190

70

Paracetamol Caffeine

Saridon®

FEVER & GENERAL PAIN

500 mg/50 mg Tablet
Analgesic/Antipyretic

1. Name of the Product
Paracetamol + Caffeine (Saridon®)

2. Description of the Product
The product contains paracetamol and caffeine in tablet form which is classified as antipyretic and analgesic.

3. What is in the Medicine?
Each tablet contains:
Paracetamol…….. 500 mg
Caffeine…… 50 mg

4. Strength of the Medicine
Paracetamol + Caffeine (Saridon®) 500 mg/50 mg Tablet

5. What is the Medicine used for?
- For fast and effective relief of mild to severe headaches.
- For the relief of pain such as headache, toothache, menstrual discomfort, postoperative and rheumatic pain.
- For the relief of pain and fever associated with colds and "flu".

6. How much and how often should you use this Medicine?
Single dose.
Adults: 1-2 tablets
Adolescents aged 12-16 years: 1 tablet
If necessary, up to three single doses may be taken within 24 hours.
The tablets should be taken with ample water or other fluid.
It should not be taken for more than one week or at doses higher than those recommended, except as directed by a doctor.

Patients with hepatic impairment
Paracetamol is contraindicated in patients with severe hepatic insufficiency (Child-Pugh>9). Paracetamol should be used with caution in patients with mild to moderate hepatic insufficiency (Child-Pugh <9).

Patients with renal impairment
Paracetamol should be used with particular caution in patients with severe renal failure. Dose adjustment may be necessary in patients with impaired renal function.

7. When should you not take this Medicine?
It should not be used if any of the following conditions is present:
- Allergy to phenylbutazone-containing products,
- Known hypersensitivity to paracetamol, acetylsalicylic acid or proven allergy to caffeine,
- Inherited deficiency of glucose-6-phosphate dehydrogenase (evidenced by hemolytic anemia),
- Acute hepatic porphyria

Special caution should be taken (lowering of the dosage and/or spacing out of the doses) in the following conditions:
- Impaired hepatic function (in cases of chronic alcohol abuse or hepatitis for example),
- Impaired renal function,
- Gilbert's syndrome (fluctuating benign jaundice secondary to glucuronyl-transferase deficiency),
- Hematopoetic dysfunction

It should not be given to infants or children under 12 years of age.
It is recommended that it should not be used during pregnancy, particularly in the first trimester and in the final six weeks.
It should not be used by nursing women.

8. Care that should be taken when taking this Medicine?
- Should not be taken regularly for a prolonged period, except as directed by a doctor.
- Prolonged intake of paracetamol-containing analgesics resulting in high cumulative doses may in rare instances lead to analgesic nephropathy and largely irreversible renal insufficiency.
- Very rare cases of serious skin reactions have been reported. In the event of skin reddening, rash, blisters or peeling, discontinue paracetamol use and seek medical attention immediately.
- Continual use of analgesics for the treatment of headache may itself lead to chronic headaches.
- Special caution for patients with asthma, chronic rhinitis or chronic urticaria, particularly those hypersensitive to other anti-inflammatory agents.
- There have been isolated reports of asthma attacks and anaphylactic reaction associated with paracetamolcontaining preparations in susceptible individuals.
- Should not be used after the expiry date shown in the pack.

70

TBD

Pharmacode

TBD

Pharmacode

- Pregnancy and Lactation: Category C.
  - Available data from epidemiological studies on the therapeutic use of paracetamol and possible undesirable effects regarding pregnancy or fetal/neonatal development continue to support a positive benefit/risk balance. In reproduction studies of oral paracetamol, no evidence of malformations or fetotoxicity was found, and prospective data regarding paracetamol overdose in pregnancy showed no increase in risk of malformations. Paracetamol may be used throughout pregnancy, but as a precautionary measure should only be taken under a doctor's advice. It is not recommended that paracetamol be used over a prolonged time period, in higher dosages or in combination with other drugs during pregnancy, as the safe use under such conditions has not been confirmed.
- Fertility
  - There is limited evidence that drugs which inhibit cyclooxygenase/ prostaglandin synthesis may cause impairment of female fertility by an effect on ovulation, which is reversible upon withdrawal of treatment. Because paracetamol is believed to inhibit prostaglandin synthesis it is possible that it may impair fertility, although this has not been demonstrated.

9. Undesirable Effects of this Medicine
- Allergic reactions such as skin rash and urticaria – occasional
- Hypersensitivity reactions with symptoms such as pruritus, erythema, urticaria, angiodema, dyspnea or asthma – very rare
- Isolated reports of anaphylactoid reactions or anaphylactic shocks
- Isolated cases of thrombocytopenia, leucopenia, agranulocytosis and pancytopenia associated with paracetamol have been reported.

10. What other medicine or food should be avoided while taking this Medicine?
Hypnotics, Anti-epileptic drugs (phenobarbital, phenytoin, carbamazepine), Rifampicin, Propanthelin, Metoclopropamide, Chloramphenicol, Zidovudine, Barbiturates, Antihistaminics, Benzodiazepines, Oral contraceptives, Cimetidine, Disulfiram, Theophylline, Ephedrine, Smoking, Alcohol.

11. What should you do if you miss a dose?
No available information.

12. Signs and Symptoms of Overdose
Following overdose, symptoms of paracetamol poisoning often appear within 24-48 hours, but may be delayed beyond this time.
Symptoms: Poisoning is more likely to occur in the presence of alcohol. Liver damage (hepatocellular necrosis) and impairment of liver function may occur and may progress to hepatic coma. Clinical indications of liver damage may not manifest until 2-4 days following overdose. Hypokalemia and metabolic acidosis (including lactic acidosis) may also develop in the setting of acute and/or chronic overdose.
There have been isolated reports of acute renal failure following paracetamol overdose.

13. What to do when you have taken more than the recommended dosage?
Management of overdose: The cytotoxic effects can be lessened by intravenous administration of sulfhydryl compounds such as cysteamine or N-acetylcysteine, if possible within eight hours of ingestion of the overdose.

14. How should you keep this Medicine?
Store at temperatures not exceeding 30ºC.

15. When should you consult your doctor?
If symptoms persist with the administration of the product, consult your doctor.

Marketing Authorization Holder:
Bayer Philippines, Inc.
8th Floor Science Hub Tower 1, Campus Avenue Corner Turin St.,
McKinley Hill Cyberpark, Pinagsama, Taguig City

Manufacturer:
P.T. Bayer Indonesia
Jalan Raya Jakarta Bogor KM 32
Cisalak, Kec. Sukmajaya, Kota Depok,
Jawa Barat, Indonesia

At the first sign of any adverse drug reaction, patient must seek medical attention immediately.

Report any suspected adverse drug reaction or product complaint to MedInfoCH-PH@bayer.com or drugsafety.philippines@bayer.com or to the Philippines FDA at adr@fda.gov.ph.
Inquiries can also be directed to:
Bayer Philippines, Inc.
Taguig City, Philippines
E-mail:MedInfoCH-PH@bayer.com
drugsafety.philippines@bayer.com

Registration No.: DR-XY46456

Date of First Authorization:
15 November 2018

Date of Revision of Patient Information Leaflet:
November 2022 (Based from CCDS version 3 dated 22 May 2020)

Bayer

NNNNNN

Saridon Triple Action (Paracetamol + Propyphenazone + Caffeine)

250 mg/150 mg/50 mg Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY36639

Saridon Triple Action: Official FDA Philippines registration record

Brand name	Saridon Triple Action
Generic name	Paracetamol + Propyphenazone + Caffeine
Dosage strength and form	250 mg/150 mg/50 mg Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Analgesic/Antipyretic
Registration number	DR-XY36639
Manufacturer	PT. Bayer Indonesia

Saridon Triple Action: OTC forms registered under the Saridon brand (capsule availability)

OTC capsule form registered under the Saridon brand: NO. 3 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Saridon Triple Action	250 mg/150 mg/50 mg Tablet	Over-the-Counter (OTC) Drug	DR-XY36639
Saridon Max	275 mg Film-Coated Tablet	Over-the-Counter (OTC) Drug	DRP-1872-02
Saridon	500 mg/50 mg Tablet	Over-the-Counter (OTC) Drug	DR-XY46456

Saridon Triple Action: Official leaflet (Package Insert / Patient Information Leaflet)

An official leaflet for this registration is published on the FDA Verification Portal: PI_01 (https://drive.google.com/file/d/1Hfq5y_oT1hSau0Hh0uc6BxB0-1SFW-6Z/view?usp=drivesdk). Its text (OCR) and the original leaflet pages follow.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Saridon"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Saridon Triple Action: Official leaflet text (OCR transcription)

Machine transcription of the official FDA leaflet for Reg. No. DR-XY36639. OCR may contain errors; the original leaflet pages are attached after this section.

Saridon Triple Action leaflet PI_01

91141250
1. Name of the Product
Paracetamol / Propyphenazone / Caffeine (Saridon Triple Action)
2. Description of the Product
Paracetamol / Propyphenazone / Caffeine (Saridon Triple Action) is an analgesic/antipyretic.
3. What is in the Medicine?
450 mg tablet:
Each tablet contains
Paracetamol.....250 mg
Propyphenazone.....150 mg
Caffeine.....50 mg
4. Strength of the Medicine
5. What is the Medicine used for?
• For fast and effective relief of mild to severe headaches
• For the relief of pain such as headache, toothache, menstrual discomfort, postoperative and rheumatic pain
• For the relief of pain and fever associated with colds and
u
6. How much and how often should you use this Medicine?
Single dose.
Adults: 1-2 tablets
Adolescents aged 12-16 years: 1 tablet
If necessary, up to three single doses may be taken within 24 hours.
The tablets should be taken with ample water or other
uid.
Should not be taken for more than one week or at dose higher than those recommended, except as directed by a doctor.
Patients with hepatic impairment:
Paracetamol is contraindicated in patients with severe hepatic insu
ciency (Child-Pugh>9). Paracetamol should be used with caution in
patients with mild to moderate hepatic insu
ciency (Child-Pugh <9).
Patients with renal impairment:
Paracetamol should be used with particular caution in patients with severe renal failure. Dose adjustment may be necessary in patients
with impaired renal function.
7. When should you not take this Medicine?
Saridon Triple Action should not be used if any of the following conditions is present:
• Allergy to pyrazolones or related compounds (hypersensitivity to phenazone, propyphenazone, aminophenazone, metamizole-containing products)
• Allergy to phenylbutazone-containing products
• Known hypersensitivity to paracetamol, acetylsalicylic acid or proven allergy to caffeine
• Inherited de
ciency of glucose-6-phosphate dehydrogenase (evidenced by hemolytic anemia)
• Acute hepatic porphyria
Special caution should be taken (lowering of the dosage and/or spacing out of the doses) in the following conditions:
• Impaired hepatic function (e.g. in cases of chronic alcohol abuse or hepatitis)
• Impaired renal function
• Gilbert’s syndrome (
uctuating benign jaundice secondary to glucoronyl-transferase de
ciency)
• Hematopoetic dysfunction
Saridon Triple Action should not be given to infants or children under 12 years of age.
Pregnancy and Lactation:
• Category C
• It is recommended that Saridon Triple Action should not be used during pregnancy, particularly in the
rst trimester and in the
nal six
weeks.
• Saridon Triple Action should not be used by nursing women.
8. Care that should be taken when taking this Medicine?
• Should not be taken regularly for a prolonged period, except as directed by a doctor.

• Prolonged intake of paracetamol-containing analgesics resulting in high cumulative doses may in rare instances lead to analgesic nephropathy and largely irreversible renal insu
ciency.
• Very rare cases of serious skin reactions have been reported. In the event of skin reddening, rash blisters or peeling, discontinue paracetamol use and seek medical attention immediately.
• Continual use of analgesics for the treatment of headache may itself lead to chronic headaches.
• Special caution for patients with asthma, chronic rhinitis or chronic urticaria, particularly those hypersensitive to other anti-in ammatory agents.
• There have been isolated reports of asthma attacks and anaphylactic reaction associated with propyphenazone and paracetamol-containing preparations in susceptible individuals.
• Should not be used after the expiry date shown in the pack.
• Ability to Drive: Unlikely to produce an impairment of driving performance or performance related to driving.
9. Undesirable Effects of this Medicine
• Allergic reactions such as skin rash and urticaria – occasional
• Hypersensitivity reactions with symptoms such as pruritus, erythema, urticaria, angioedema, dyspnea or asthma – very rare
• Isolated reports of anaphylactoid reactions or anaphylactic shocks
• Isolated cases of thrombocytopenia, leucopenia, agranulocytosis and pancytopenia associated with paracetamol and propyphenazone
have been reported.
10. What other Medicine or Food should be avoided while taking this Medicine?
Hypnotics, Anti-epileptic drugs (phenobarbital, phenytoin, carbamazepine), Rifampicin, Propanthelin, Metoclopropamide, Chloramphenicol, Zidovudine, Barbiturates, Antihistaminics, Benzodiazepines, Oral contraceptives, Cimetidine, Disul
ram, Theophylline,
Ephedrine, Smoking, Alcohol.
11. What should you do if you miss a dose?
No available information.
12. Signs and Symptoms of Overdose
Following overdose, symptoms of paracetamol poisoning often appear within 24-48 hours, but may be delayed beyond this time.
Symptoms: Poisoning is more likely to occur in the presence of alcohol. Liver damage (hepatocellular necrosis) and impairment of liver
function may occur and may progress to hepatic coma. Clinical indications of liver damage may not manifest until 24 days following overdose.
Hypokalemia and metabolic acidosis (including lactic acidosis) may also develop in the setting of acute and/or chronic overdose.
There have been isolated reports of acute renal failure following paracetamol overdose.
13. What to do when you have taken more than the recommended dosage?
Management of overdose: The cytotoxic effects can be lessened by intravenous administration of sulfhydryl compounds such as cysteamine or N-acetylcysteine, if possible within eight hours of ingestion of the overdose.
14. How should you keep this Medicine?
Store at temperatures not exceeding 25o
C.
15. When should you consult your doctor?
If symptoms persist, consult your doctor.
Instructions for Use/Handling: The tablets should be taken with ample water or other
uid.
Manufacturer: PT. Bayer Indonesia
Jalan Raya Jakarta Bogor KM 32,
Cisalak, Kec. Sukmajaya, Kota Depok,
Jawa Barat, Indonesia
Marketing Authorization Holder: Bayer Philippines, Inc.
8th Floor, Science Hub, Tower 1,
Campus Avenue Corner Turin Street,
McKinley Hill Cyberpark, Pinagsama, Taguig City
At the
rst sign of any adverse drug reaction, patient must seek medical attention immediately.
Report any suspected adverse drug reaction or product complaint to MedInfoCH-PH@bayer.com or
drugsafety.philippines@bayer.com or
to the Philippines FDA at adr@fda.gov.ph.
If you want to report a product complaint or side effect, please contact your health care professional or the Philippines FDA at adr@fda.gov.ph.
Inquiries can also be directed to:
Bayer Philippines, Inc.
Taguig City, Philippines
E-mail: MedInfoCH-PH@bayer.com
drugsafety.philippines@bayer.com
Registration Number: DR-XY36639
Date of First Authorization: 11 September 2009 Date of Revision of Patient Information Lea
et: July 2024

(Based on Paracetamol CCDS version 3 dated 22 May 2020)
117
117 117
PT. BAYER INDONESIA
ARTWORK DEVELOPMENT COMMERCIAL
Material Information
Component Type : LFLT
Material Number : 91141250
Pharmacode : 117
Pack-Mat Supplier : GPI/PURA
Artwork Agency : AGP
History
00 / 08/07/24 : TL032620, Revise as per comment from country
01 / 11/07/24 : Revise as per comment from country
02 / 11/07/24 : Adding period after “
uid” and “overdose”, adding comma after “Street”
Material Speciffcation
Dimension : 140 x 190 mm
Material : HVS 60 gsm
Product Information
Product Name : Saridon
Dossage Form (EFF, FCT, Others) : Plain tablet
Pack Size : 120 s ; 10s
Market : PH
SKU : 80483660; 87426327
Change Request : TL032620
Replacement Artwork (for TL) : 89955874
Coating Type :
WB Varnish
Color 1
Black
UV Varnish Matt Varnish
Calendering
Embossing :
Yes No
Hot Stamp/Hot Foil :
Braille
FOLDED
Remark : Cut, B1-B2 Vertical 2x (Folded)
UNFOLDED
Non Coating Color 2
N/A
Color 3
Color 4
Color 5
Color 6
Color 7
Color 8
N/A
Carton Finishing:
Technical & Non Printing Items:
Lea
et Type :

Paracetamol Propyphenazone Caffeine

Saridon TRIPLE ACTION
HEADACHE & MIGRAINE

1. Name of the Product
Paracetamol / Propyphenazone / Caffeine (Saridon Triple Action)

2. Description of the Product
Paracetamol / Propyphenazone / Caffeine (Saridon Triple Action) is an analgesic/antipyretic.

3. What is in the Medicine?
450 mg tablet:
Each tablet contains
Paracetamol.....250 mg
Propyphenazone.....150 mg
Caffeine.....50 mg

4. Strength of the Medicine
Paracetamol 250 mg / Propyphenazone 150 mg / Caffeine 50 mg

5. What is the Medicine used for?
• For fast and effective relief of mild to severe headaches
• For the relief of pain such as headache, toothache, menstrual discomfort, postoperative and rheumatic pain
• For the relief of pain and fever associated with colds and flu

6. How much and how often should you use this Medicine?
Single dose.
Adults: 1-2 tablets
Adolescents aged 12-16 years: 1 tablet
If necessary, up to three single doses may be taken within 24 hours.
The tablets should be taken with ample water or other fluid.
Should not be taken for more than one week or at dose higher than those recommended, except as directed by a doctor.

Patients with hepatic impairment:
Paracetamol is contraindicated in patients with severe hepatic insufficiency (Child-Pugh>9). Paracetamol should be used with caution in patients with mild to moderate hepatic insufficiency (Child-Pugh <9).

Patients with renal impairment:
Paracetamol should be used with particular caution in patients with severe renal failure. Dose adjustment may be necessary in patients with impaired renal function.

7. When should you not take this Medicine?
Saridon Triple Action should not be used if any of the following conditions is present:
• Allergy to pyrazolones or related compounds (hypersensitivity to phenazone, propyphenazone, aminophenazone, metamizole-containing products)
• Allergy to phenylbutazone-containing products
• Known hypersensitivity to paracetamol, acetylsalicylic acid or proven allergy to caffeine
• Inherited deficiency of glucose-6-phosphate dehydrogenase (evidenced by hemolytic anemia)
• Acute hepatic porphyria

Special caution should be taken (lowering of the dosage and/or spacing out of the doses) in the following conditions:
• Impaired hepatic function (e.g. in cases of chronic alcohol abuse or hepatitis)
• Impaired renal function
• Gilbert's syndrome (fluctuating benign jaundice secondary to glucoronyl-transferase deficiency)
• Hematopoetic dysfunction

Saridon Triple Action should not be given to infants or children under 12 years of age.

Pregnancy and Lactation:
• Category C
• It is recommended that Saridon Triple Action should not be used during pregnancy, particularly in the first trimester and in the final six weeks.
• Saridon Triple Action should not be used by nursing women.

8. Care that should be taken when taking this Medicine?
• Should not be taken regularly for a prolonged period, except as directed by a doctor.
• Prolonged intake of paracetamol-containing analgesics resulting in high cumulative doses may in rare instances lead to analgesic nephropathy and largely irreversible renal insufficiency.
• Very rare cases of serious skin reactions have been reported. In the event of skin reddening, rash blisters or peeling, discontinue paracetamol use and seek medical attention immediately.
• Continual use of analgesics for the treatment of headache may itself lead to chronic headaches.
• Special caution for patients with asthma, chronic rhinitis or chronic urticaria, particularly those hypersensitive to other anti-inflammatory agents.

• There have been isolated reports of asthma attacks and anaphylactic reaction associated with propyphenazone and paracetamol-containing preparations in susceptible individuals.
• Should not be used after the expiry date shown in the pack.
• Ability to Drive: Unlikely to produce an impairment of driving performance or performance related to driving.

9. Undesirable Effects of this Medicine
• Allergic reactions such as skin rash and urticaria – occasional
• Hypersensitivity reactions with symptoms such as pruritus, erythema, urticaria, angioedema, dyspnea or asthma – very rare
• Isolated reports of anaphylactoid reactions or anaphylactic shocks
• Isolated cases of thrombocytopenia, leucopenia, agranulocytosis and pancytopenia associated with paracetamol and propyphenazone have been reported.

10. What other Medicine or Food should be avoided while taking this Medicine?
Hypnotics, Anti-epileptic drugs (phenobarbital, phenytoin, carbamazepine), Rifampicin, Propanthelin, Metoclopropamide, Chloramphenicol, Zidovudine, Barbiturates, Antihistaminics, Benzodiazepines, Oral contraceptives, Cimetidine, Disulfiram, Theophylline, Ephedrine, Smoking, Alcohol.

11. What should you do if you miss a dose?
No available information.

12. Signs and Symptoms of Overdose
Following overdose, symptoms of paracetamol poisoning often appear within 24-48 hours, but may be delayed beyond this time.

Symptoms: Poisoning is more likely to occur in the presence of alcohol. Liver damage (hepatocellular necrosis) and impairment of liver function may occur and may progress to hepatic coma. Clinical indications of liver damage may not manifest until 24 days following overdose.

Hypokalemia and metabolic acidosis (including lactic acidosis) may also develop in the setting of acute and/or chronic overdose. There have been isolated reports of acute renal failure following paracetamol overdose.

13. What to do when you have taken more than the recommended dosage?
Management of overdose: The cytotoxic effects can be lessened by intravenous administration of sulfhydryl compounds such as cysteamine or N-acetylcysteine, if possible within eight hours of ingestion of the overdose.

14. How should you keep this Medicine?
Store at temperatures not exceeding 25°C.

15. When should you consult your doctor?
If symptoms persist, consult your doctor.

Instructions for Use/Handling: The tablets should be taken with ample water or other fluid.

Manufacturer: **PT. Bayer Indonesia**
Jalan Raya Jakarta Bogor KM 32,
Cisalak, Kec. Sukmajaya, Kota Depok,
Jawa Barat, Indonesia

Marketing Authorization Holder: **Bayer Philippines, Inc.**
8th Floor, Science Hub, Tower 1,
Campus Avenue Corner Turin Street,
McKinley Hill Cyberpark, Pinagsama, Taguig City

At the first sign of any adverse drug reaction, patient must seek medical attention immediately.

Report any suspected adverse drug reaction or product complaint to MedInfoCH-PH@bayer.com or drugsafety.philippines@bayer.com or to the Philippines FDA at adr@fda.gov.ph.

If you want to report a product complaint or side effect, please contact your health care professional or the Philippines FDA at adr@fda.gov.ph.

Inquiries can also be directed to:
Bayer Philippines, Inc.
Taguig City, Philippines
E-mail: MedInfoCH-PH@bayer.com
drugsafety.philippines@bayer.com

Registration Number: DR-XY36639
Date of First Authorization: 11 September 2009
Date of Revision of Patient Information Leaflet: July 2024
(Based on Paracetamol CCDS version 3 dated 22 May 2020)

Bayer
91141250

PT. BAYER INDONESIA
ARTWORK DEVELOPMENT
COMMERCIAL

Product Information

Product Name	**: Saridon**
Dossage Form (EFF, FCT, Others)	**: Plain tablet**
Pack Size	**: 120 s ; 10s**
Market	**: PH**
SKU	**: 80483660; 87426327**
Change Request	**: TL032620**
Replacement Artwork (for TL)	**: 89955874**

Material Information

Component Type	**: LFLT**
Material Number	**: 91141250**
Pharmacode	**: 117**
Pack-Mat Supplier	**: GPI/PURA**
Artwork Agency	**: AGP**

Material Specification

Dimension : 140 x 190 mm
Material : HVS 60 gsm

Color 1	Color 2	Color 3	Color 4
Black	N/A	N/A	N/A
Color 5	**Color 6**	**Color 7**	**Color 8**
N/A	N/A	N/A	N/A

Technical & Non Printing Items:
N/A

Carton Finishing:
Coating Type :
◯ WB Varnish ◯ UV Varnish ◯ Matt Varnish
◯ Calendering ◯ Non Coating
Embossing :
◯ Yes ◯ No
Hot Stamp/Hot Foil :
◯ Yes ◯ No
Braille
◯ Yes ◯ No

Leaflet Type :
● FOLDED ◯ UNFOLDED
Remark : Cut, B1-B2 Vertical 2x (Folded)

History

00 / 08/07/24 : TL032620, Revise as per comment from country
01 / 11/07/24 : Revise as per comment from country
02 / 11/07/24 : Adding period after "fluid" and "overdose", adding comma after "Street"

Saridon Max (Naproxen Sodium)

275 mg Film-Coated Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DRP-1872-02

Saridon Max: Official FDA Philippines registration record

Brand name	Saridon Max
Generic name	Naproxen Sodium
Dosage strength and form	275 mg Film-Coated Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Non-Selective COX Inhibitor
Registration number	DRP-1872-02
Manufacturer	Interphil Laboratories, Inc.

Saridon Max: OTC forms registered under the Saridon brand (capsule availability)

OTC capsule form registered under the Saridon brand: NO. 3 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Saridon Triple Action	250 mg/150 mg/50 mg Tablet	Over-the-Counter (OTC) Drug	DR-XY36639
Saridon Max	275 mg Film-Coated Tablet	Over-the-Counter (OTC) Drug	DRP-1872-02
Saridon	500 mg/50 mg Tablet	Over-the-Counter (OTC) Drug	DR-XY46456

Saridon Max: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRP-1872-02 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Saridon"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Tempra (Paracetamol)

325 mg Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-4396

Tempra: Official FDA Philippines registration record

Brand name	Tempra
Generic name	Paracetamol
Dosage strength and form	325 mg Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Analgesic/ Antipyretic
Registration number	DR-4396
Manufacturer	Interphil Laboratories, Inc.

Tempra: OTC forms registered under the Tempra brand (capsule availability)

OTC capsule form registered under the Tempra brand: NO. 24 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Tempra	100 mg/mL Syrup (Pediatric Drops) (Orange Flavor)	Over-the-Counter (OTC) Drug	DR-XY49330
Tempra	100 mg/mL Syrup (Pediatric Drops) Orange Flavor	Over-the-Counter (OTC) Drug	DR-XY46708
Tempra	100 mg/mL Syrup (Pediatric Drops) Strawberry Flavor	Over-the-Counter (OTC) Drug	DR-XY46709
Tempra (Orange Flavor)	100mg/mL Syrup (Pediatric Drops)	Over-the-Counter (OTC) Drug	DR-XY26514
Tempra	120 mg/ 5 mL Syrup (Orange Flavor)	Household Remedy (HR)	DRHR-1766
Tempra	120 mg/ 5 mL Syrup (Orange Flavor)	Household Remedy (HR)	DRHR-460
Tempra	120 mg/ 5 mL Syrup (Strawberry Flavor)	Household Remedy (HR)	DRHR-853
Tempra	120 mg/5 mL Syrup (Orange Flavor)	Household Remedy (HR)	DRHR-2431
Tempra	120 mg/5 mL Syrup (Strawberry Flavor)	Household Remedy (HR)	DRHR-1944
Tempra	120mg/5mL Syrup (Orange Flavor)	Household Remedy (HR)	DRHR-1942
Tempracof	15mg/5mL Syrup	Over-the-Counter (OTC) Drug	DR-XY46459
Tempra Forte	250 mg/ 5 mL Syrup (Strawberry Flavor)	Over-the-Counter (OTC) Drug	DR-XY24336
Tempra Forte (Orange Flavor)	250 mg/5 mL Syrup	Over-the-Counter (OTC) Drug	DR-XY46515
Tempra Forte	250 mg/5 mL Syrup (Strawberry Flavor)	Over-the-Counter (OTC) Drug	DR-XY46514

Brand name	Dosage strength and form	Classification	Reg. No.
Tempra Forte	250mg/ 5mL Syrup (Strawberry Flavor)	Over-the-Counter (OTC) Drug	DR-XY45843
Tempracof Forte	30 mg/5 mL Syrup	Over-the-Counter (OTC) Drug	DR-XY46457
Tempracof	30 mg/5 mL Syrup (Orange Vanilla Butterscotch Flavor)	Over-the-Counter (OTC) Drug	DRP-1987-04
Tempra	325 mg Tablet	Over-the-Counter (OTC) Drug	DR-4396
Temprazin	5 mg/5 mL Syrup	Over-the-Counter (OTC) Drug	DRP-1468-05
Tempra Blizz	500 mg Effervescent Tablet (Orange Flavor)	Over-the-Counter (OTC) Drug	DR-XY48911
Tempra Blizz	500 mg Granules for Solution (Orange Flavor)	Over-the-Counter (OTC) Drug	DR-XY48940
Tempra Forte	500 mg Tablet	Over-the-Counter (OTC) Drug	DR-X2535
Tempra Boost-C	500 mg/40 mg Granules for Solution	Over-the-Counter (OTC) Drug	DRP-9330-01
Tempracof	7.5 mg/mL Syrup (Pediatric Drops)	Over-the-Counter (OTC) Drug	DR-XY46458

Tempra: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-4396 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Tempra"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Tempra Blizz (Paracetamol)

500 mg Effervescent Tablet (Orange Flavor) | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY48911

Tempra Blizz: Official FDA Philippines registration record

Brand name	Tempra Blizz
Generic name	Paracetamol
Dosage strength and form	500 mg Effervescent Tablet (Orange Flavor)
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Analgesic / Antipyretic
Registration number	DR-XY48911
Manufacturer	DHG Pharmaceutical Joint-Stock Company

Tempra Blizz: OTC forms registered under the Tempra brand (capsule availability)

OTC capsule form registered under the Tempra brand: NO. 24 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Tempra	100 mg/mL Syrup (Pediatric Drops) (Orange Flavor)	Over-the-Counter (OTC) Drug	DR-XY49330
Tempra	100 mg/mL Syrup (Pediatric Drops) Orange Flavor	Over-the-Counter (OTC) Drug	DR-XY46708
Tempra	100 mg/mL Syrup (Pediatric Drops) Strawberry Flavor	Over-the-Counter (OTC) Drug	DR-XY46709
Tempra (Orange Flavor)	100mg/mL Syrup (Pediatric Drops)	Over-the-Counter (OTC) Drug	DR-XY26514
Tempra	120 mg/ 5 mL Syrup (Orange Flavor)	Household Remedy (HR)	DRHR-1766
Tempra	120 mg/ 5 mL Syrup (Orange Flavor)	Household Remedy (HR)	DRHR-460
Tempra	120 mg/ 5 mL Syrup (Strawberry Flavor)	Household Remedy (HR)	DRHR-853
Tempra	120 mg/5 mL Syrup (Orange Flavor)	Household Remedy (HR)	DRHR-2431
Tempra	120 mg/5 mL Syrup (Strawberry Flavor)	Household Remedy (HR)	DRHR-1944
Tempra	120mg/5mL Syrup (Orange Flavor)	Household Remedy (HR)	DRHR-1942
Tempracof	15mg/5mL Syrup	Over-the-Counter (OTC) Drug	DR-XY46459
Tempra Forte	250 mg/ 5 mL Syrup (Strawberry Flavor)	Over-the-Counter (OTC) Drug	DR-XY24336
Tempra Forte (Orange Flavor)	250 mg/5 mL Syrup	Over-the-Counter (OTC) Drug	DR-XY46515
Tempra Forte	250 mg/5 mL Syrup (Strawberry Flavor)	Over-the-Counter (OTC) Drug	DR-XY46514

Brand name	Dosage strength and form	Classification	Reg. No.
Tempra Forte	250mg/ 5mL Syrup (Strawberry Flavor)	Over-the-Counter (OTC) Drug	DR-XY45843
Tempracof Forte	30 mg/5 mL Syrup	Over-the-Counter (OTC) Drug	DR-XY46457
Tempracof	30 mg/5 mL Syrup (Orange Vanilla Butterscotch Flavor)	Over-the-Counter (OTC) Drug	DRP-1987-04
Tempra	325 mg Tablet	Over-the-Counter (OTC) Drug	DR-4396
Temprazin	5 mg/5 mL Syrup	Over-the-Counter (OTC) Drug	DRP-1468-05
Tempra Blizz	500 mg Effervescent Tablet (Orange Flavor)	Over-the-Counter (OTC) Drug	DR-XY48911
Tempra Blizz	500 mg Granules for Solution (Orange Flavor)	Over-the-Counter (OTC) Drug	DR-XY48940
Tempra Forte	500 mg Tablet	Over-the-Counter (OTC) Drug	DR-X2535
Tempra Boost-C	500 mg/40 mg Granules for Solution	Over-the-Counter (OTC) Drug	DRP-9330-01
Tempracof	7.5 mg/mL Syrup (Pediatric Drops)	Over-the-Counter (OTC) Drug	DR-XY46458

Tempra Blizz: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY48911 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Tempra"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.