MediFind PH: OTC Medicine Product Leaflets

Volume 3: Digestive Health, Antiseptic, Antifungal and Eye Care

32 product leaflets compiled in registration-file order. Each leaflet is included unchanged after this contents page and has its own bookmark.

No.	Product	Reg. No.	Page
1	Bactidol (Hexetidine) 0.1% Solution	DRHR-102	3
2	Bactidol Extra Soothing (Dichlorobenzyl Alcohol + Amylmetacresol) 1.2 mg/600 mcg Lozenge (Orange Flavor)	DR-XY46809	4
3	Betadine (Povidone-Iodine) 4.5 mg/mL (0.45% w/v) Oromucosal Solution (Throat Spray)	DRHR-1749	5
4	Betadine (Povidone-Iodine) 2.5% w/w Dry Powder Spray	DRHR-1764	6
5	Bonamine (Meclizine Hydrochloride) 12.5 mg Chewable Tablet	DR-XY19867	7
6	Bonamine (Meclizine Hydrochloride) 25 mg Chewable Tablet	DR-XY22028	11
7	Buscopan (Hyoscine N-Butylbromide) 10 mg Tablet	DR-XY41205	16
8	Canesten (Clotrimazole) 100 mg Vaginal Tablet	DR-XY47618	17
9	Canesten (Clotrimazole) 10 mg/mL (1%) Topical Solution	DR-8992	22
10	Canesten (Clotrimazole) 10 mg/g (1% w/w) Cream	DR-XY42700	27
11	Daktarin (Miconazole nitrate) 20 mg per 1 g cream	DR-XY14359	32
12	Daktarin (Miconazole) 20 mg/g Oral Gel	DR-XY14361	33
13	Diatabs (Loperamide Hydrochloride) 2 mg Capsule	DR-XY24964	34
14	Diatabs Advance (Loperamide + Simeticone) 2 mg/ 125 mg Softgel Capsule	DR-XY46907	35
15	Dulcolax (Bisacodyl) 5 mg Enteric-Coated Tablet	DR-XY42393	36
16	Dulcolax (Bisacodyl) 10 mg Rectal Suppository	DR-XY44557	37
17	Eye Mo Cool Red Eyes (Tetrahydrozoline Hydrochloride) 0.05% Solution (Eyedrops)	DRHR-2428	38
18	Eye Mo Moisturizing Formula (Hypromellose) 3 mg/mL Ophthalmic Solution	DR-XY35419	39
19	Gaviscon (Sodium Alginate + Sodium Bicarbonate + Calcium Carbonate) 250 mg/ 133.5 mg/ 80 mg Chewable Tablet	DR-XY35582	40
20	Gaviscon Double Action (Sodium Alginate + Sodium Bicarbonate + Calcium Carbonate) 250 mg/106.5 mg/187.5 mg Chewable Tablet	DR-XY43658	44
21	Gaviscon Infant (Sodium Alginate + Magnesium Alginate) 225 mg/ 87.5 mg Powder for Oral Suspension	DR-XY46801	49
22	Gaviscon (Sodium Alginate + Sodium Bicarbonate + Calcium Carbonate) 500 mg/ 267 mg/ 160 mg per 10 mL Oral Suspension	DR-XY34746	55
23	Gaviscon Double Action (Sodium Alginate + Sodium Bicarbonate + Calcium Carbonate) 500 mg/213 mg/325 mg per 10 mL Suspension	DR-XY43337	56
24	Hydrite (Oral Rehydration Salts) Granules for Solution (Melon Flavor)	DRHR-1810	57
25	Imodium (Loperamide hydrochloride) 2 mg Capsule	DR-XY14916	58
26	Imodium Complete Relief (Loperamide + Simethicone) 2 mg/125 mg Tablet	DR-XY49096	59
27	Kremil-S (Aluminum Hydroxide + Magnesium Hydroxide + Simethicone) 178 mg/233 mg/30 mg Chewable Tablet	DR-XY29748	60
28	Kremil-S Spasm Relief (Hyoscine-N-Butylbromide) 10 mg Film-Coated Tablet	DRP-16071-01	61
29	Maalox No. 1 (Aluminum hydroxide + Magnesium hydroxide) 200 mg/200 mg Chewable Tablet	DRHR-697	62

No.	Product	Reg. No.	Page
30	Maalox (Aluminum hydroxide/ Magnesium hydroxide) 225mg/ 200mg per 5mL Suspension	DRHR-698	63
31	Pedialyte 45 (Oral Rehydration Salts (Maintenance)) (see reverse for formulation) Oral Solution (Fruit Flavor)	DRHR-202	64
32	Visine (Tetrahydrozoline Hydrochloride) 500 mcg/mL (0.05% w/v) Ophthalmic Solution (Eye Drops)	DR-XY37227	65

Bactidol (Hexetidine)

0.1% Solution | Household Remedy (HR) | FDA Reg. No. DRHR-102

Bactidol: Official FDA Philippines registration record

Brand name	Bactidol
Generic name	Hexetidine
Dosage strength and form	0.1% Solution
Classification	Household Remedy (HR)
Pharmacologic category	Oral Antiseptic
Registration number	DRHR-102
Manufacturer	Interphil Laboratories, Inc.

Bactidol: OTC forms registered under the Bactidol brand (capsule availability)

OTC capsule form registered under the Bactidol brand: NO. 3 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Bactidol	0.1% Solution	Household Remedy (HR)	DRHR-102
Bactidol Extra Soothing	1.2 mg/600 mcg Lozenge (Orange Flavor)	Over-the-Counter (OTC)	DR-XY46809
Bactidol Heavy Phlegm Relief	100 mg/4 mg per 5mL Syrup	Over-the-Counter (OTC) Drug	DR-XY47306

Bactidol: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-102 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Bactidol"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Bactidol Extra Soothing (Dichlorobenzyl Alcohol + Amylmetacresol)

1.2 mg/600 mcg Lozenge (Orange Flavor) | Over-the-Counter (OTC) | FDA Reg. No. DR-XY46809

Bactidol Extra Soothing: Official FDA Philippines registration record

Brand name	Bactidol Extra Soothing
Generic name	Dichlorobenzyl Alcohol + Amylmetacresol
Dosage strength and form	1.2 mg/600 mcg Lozenge (Orange Flavor)
Classification	Over-the-Counter (OTC)
Pharmacologic category	Antiseptic
Registration number	DR-XY46809
Manufacturer	Unique Pharmaceutical Laboratories (Division of J.B. Chemicals & Pharmaceuticals Ltd.)

Bactidol Extra Soothing: OTC forms registered under the Bactidol brand (capsule availability)

OTC capsule form registered under the Bactidol brand: NO. 3 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Bactidol	0.1% Solution	Household Remedy (HR)	DRHR-102
Bactidol Extra Soothing	1.2 mg/600 mcg Lozenge (Orange Flavor)	Over-the-Counter (OTC)	DR-XY46809
Bactidol Heavy Phlegm Relief	100 mg/4 mg per 5mL Syrup	Over-the-Counter (OTC) Drug	DR-XY47306

Bactidol Extra Soothing: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY46809 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Bactidol"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Betadine (Povidone-Iodine)

4.5 mg/mL (0.45% w/v) Oromucosal Solution (Throat Spray) | Over-the-Counter (OTC) Drug | FDA Reg. No. DRHR-1749

Betadine: Official FDA Philippines registration record

Brand name	Betadine
Generic name	Povidone-Iodine
Dosage strength and form	4.5 mg/mL (0.45% w/v) Oromucosal Solution (Throat Spray)
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antiseptic
Registration number	DRHR-1749
Manufacturer	Mundipharma Pharmaceuticals Limited

Betadine: OTC forms registered under the Betadine brand (capsule availability)

OTC capsule form registered under the Betadine brand: NO. 8 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Betadine	1% Oral Antiseptic Solution	Household Remedy (HR)	DRHR-328
Betadine	10% Topical Solution	Household Remedy (HR)	DRHR-1809
Betadine	10% w/w Ointment	Over-the-Counter (OTC) Drug	DRHR-443
Betadine	100 mg/mL (10% w/v) Povidone-Iodine	Household Remedy (HR)	DRHR-2130
Betadine	100 mg/mL (10%) Vaginal Douche	Over-the-Counter (OTC) Drug	DRHR-455
Betadine	2.5% w/w Dry Powder Spray	Household Remedy (HR)	DRHR-1764
Betadine	4.5 mg/mL (0.45% w/v) Oromucosal Solution (Throat Spray)	Over-the-Counter (OTC) Drug	DRHR-1749
Betadine	50 mg/g (5% w/w) Cream	Over-the-Counter (OTC) Drug	DRHR-384

Betadine: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-1749 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Betadine"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Betadine (Povidone-Iodine)

2.5% w/w Dry Powder Spray | Household Remedy (HR) | FDA Reg. No. DRHR-1764

Betadine: Official FDA Philippines registration record

Brand name	Betadine
Generic name	Povidone-Iodine
Dosage strength and form	2.5% w/w Dry Powder Spray
Classification	Household Remedy (HR)
Pharmacologic category	Antiseptic and Disinfectant
Registration number	DRHR-1764
Manufacturer	Mundipharma Pharmaceuticals Ltd.

Betadine: OTC forms registered under the Betadine brand (capsule availability)

OTC capsule form registered under the Betadine brand: NO. 8 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Betadine	1% Oral Antiseptic Solution	Household Remedy (HR)	DRHR-328
Betadine	10% Topical Solution	Household Remedy (HR)	DRHR-1809
Betadine	10% w/w Ointment	Over-the-Counter (OTC) Drug	DRHR-443
Betadine	100 mg/mL (10% w/v) Povidone-Iodine	Household Remedy (HR)	DRHR-2130
Betadine	100 mg/mL (10%) Vaginal Douche	Over-the-Counter (OTC) Drug	DRHR-455
Betadine	2.5% w/w Dry Powder Spray	Household Remedy (HR)	DRHR-1764
Betadine	4.5 mg/mL (0.45% w/v) Oromucosal Solution (Throat Spray)	Over-the-Counter (OTC) Drug	DRHR-1749
Betadine	50 mg/g (5% w/w) Cream	Over-the-Counter (OTC) Drug	DRHR-384

Betadine: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-1764 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Betadine"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Bonamine (Meclizine Hydrochloride)

12.5 mg Chewable Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY19867

Bonamine: Official FDA Philippines registration record

Brand name	Bonamine
Generic name	Meclizine Hydrochloride
Dosage strength and form	12.5 mg Chewable Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Anti-emetic
Registration number	DR-XY19867
Manufacturer	Interphil Laboratories, Inc.

Bonamine: OTC forms registered under the Bonamine brand (capsule availability)

OTC capsule form registered under the Bonamine brand: NO. 2 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Bonamine	12.5 mg Chewable Tablet	Over-the-Counter (OTC) Drug	DR-XY19867
Bonamine	25 mg Chewable Tablet	Over-the-Counter (OTC) Drug	DR-XY22028

Bonamine: Official leaflet (Package Insert / Patient Information Leaflet)

An official leaflet for this registration is published on the FDA Verification Portal: PIL_01 (https://drive.google.com/file/d/16PqnBAIcxe1Neg5-aB37fNfAKZfL5o8l/view?usp=drivesdk). Its text (OCR) and the original leaflet pages follow.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Bonamine"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Bonamine: Official leaflet text (OCR transcription)

Machine transcription of the official FDA leaflet for Reg. No. DR-XY19867. OCR may contain errors; the original leaflet pages are attached after this section.

Bonamine leaflet PIL_01

•·
Item Code: Item Description:
Bonamine 12.5 mg - Chewable
Material Code: Product Information Leaflet
Reason of Revision: Dimension:
Revision No.:
Note
I-BNM-0 1 1111111 111111111 IIII 11111 1111
C82Z
I Meclizine HCI I
Bona mine®
12.5 mg Chewable Tablet
Anti-emetic
DESCRIPTION OF THE PRODUCT
Meclizine Hydrochloride (Bonamine®
) Chewable Tablet is a pale orange, mottled, round,
biconvex tablet, plain on one side and scored on the other side with characteristic sweet
yogurt-like flavor.
WHAT IS IN THE MEDICINE?
Active Pharmaceutical Ingredient (API):
Meclizine Hydrochloride
Excipients used are:
. . Glycine, Corn Starch, Carmellose Calcium, Flavor, Magnesium Stearate, Sodium
Saccharin, Colorant, Sodium Lauryl Sulfate, and Purified Water.
STRENGTH OF THE MEDICINE
Meclizine Hydrochloride (Bonamine®
) Chewable_Tablet for Kids:
. Each chewable tablet contains 12.5 mg of MecIIzme Hydrochloride.
WHAT IS THIS MEDICINE USED FOR?
) Chewable Tablet i_s used_ for _the prevention and
treatment of nausea, vomiting, or dizziness associated with motion sickness.
HOW MUCH AND HOW OFTEN SHOULD YOU USE THIS MEDICINE?
A daily dose should be taken one hour before travel starts for 24-hour protection.
. The tablet may be chewed and swallowed with or without water or swallowed whole with
water.
MECLIZINE HYDROCHLORIDE (BONAMINE®) 12.5 MG CHEWABLE
TABLET FOR KIDS
Age Recommended Dose
Children > 11 yrs. old 2 tablets
Or, as prescribed by a physician. If additional dose is needed, interval should be 4 hours or longer.
Children 5-15 yrs old should be limited to two single doses only.
WHEN SHOULD YOU NOT TAKE THIS MEDICINE?
• A known hypersensitivity to Meclizine or any ingredient in this product.
• Patients taking monoamine oxidase inhibitor antidepressant.
• Pregnant or breastfeeding mothers.
CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE
• This drug may cause drowsiness. Take due precautions against driving a car or
operating dangerous machinery.
• A doctor must always be consulted before giving Meclizine especially if the patient
is taking other medications (antidepressant) and patients who are alcoholics or who
regularly ingest alcohol in excess before giving this drug.
• It is also wise to remember to check the expiration date on the package or label of
this product. Do not use after expiry date.
FRONT
Dimensions are in millimeters unless otherwise noted
Supplier: Revision Date:
Color Used: Graphics Artist:
Black
UNDESIRABLE EFFECTS OF THIS MEDICINE
Drowsiness, fatigue, dry mouth, constipation, headache, stomach upset, and rarely

blurred vision have been reported with the use of Meclizine.
WHAT OTHER MEDICINE OR FOOD SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?
Avoid alcohol, sedatives, anti-allergic drugs and tranquilizers while taking Meclizine as this may increase the drowsiness effect.
WHAT SHOULD YOU DO IF YOU MISS A DOSE?
Do not double the dose. If you miss a dose, just take the next dose as scheduled_ (i.e., once a day) if needed to prevent and treat nausea, vomiting, or dizziness associated with motion sickness.
SIGNS AND SYMPTOMS OF OVERDOSE
Overdose of Meclizine may manifest as follows:
• Drowsiness
• Dry mouth
• Fatigue
• Rarely, blurred vision
WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN THE RECOMMENDED DOSAGE?
If more than the recommended dose has been given, consult a health professional immediately. Do not exceed recommended dosage.
HOW SHOULD YOU KEEP THIS MEDICINE?
Store at temperatures not exceeding 30°
C. Keep out of reach of children.
WHEN SHOULD YOU CONSULT YOUR DOCTOR?
Pregnant or breastfeeding mothers, patients with increased pressure in the e_ye, asthma, emphysema, chronic pulmonary disease, difficulty in breathing and th_ose with difficulty in urination due to enlargement of the prostate gland and those taking sedatives and tranquilizers should take Meclizine only under the direction of a doctor.
HOW SUPPLIED?
Meclizine Hydrochloride (Bonamine®
) 12.5 mg Chewable Tablet for Kids: Alu/PVC Blister
Pack x 1 O's (Box of 120's)
NAME AND ADDRESS OF MARKETING AUTHORIZATION HOLDER
Taisho Pharmaceuticals (Philippines), Inc.
8th Floor CLIPP Center Building,
11th Avenue corner 39th Street,
Bonifacio Global City, Taguig 1634, Philippines
Under Authority of:
Taisho Pharmaceutical Co., Ltd.,
Tokyo, Japan
Trademark Owner
NAME AND ADDRESS OF MANUFACTURER
Interphil Laboratories, Inc.
Canlubang Industrial Estate,
Bo. Pittland, Cabuyao, Laguna, Philippines
ADR REPORTING STATEMENT
For suspected adverse drug reaction, report_ to FDA: www.fda.gov.ph
Seek medical attention immediately at the first sign of any adverse drug reaction.
REGISTRATION NUMBER
Meclizine Hydrochloride (Bonamine®
) 12.5 mg Chewable Tablet for Kids: DR-XY19867
RENEWAL OF THE AUTHORIZATION
) 12.5 mg Chewable Tablet for Kids: 21 October 2029
DATE OF REVISION OF THE PATIENT INFORMATION LEAFLET (PIL)
May 2024
@-1 0524
BACK
r-'
O'I
0

Item Code:	Item Description: **Bonamine 12.5 mg - Chewable** **Product Information Leaflet**		Supplier:	Revision Date:
Material Code:				
Revision No.:	Reason of Revision:	Dimension: 75 x 160 mm Note: Dimensions are in millimeters unless otherwise noted. Tolerances are ± 1 mm.	Color Used: Black	Graphics Artist:

75 mm

160 mm

I-BNM-0

C82Z

Meclizine HCl
Bonamine®

12.5 mg Chewable Tablet
Anti-emetic

DESCRIPTION OF THE PRODUCT

Meclizine Hydrochloride (Bonamine®) Chewable Tablet is a pale orange, mottled, round, biconvex tablet, plain on one side and scored on the other side with characteristic sweet yogurt-like flavor.

WHAT IS IN THE MEDICINE?

Active Pharmaceutical Ingredient (API):
Meclizine Hydrochloride

Excipients used are:
Glycine, Corn Starch, Carmellose Calcium, Flavor, Magnesium Stearate, Sodium Saccharin, Colorant, Sodium Lauryl Sulfate, and Purified Water.

STRENGTH OF THE MEDICINE

Meclizine Hydrochloride (Bonamine®) Chewable Tablet for Kids:
Each chewable tablet contains 12.5 mg of Meclizine Hydrochloride.

WHAT IS THIS MEDICINE USED FOR?

Meclizine Hydrochloride (Bonamine®) Chewable Tablet is used for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.

HOW MUCH AND HOW OFTEN SHOULD YOU USE THIS MEDICINE?

A daily dose should be taken one hour before travel starts for 24-hour protection.
The tablet may be chewed and swallowed with or without water or swallowed whole with water.

MECLIZINE HYDROCHLORIDE (BONAMINE®) 12.5 MG CHEWABLE TABLET FOR KIDS

Age	Recommended Dose
Children ≥ 11 yrs. old	2 tablets
5 to 10 yrs. old	1 tablet
Or, as prescribed by a physician. If additional dose is needed, interval should be 4 hours or longer. Children 5-15 yrs old should be limited to two single doses only.	

WHEN SHOULD YOU NOT TAKE THIS MEDICINE?

- A known hypersensitivity to Meclizine or any ingredient in this product.
- Patients taking monoamine oxidase inhibitor antidepressant.
- Pregnant or breastfeeding mothers.

CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE

- This drug may cause drowsiness. Take due precautions against driving a car or operating dangerous machinery.
- A doctor must always be consulted before giving Meclizine especially if the patient is taking other medications (antidepressant) and patients who are alcoholics or who regularly ingest alcohol in excess before giving this drug.
- It is also wise to remember to check the expiration date on the package or label of this product. Do not use after expiry date.

FRONT

75 mm

160 mm

UNDESIRABLE EFFECTS OF THIS MEDICINE

Drowsiness, fatigue, dry mouth, constipation, headache, stomach upset, and rarely blurred vision have been reported with the use of Meclizine.

WHAT OTHER MEDICINE OR FOOD SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?

Avoid alcohol, sedatives, anti-allergic drugs and tranquilizers while taking Meclizine as this may increase the drowsiness effect.

WHAT SHOULD YOU DO IF YOU MISS A DOSE?

Do not double the dose. If you miss a dose, just take the next dose as scheduled (i.e., once a day) if needed to prevent and treat nausea, vomiting, or dizziness associated with motion sickness.

SIGNS AND SYMPTOMS OF OVERDOSE

Overdose of Meclizine may manifest as follows:

- Drowsiness
- Dry mouth
- Fatigue
- Rarely, blurred vision

WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN THE RECOMMENDED DOSAGE?

If more than the recommended dose has been given, consult a health professional immediately. Do not exceed recommended dosage.

HOW SHOULD YOU KEEP THIS MEDICINE?

Store at temperatures not exceeding 30°C. Keep out of reach of children.

WHEN SHOULD YOU CONSULT YOUR DOCTOR?

Pregnant or breastfeeding mothers, patients with increased pressure in the eye, asthma, emphysema, chronic pulmonary disease, difficulty in breathing and those with difficulty in urination due to enlargement of the prostate gland and those taking sedatives and tranquilizers should take Meclizine only under the direction of a doctor.

HOW SUPPLIED?

Meclizine Hydrochloride (Bonamine®) 12.5 mg Chewable Tablet for Kids: Alu/PVC Blister Pack x 10's (Box of 120's)

NAME AND ADDRESS OF MARKETING AUTHORIZATION HOLDER

Taisho Pharmaceuticals (Philippines), Inc.
8th Floor CLIPP Center Building,
11th Avenue corner 39th Street,
Bonifacio Global City, Taguig 1634, Philippines

Under Authority of:
Taisho Pharmaceutical Co., Ltd.,
Tokyo, Japan
Trademark Owner

NAME AND ADDRESS OF MANUFACTURER

Interphil Laboratories, Inc.
Canlubang Industrial Estate,
Bo. Pittland, Cabuyao, Laguna, Philippines

ADR REPORTING STATEMENT

For suspected adverse drug reaction, report to FDA: **www.fda.gov.ph**
Seek medical attention immediately at the first sign of any adverse drug reaction.

REGISTRATION NUMBER

Meclizine Hydrochloride (Bonamine®) 12.5 mg Chewable Tablet for Kids: DR-XY19867

RENEWAL OF THE AUTHORIZATION

Meclizine Hydrochloride (Bonamine®) 12.5 mg Chewable Tablet for Kids: 21 October 2029

DATE OF REVISION OF THE PATIENT INFORMATION LEAFLET (PIL)

May 2024

-1 0524

BACK

Bonamine (Meclizine Hydrochloride)

25 mg Chewable Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY22028

Bonamine: Official FDA Philippines registration record

Brand name	Bonamine
Generic name	Meclizine Hydrochloride
Dosage strength and form	25 mg Chewable Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Anti-emetic/ Anti-vertigo
Registration number	DR-XY22028
Manufacturer	Interphil Laboratories, Inc.

Bonamine: OTC forms registered under the Bonamine brand (capsule availability)

OTC capsule form registered under the Bonamine brand: NO. 2 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Bonamine	12.5 mg Chewable Tablet	Over-the-Counter (OTC) Drug	DR-XY19867
Bonamine	25 mg Chewable Tablet	Over-the-Counter (OTC) Drug	DR-XY22028

Bonamine: Official leaflet (Package Insert / Patient Information Leaflet)

An official leaflet for this registration is published on the FDA Verification Portal: PIL_01 (https://drive.google.com/file/d/1cmQGJdB0wRZiOZUKiTtT_dNxazpRGIQ2/view?usp=drivesdk). Its text (OCR) and the original leaflet pages follow.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Bonamine"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Bonamine: Official leaflet text (OCR transcription)

Machine transcription of the official FDA leaflet for Reg. No. DR-XY22028. OCR may contain errors; the original leaflet pages are attached after this section.

Bonamine leaflet PIL_01

CX)
<:I'
.-I
Item Code: Item Description:
Bonamine 25mg Chewable Tablet
Material Code: Product Information Leaflet
Revision(s): Dimension:
Revision No.: 148 x 148 mm
Note Dimensions are in millimeters unless otherwise noted.
Tolerances are+ 1 mm
-I
1 1111111 11111111111 II 11111 1111
C90Z
I Meclizine HCI I
Bonamine®
25 mg Chewable Tablet
Anti-emetic/ Anti-vertigo
DESCRIPTION OF THE PRODUCT
Meclizine Hydrochloride (Bonamine®) Chewable Tablet is a pale orange, mottled, round, biconvex tablet, plain on one side and scored on the other side with characteristic sweet yogurt-like flavor.
WHAT IS IN THE MEDICINE?
Active Pharmaceutical Ingredient (API):
Meclizine Hydrochloride
Excipients used are:
Glycine, Corn Starch, Carmellose Calcium, Flavor, Magnesium Stearate, Sodium Saccharin, Colorant, Sodium Lauryl Sulfate, and Water.
STRENGTH OF THE MEDICINE
Meclizine Hydrochloride (Bonamine®) Chewable Tablet for Adults:
Each chewable tablet contains 25 mg of Meclizine Hydrochloride.
WHAT IS THIS MEDICINE USED FOR?
Meclizine Hydrochloride (Bonamine®) Chewable Tablet is used for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.
HOW MUCH AND HOW OFTEN SHOULD YOU USE THIS MEDICINE?
A daily dose should be taken one hour before travel starts for 24-hr protection.
The tablet may be chewed and swallowed with or without water or swallowed whole with water.
MECLIZINE HYDROCHLORIDE (BONAMINE®) 25 MG CHEWABLE TABLET FOR ADULTS
Age I
Recommended Dose
Adults and Children ;e 11 yrs. old I 1 to 2 tablets
Or, as prescribed by a physician. If additional dose is needed, interval should be 4 hours or longer.
Children 5-15 yrs old should be limited to two single doses only.
WHEN SHOULD YOU NOT TAKE THIS MEDICINE?
• A known hypersensitivity to Meclizine or any ingredient in this product.
• Patients taking monoamine oxidase inhibitor antidepressant
• Pregnant or breastfeeding mothers
CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE
• This drug may cause drowsiness. Take due precautions against driving a car or operating dangerous machinery.
• A doctor must always be consulted before giving Meclizine especially if the patient is taking other medications (antidepressant) and patients who are alcoholics or who regularly ingest alcohol in excess before giving this drug.
• It is also wise to remember to check the expiration date on the package or label of this product. Do not use after expiry date.

UNDESIRABLE EFFECTS OF THIS MEDICINE
Drowsiness, fatigue, dry mouth, constipation, headache, stomach upset, and rarely blurred vision have been reported with the use of Meclizine.
WHAT OTHER MEDICINE OR FOOD SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?
Avoid alcohol, sedatives, anti-allergic drugs and tranquilizers while taking Meclizine as this may increase the drowsiness effect.
WHAT SHOULD YOU DO IF YOU MISS A DOSE?
Do not double the dose. If you miss a dose, just take the next dose as scheduled (i.e., once a day) if needed to prevent and treat nausea, vomiting, or dizziness associated with motion sickness.
FRONT
CX)
<:I'
.-I
Supplier: Revision Date:
_j,&.CJl)Iti Printing Corp.
Tel. No.: 532-5870 • 532-0721 • 532-7645
Color Used: Graphics Artist:
Black
SIGNS AND SYMPTOMS OF OVERDOSE
Overdose of Meclizine may manifest as follows:
Drowsiness
Dry mouth
Fatigue
Rarely, blurred vision
WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN RECOMMENDED DOSAGE?
If more than the recommended dose has been given, consult a health professional immediately. Do not exceed recommended dosage.
HOW SHOULD YOU KEEP THIS MEDICINE?
Store at temperatures not exceeding 30°
C. Keep out of reach of
children.
WHEN SHOULD YOU CONSULT YOUR DOCTOR?
Pregnant or breastfeeding mothers, patients with increased pressure in the eye, asthma, emphysema, chronic pulmonary disease, difficulty in breathing and those with difficulty in urination due to enlargement of the prostate gland and those taking sedatives and tranquilizers should take Meclizine only under the direction of a doctor.
HOW SUPPLIED?
Meclizine Hydrochloride (Bonamine®) 25 mg Chewable Tablet for Adult: Box of 240's
NAME AND ADDRESS OF MARKETING AUTHORIZATION HOLDER
Taisho Pharmaceuticals (Philippines), Inc.
8th Floor CLIPP Center Building,
11th Avenue corner 39th Street,
Bonifacio Global City, Taguig 1634, Philippines
Under Authority of:
Taisho Pharmaceutical Co., Ltd.,
Tokyo, Japan
Trademark Owner
NAME AND ADDRESS OF MANUFACTURER
Interphil Laboratories, Inc.
Canlubang Industrial Estate,
Bo. Pittland, Cabuyao, Laguna
ADR REPORTING STATEMENT
For suspected adverse drug reaction, report to FDA: www.fda.gov.ph
Seek medical attention immediately at the first sign of any adverse drug reaction .
REGISTRATION NUMBER
Meclizine Hydrochloride (Bonamine®) 25 mg Chewable Tablet for Adult: DR-XY22028
RENEWAL OF THE AUTHORIZATION

Adult: 11 September 2023
DATE OF REVISION OF THE PATIENT INFORMATION LEAFLET (PIL)
September 2023
@-1 1023
BACK
-I

Item Code:	Item Description: **Bonamine 25mg Chewable Tablet** Product Information Leaflet		Supplier: JARDIS *Printing Corp.* Tel. No.: 532-5870 • 532-0721 • 532-7645	Revision Date:
Material Code:				
Revision No.:	Revision(s):	Dimension: 148 x 148 mm Note: Dimensions are in millimeters unless otherwise noted. Tolerances are ± 1 mm.	Color Used: Black	Graphics Artist:

148 mm

148 mm

C9●Z

Meclizine HCl

Bonamine®

25 mg Chewable Tablet
Anti-emetic / Anti-vertigo

DESCRIPTION OF THE PRODUCT

Meclizine Hydrochloride (Bonamine®) Chewable Tablet is a pale orange, mottled, round, biconvex tablet, plain on one side and scored on the other side with characteristic sweet yogurt-like flavor.

WHAT IS IN THE MEDICINE?

Active Pharmaceutical Ingredient (API):
Meclizine Hydrochloride

Excipients used are:
Glycine, Corn Starch, Carmellose Calcium, Flavor, Magnesium Stearate, Sodium Saccharin, Colorant, Sodium Lauryl Sulfate, and Water.

STRENGTH OF THE MEDICINE

Meclizine Hydrochloride (Bonamine®) Chewable Tablet for Adults:
Each chewable tablet contains 25 mg of Meclizine Hydrochloride.

WHAT IS THIS MEDICINE USED FOR?

Meclizine Hydrochloride (Bonamine®) Chewable Tablet is used for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.

HOW MUCH AND HOW OFTEN SHOULD YOU USE THIS MEDICINE?

A daily dose should be taken one hour before travel starts for 24-hr protection.

The tablet may be chewed and swallowed with or without water or swallowed whole with water.

MECLIZINE HYDROCHLORIDE (BONAMINE®) 25 MG CHEWABLE TABLET FOR ADULTS	
Age	**Recommended Dose**
Adults and Children ≥ 11 yrs. old	1 to 2 tablets
Or, as prescribed by a physician. If additional dose is needed, interval should be 4 hours or longer. Children 5-15 yrs old should be limited to two single doses only.	

WHEN SHOULD YOU NOT TAKE THIS MEDICINE?

- A known hypersensitivity to Meclizine or any ingredient in this product.
- Patients taking monoamine oxidase inhibitor antidepressant
- Pregnant or breastfeeding mothers

CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE

- This drug may cause drowsiness. Take due precautions against driving a car or operating dangerous machinery.
- A doctor must always be consulted before giving Meclizine especially if the patient is taking other medications (antidepressant) and patients who are alcoholics or who regularly ingest alcohol in excess before giving this drug.
- It is also wise to remember to check the expiration date on the package or label of this product. Do not use after expiry date.

UNDESIRABLE EFFECTS OF THIS MEDICINE

Drowsiness, fatigue, dry mouth, constipation, headache, stomach upset, and rarely blurred vision have been reported with the use of Meclizine.

WHAT OTHER MEDICINE OR FOOD SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?

Avoid alcohol, sedatives, anti-allergic drugs and tranquilizers while taking Meclizine as this may increase the drowsiness effect.

WHAT SHOULD YOU DO IF YOU MISS A DOSE?

Do not double the dose. If you miss a dose, just take the next dose as scheduled (i.e., once a day) if needed to prevent and treat nausea, vomiting, or dizziness associated with motion sickness.

148 mm

FRONT

148 mm

148 mm

SIGNS AND SYMPTOMS OF OVERDOSE

Overdose of Meclizine may manifest as follows:
Drowsiness
Dry mouth
Fatigue
Rarely, blurred vision

WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN RECOMMENDED DOSAGE?

If more than the recommended dose has been given, consult a health professional immediately. Do not exceed recommended dosage.

HOW SHOULD YOU KEEP THIS MEDICINE?

Store at temperatures not exceeding 30°C. Keep out of reach of children.

WHEN SHOULD YOU CONSULT YOUR DOCTOR?

Pregnant or breastfeeding mothers, patients with increased pressure in the eye, asthma, emphysema, chronic pulmonary disease, difficulty in breathing and those with difficulty in urination due to enlargement of the prostate gland and those taking sedatives and tranquilizers should take Meclizine only under the direction of a doctor.

HOW SUPPLIED?

Meclizine Hydrochloride (Bonamine®) 25 mg Chewable Tablet for Adult: Box of 240's

NAME AND ADDRESS OF MARKETING AUTHORIZATION HOLDER

Taisho Pharmaceuticals (Philippines), Inc.
8th Floor CLIPP Center Building,
11th Avenue corner 39th Street,
Bonifacio Global City, Taguig 1634, Philippines

Under Authority of:
Taisho Pharmaceutical Co., Ltd.,
Tokyo, Japan
Trademark Owner

NAME AND ADDRESS OF MANUFACTURER

Interphil Laboratories, Inc.
Canlubang Industrial Estate,
Bo. Pittland, Cabuyao, Laguna

ADR REPORTING STATEMENT

For suspected adverse drug reaction, report to FDA: www.fda.gov.ph
Seek medical attention immediately at the first sign of any adverse drug reaction.

REGISTRATION NUMBER

Meclizine Hydrochloride (Bonamine®) 25 mg Chewable Tablet for Adult: DR-XY22028

RENEWAL OF THE AUTHORIZATION

Meclizine Hydrochloride (Bonamine®) 25 mg Chewable Tablet for Adult: 11 September 2023

DATE OF REVISION OF THE PATIENT INFORMATION LEAFLET (PIL)
September 2023

-1 1023

148 mm

BACK

Buscopan (Hyoscine N-Butylbromide)

10 mg Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY41205

Buscopan: Official FDA Philippines registration record

Brand name	Buscopan
Generic name	Hyoscine N-Butylbromide
Dosage strength and form	10 mg Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antispasmodic
Registration number	DR-XY41205
Manufacturer	PT. BOEHRINGER INGELHEIM INDONESIA

Buscopan: OTC forms registered under the Buscopan brand (capsule availability)

OTC capsule form registered under the Buscopan brand: NO. 1 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Buscopan	10 mg Tablet	Over-the-Counter (OTC) Drug	DR-XY41205

Buscopan: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY41205 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Buscopan"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Canesten (Clotrimazole)

100 mg Vaginal Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY47618

Canesten: Official FDA Philippines registration record

Brand name	Canesten
Generic name	Clotrimazole
Dosage strength and form	100 mg Vaginal Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antifungal
Registration number	DR-XY47618
Manufacturer	Haupt Pharma Wülfing GmbH

Canesten: OTC forms registered under the Canesten brand (capsule availability)

OTC capsule form registered under the Canesten brand: NO. 4 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Canesten	10 mg/g (1% w/w) Cream	Over-the-Counter (OTC) Drug	DR-XY42700
Canesten	10 mg/mL (1%) Topical Solution	Over-the-Counter (OTC) Drug	DR-8992
Canesten	100 mg Vaginal Tablet	Over-the-Counter (OTC) Drug	DR-XY32759
Canesten	100 mg Vaginal Tablet	Over-the-Counter (OTC) Drug	DR-XY47618

Canesten: Official leaflet (Package Insert / Patient Information Leaflet)

An official leaflet for this registration is published on the FDA Verification Portal: PIL_01 (https://drive.google.com/file/d/1ZRjRz4L9_JyYUokUHsZAEUdFkgnwerbS/view?usp=drivesdk). Its text (OCR) and the original leaflet pages follow.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Canesten"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Canesten: Official leaflet text (OCR transcription)

Machine transcription of the official FDA leaflet for Reg. No. DR-XY47618. OCR may contain errors; the original leaflet pages are attached after this section.

Canesten leaflet PIL_01

1. Name of the Product
Clotrimazole (Canesten®) 100 mg Vaginal Tablet
2. Description of the Product
Vaginal tablet contains Clotrimazole which acts against fungi by inhibiting ergosterol synthesis. Inhibition of ergosterol synthesis leads to
structural and functional impairment of the fungal cytoplasmic membrane.
3. What is in the Medicine?
100 mg Vaginal Tablet:
Each vaginal tablet contains:
Clotrimazole…….100 mg
4. Strength of the Medicine
100 mg Vaginal Tablet: Clotrimazole 100 mg
5. What is this Medicine used for?
Infections of the genital region (vaginitis) caused by fungi (usually Candida), and superinfections caused by clotrimazole-sensitive bacteria.
6. How much and how often should you use this Medicine?
The vaginal tablet should be inserted as deeply as possible into the vagina in the evening before going to bed. Insertion is best achieved
when lying back with the legs slightly drawn up.
Clotrimazole vaginal tablets need moisture in the vagina to dissolve completely, otherwise undissolved pieces of the vaginal tablet might
crumble out of the vagina. To prevent this it is important to insert the medication as deeply as possible into the vagina at bedtime.
Should the vaginal tablet not dissolve completely within one night, the use of a vaginal cream should be considered.
For treatment of vaginal infections:
6-day treatment with Clotrimazole (Canesten®) 100 mg:
1 vaginal tablet to be introduced each evening on 6 successive days.
Generally:
If symptoms persist, the patient should consult a doctor.
The treatment can be repeated if necessary, however, recurrent infections may indicate an underlying medical cause.
Patient should seek medical advice if symptoms return within 2 months.
If the labia and adjacent areas are simultaneously infected, local treatment with an external cream should also be given in addition to the intravaginal treatment (combination treatment).
Intended for use by adults and children 12 years of age and older.
7. When should you not take this Medicine?
Hypersensitivity to the active substance or excipients (Calcium Lactate pentahydrate, Cellulose microcrystalline, Crospovidone, Hypromellose 15cP, Lactic Acid, Lactose monohydrate, Magnesium stearate, Maize starch, Silica colloidal anhydrous)
8. Care that should be taken when taking this Medicine?
If the patient has a fever (temperature of 38°C or above), lower abdominal pain, back pain, foul smelling vaginal discharge, nausea, vaginal haemorrhage, and/or associated shoulder pain the patient should consult a doctor.
Treatment during the menstrual period should not be performed. The treatment should be finished before the onset of menstruation.
Do not use tampons, intravaginal douches, spermicides or other vaginal products while using this product.
Avoidance of vaginal intercourse is recommended in case of vaginal infection and while using this product, the partner could become
infected. The sexual partner should also undergo local treatment if symptoms, e.g. pruritus, inflammation, etc. are present.
When used in pregnancy, the vaginal tablets should be inserted without using an applicator.
Avoid contact with eyes. Do not swallow.
100 mg Vaginal Tablet
_1.5 laetus 143
143 laetus_1.5
143 laetus_1.5 90193877_p02.indd 1 18.09.23 11:40
9. Undesirable Effects of this Medicine
Immune system disorders: anaphylactic reaction, angioedema, hypersensitivity
Vascular disorder: syncope, hypotension
Respiratory, thoracic and mediastinal disorders: dyspnea
Reproductive system and breast disorders: vaginal exfoliation, vaginal discharge, vaginal haemorrhage, vulvovaginal discomfort, vulvovaginal erythema, vulvovaginal burning sensation, vulvovaginal pruritus, vulvovaginal pain
Gastrointestinal disorders: abdominal pain, nausea
General disorders and administration site conditions: application site irritation, edema, pain
10.What other medicine or food should be avoided while taking this Medicine?

Concomitant medication with vaginal clotrimazole and oral tacrolimus (FK-506; immunosuppressant) might lead to increased tacrolimus
plasma levels and similarly with sirolimus. Patients should thus be thoroughly monitored for symptoms of tacrolimus or sirolimus overdosage, if necessary by determination of the respective plasma levels.
11. What should you do if you miss a dose?
No available information.
12. Signs and Symptoms of Overdose
No risk of acute intoxication is seen as it is unlikely to occur following a single vaginal or dermal application of an overdose (application over a large area under conditions favorable to absorption) or inadvertent oral ingestion.
13. What to do when you have taken more than the recommended dosage?
There is no specific antidote.
14. How should you keep this Medicine?
Store at temperatures not exceeding 30ºC.
Keep medicine out of the reach of children.
15. When should you consult your doctor?
If symptoms persist for more than 7 days, the patient may have a medical condition that requires treatment by a doctor.
Patient should seek medical advice if symptoms return within 2 months.
If the patient has a fever (temperature of 38°C or above), lower abdominal pain, back pain, foul smelling vaginal discharge, nausea, vaginal haemorrhage, and/or associated shoulder pain the patient should consult a doctor.
Manufacturer:
Haupt Pharma Wülfing GmbH
Bethelner Landstrasse 18, 31028 Gronau/Leine, Germany
Packed by:
GP Grenzach Produktions GmbH
Emil-Barell-Strasse 7, 79639 Grenzach-Wyhlen, Germany
Marketing Authorization Holder:
Bayer Philippines, Inc
8th Floor Science Hub Tower 1, Campus Avenue
Corner Turin St., McKinley Hill Cyberpark,
Pinagsama, Taguig City
If you want to report a product complaint or side effect, please contact your health care professional
or the Philippine FDA at adr@fda.gov.ph
Inquiries can also be directed to:
Bayer Philippines, Inc.
Taguig City, Philippines
E-mail: MedInfoCH-PH@bayer.com
drugsafety.philippines@bayer.com
Registration Number:
DR-XY47618
Date of First Authorization:
01 March 1976
Date of Revision of PIL:
November 2022
(Based on CCDS version 5 dated 28-Oct-2020)
90193877
_1.5 laetus 143
143 laetus_1.5
143 laetus_1.5 90193877_p02.indd 2 18.09.23 11:40

I 143 I laetus_1.5

Clotrimazole Canesten®

100 mg Vaginal Tablet

1. **Name of the Product**

   Clotrimazole (Canesten®) 100 mg Vaginal Tablet

2. **Description of the Product**

   Vaginal tablet contains Clotrimazole which acts against fungi by inhibiting ergosterol synthesis. Inhibition of ergosterol synthesis leads to structural and functional impairment of the fungal cytoplasmic membrane.

3. **What is in the Medicine?**

   100 mg Vaginal Tablet:
   Each vaginal tablet contains:
   Clotrimazole.....100 mg

4. **Strength of the Medicine**

   100 mg Vaginal Tablet: Clotrimazole 100 mg

5. **What is this Medicine used for?**

   Infections of the genital region (vaginitis) caused by fungi (usually *Candida*), and superinfections caused by clotrimazole-sensitive bacteria.

6. **How much and how often should you use this Medicine?**

   The vaginal tablet should be inserted as deeply as possible into the vagina in the evening before going to bed. Insertion is best achieved when lying back with the legs slightly drawn up.

   Clotrimazole vaginal tablets need moisture in the vagina to dissolve completely, otherwise undissolved pieces of the vaginal tablet might crumble out of the vagina. To prevent this it is important to insert the medication as deeply as possible into the vagina at bedtime.
   Should the vaginal tablet not dissolve completely within one night, the use of a vaginal cream should be considered.

   For treatment of vaginal infections:

   6-day treatment with Clotrimazole (Canesten®) 100mg:

   1 vaginal tablet to be introduced each evening on 6 successive days.

   Generally:
   If symptoms persist, the patient should consult a doctor.

   The treatment can be repeated if necessary, however, recurrent infections may indicate an underlying medical cause.
   Patient should seek medical advice if symptoms return within 2 months.

   If the labia and adjacent areas are simultaneously infected, local treatment with an external cream should also be given in addition to the intravaginal treatment (combination treatment).
   Intended for use by adults and children 12 years of age and older.

7. **When should you not take this Medicine?**

   Hypersensitivity to the active substance or excipients (Calcium Lactate pentahydrate, Cellulose microcrystalline, Crospovidone, Hypromellose 15cP, Lactic Acid, Lactose monohydrate, Magnesium stearate, Maize starch, Silica colloidal anhydrous)

8. **Care that should be taken when taking this Medicine?**

   If the patient has a fever (temperature of 38°C or above), lower abdominal pain, back pain, foul smelling vaginal discharge, nausea, vaginal haemorrhage, and/or associated shoulder pain the patient should consult a doctor.

   Treatment during the menstrual period should not be performed. The treatment should be finished before the onset of menstruation.

   Do not use tampons, intravaginal douches, spermicides or other vaginal products while using this product.

   Avoidance of vaginal intercourse is recommended in case of vaginal infection and while using this product, the partner could become infected. The sexual partner should also undergo local treatment if symptoms, e.g. pruritus, inflammation, etc. are present.

   When used in pregnancy, the vaginal tablets should be inserted without using an applicator.

   Avoid contact with eyes. Do not swallow.

I 143 laetus_1.5
90193877_p02.indd 1 18.09.23 11:40

I 143 I laetus_1.5

9. Undesirable Effects of this Medicine

Immune system disorders: anaphylactic reaction, angioedema, hypersensitivity

Vascular disorder: syncope, hypotension

Respiratory, thoracic and mediastinal disorders: dyspnea

Reproductive system and breast disorders: vaginal exfoliation, vaginal discharge, vaginal haemorrhage, vulvovaginal discomfort, vulvovaginal erythema, vulvovaginal burning sensation, vulvovaginal pruritus, vulvovaginal pain

Gastrointestinal disorders: abdominal pain, nausea

General disorders and administration site conditions: application site irritation, edema, pain

10. What other medicine or food should be avoided while taking this Medicine?

Concomitant medication with vaginal clotrimazole and oral tacrolimus (FK-506; immunosuppressant) might lead to increased tacrolimus plasma levels and similarly with sirolimus. Patients should thus be thoroughly monitored for symptoms of tacrolimus or sirolimus overdosage, if necessary by determination of the respective plasma levels.

11. What should you do if you miss a dose?

No available information.

I 143 I laetus_1.5

12. Signs and Symptoms of Overdose

No risk of acute intoxication is seen as it is unlikely to occur following a single vaginal or dermal application of an overdose (application over a large area under conditions favorable to absorption) or inadvertent oral ingestion.

13. What to do when you have taken more than the recommended dosage?

There is no specific antidote.

14. How should you keep this Medicine?

Store at temperatures not exceeding 30°C.
Keep medicine out of the reach of children.

15. When should you consult your doctor?

If symptoms persist for more than 7 days, the patient may have a medical condition that requires treatment by a doctor.

Patient should seek medical advice if symptoms return within 2 months.

If the patient has a fever (temperature of 38°C or above), lower abdominal pain, back pain, foul smelling vaginal discharge, nausea, vaginal haemorrhage, and/or associated shoulder pain the patient should consult a doctor.

Manufacturer:
Haupt Pharma Wülfing GmbH
Bethelner Landstrasse 18, 31028 Gronau/Leine, Germany

Packed by:
GP Grenzach Produktions GmbH
Emil-Barell-Strasse 7, 79639 Grenzach-Wyhlen, Germany

Marketing Authorization Holder:
Bayer Philippines, Inc
8th Floor Science Hub Tower 1, Campus Avenue
Corner Turin St., McKinley Hill Cyberpark,
Pinagsama, Taguig City

If you want to report a product complaint or side effect, please contact your health care professional or the Philippine FDA at adr@fda.gov.ph

Inquiries can also be directed to:
Bayer Philippines, Inc.
Taguig City, Philippines
E-mail: MedInfoCH-PH@bayer.com
drugsafety.philippines@bayer.com

Registration Number:
DR-XY47618

Date of First Authorization:
01 March 1976

Date of Revision of PIL:
November 2022
(Based on CCDS version 5 dated 28-Oct-2020)

I 143 I laetus_1.5

Bayer
90193877

I 143 I laetus_1.5
90193877_p02.indd 2 18.09.23 11:40

Canesten (Clotrimazole)

10 mg/mL (1%) Topical Solution | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-8992

Canesten: Official FDA Philippines registration record

Brand name	Canesten
Generic name	Clotrimazole
Dosage strength and form	10 mg/mL (1%) Topical Solution
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antifungal
Registration number	DR-8992
Manufacturer	KVP Pharma + Veterinar Produkte GmbH

Canesten: OTC forms registered under the Canesten brand (capsule availability)

OTC capsule form registered under the Canesten brand: NO. 4 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Canesten	10 mg/g (1% w/w) Cream	Over-the-Counter (OTC) Drug	DR-XY42700
Canesten	10 mg/mL (1%) Topical Solution	Over-the-Counter (OTC) Drug	DR-8992
Canesten	100 mg Vaginal Tablet	Over-the-Counter (OTC) Drug	DR-XY32759
Canesten	100 mg Vaginal Tablet	Over-the-Counter (OTC) Drug	DR-XY47618

Canesten: Official leaflet (Package Insert / Patient Information Leaflet)

An official leaflet for this registration is published on the FDA Verification Portal: PIL_01 (https://drive.google.com/file/d/12Xbcr0Mvgc7trgAVrhMG-lKVRsxyWsKT/view?usp=drivesdk). Its text (OCR) and the original leaflet pages follow.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Canesten"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Canesten: Official leaflet text (OCR transcription)

Machine transcription of the official FDA leaflet for Reg. No. DR-8992. OCR may contain errors; the original leaflet pages are attached after this section.

Canesten leaflet PIL_01

1. Name of the Product
Clotrimazole (Canesten®)
2. Description of the Product
Clotrimazole (Canesten®) appears as 10 mg/g (1% w/w) Cream and 10 mg/mL (1%) Topical Solution which is categorized as antifungal preparations for external use.
3. What is in the Medicine?
Cream:
Each gram of cream contains:
Clotrimazole.....10 mg
Topical Solution:
Each mL of topical solution contains:
4. Strength of the Medicine
Cream: 10 mg/g (1% w/w)
Topical Solution: 10 mg/mL (1%)
5. What is this Medicine used for?
For the treatment of dermatomycoses caused by fungi, yeasts and molds:
1. Athlete's foot (alipunga)
2. Hand fungus
3. Ringworm (buni)
4. Jock itch (hadhad)
5. Pityriasis versicolor (an-an)
6. Other skin fungal infections (cutaneous candidiasis, erythrasma)
6. How much and how often should you use this Medicine? How long is the treatment duration?
Cream:
Apply thinly 2 to 3 times a day and gently rub in. A strip of cream (1/2 cm long) is enough to treat an area about the size of the hand.
Topical Solution:
in. A few drops are enough to treat an area about the size of the hand.
To ensure complete healing depending on the indication, the treatment should be continued even if symptoms disappear according to the duration of treatment as specified below:
Pityriasis versicolor: 1 to 3 weeks
Candida vulvitis and
Candida balanitis: 1 to 2 weeks
Patients should consult their physician if there is no improvement after the recommended duration of treatment.
7. When should you not take this Medicine?
People who are hypersensitive to any ingredient in the preparation.
8. Care that should be taken when taking this Medicine?
Keep medicine out of the reach of children.
Avoid contact with eyes. Do not swallow.
Cream may reduce the effectiveness and safety of latex products such as condoms and diaphragms when applied on the genital area (women: labia and adjacent area of the vulva; men: prepuce and glans of the penis). The effect is temporary and occurs only during

treatment.
9. Undesirable Effects of this Medicine
Immune system disorders: angioedema, anaphylactic reaction, hypersensitivity.
10 mg/g (1% w/w) Cream
10 mg/mL (1%) Topical Solution
Antifungal
Respiratory, thoracic and mediastinal disorders: dyspnea
Skin and subcutaneous tissue disorders: blisters, contact dermatitis, erythema, paraesthesia, skin exfoliation, pruritus, rash, urticaria, stinging skin/burning sensation skin.
General disorders and administration site conditions: application site irritation, application site reaction, edema, pain.
10. What other medicine or food should be avoided while taking this Medicine?
None known.
11. What should you do if you miss a dose?
No available information.
12. Signs and Symptoms of Overdose
unlikely to occur following a single dermal application of an overdose (application over a large area under conditions favorable to absorption) or inadvertent oral ingestion.
13. What to do when you have taken more than the recommended dosage?
There is no specific antidote however patients should still consult their physician.
14. How should you keep this Medicine?
Cream: Store below 30°C
Topical Solution: Store below 25°C
15. When should you consult your doctor?
Patients should consult their physician if there is no improvement after 4 weeks of treatment.
Manufacturer:
Cream
Encube Ethicals Private Limited
Plot No. C-1, Madkaim Industrial Estate, Madkaim, Post: Mardol, Ponda, Goa 403404, India
Topical Solution
KVP Pharma + Veterinar Produkte GmbH Kiel, Germany
Projensdorfer Strasse 324 24106 Kiel,
Marketing Authorization Holder:
Bayer Philippines, Inc.
8th Floor Science Hub Tower 1, Campus Ave. Cor. Turin St., McKinley Hill Cyberpark, Pinagsama, Taguig City
Pregnancy
While there is a limited amount of clinical data in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity. Clotrimazole (Canesten®) can be used during pregnancy, but only under the direction of a health care professional.
Lactation
There are no data on the excretion of Clotrimazole (Canesten®) into human milk. However, systemic absorption is minimal after topical administration and is unlikely to lead to systemic effects. Clotrimazole

(Canesten®) may be used during lactation. If used topically on the nipple area, wash breast before feeding child.
If you want to report a product complaint or side effect, please contact your health care professional or the Philippine FDA at adr@fda.gov.ph.
Inquiries can also be directed to:
Bayer Philippines, Inc.
Taguig City, Philippines
E-mail: MedInfoCH-PH@bayer.com
drugsafety.philippines@bayer.com
Registration Number:
Cream: DR-XY42700
Topical Solution: DR-8992
Date of First Authorization:
Cream: 30 September 2013
Topical Solution: 11 July 1973
Date of Revision of PIL:
November 2022 (Based from CCDS v5 dated 15 October 2020)
Vascular disorders: hypotension, syncope
CPH5/03
N/A
86658291 86658313 86658348

148 mm

210 mm

Clotrimazole Canesten®

10 mg/g (1% w/w) Cream
10 mg/mL (1%) Topical Solution
Antifungal

1. Name of the Product
Clotrimazole (Canesten®)

2. Description of the Product
Clotrimazole (Canesten®) appears as 10 mg/g (1% w/w) Cream and 10 mg/mL (1%) Topical Solution which is categorized as antifungal preparations for external use.

3. What is in the Medicine?
Cream:
Each gram of cream contains:
Clotrimazole.....10 mg

Topical Solution:
Each mL of topical solution contains:
Clotrimazole.....10 mg

4. Strength of the Medicine
Cream: 10 mg/g (1% w/w)
Topical Solution: 10 mg/mL (1%)

5. What is this Medicine used for?
For the treatment of dermatomycoses caused by fungi, yeasts and molds:
1. Athlete's foot (alipunga)
2. Hand fungus
3. Ringworm (buni)
4. Jock itch (hadhad)
5. *Pityriasis versicolor* (an-an)
6. Other skin fungal infections (cutaneous candidiasis, erythrasma)

6. How much and how often should you use this Medicine? How long is the treatment duration?
Cream:
Apply thinly 2 to 3 times a day and gently rub in. A strip of cream (1/2 cm long) is enough to treat an area about the size of the hand.

Topical Solution:
Apply thinly 2 to 3 times a day and gently rub in. A few drops are enough to treat an area about the size of the hand.

To ensure complete healing depending on the indication, the treatment should be continued even if symptoms disappear according to the duration of treatment as specified below:

Dermatomycoses:	3 to 4 weeks
Erythrasma:	2 to 4 weeks
Pityriasis versicolor:	1 to 3 weeks
Candida vulvitis and Candida balanitis:	1 to 2 weeks

Patients should consult their physician if there is no improvement after the recommended duration of treatment.

7. When should you not take this Medicine?
People who are hypersensitive to any ingredient in the preparation.

8. Care that should be taken when taking this Medicine?
Keep medicine out of the reach of children. Avoid contact with eyes. Do not swallow.

Cream may reduce the effectiveness and safety of latex products such as condoms and diaphragms when applied on the genital area (women: labia and adjacent area of the vulva; men: prepuce and glans of the penis). The effect is temporary and occurs only during treatment.

Pregnancy
While there is a limited amount of clinical data in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity.

148 mm

210 mm

Clotrimazole (Canesten®) can be used during pregnancy, but only under the direction of a health care professional.

Lactation
There are no data on the excretion of Clotrimazole (Canesten®) into human milk. However, systemic absorption is minimal after topical administration and is unlikely to lead to systemic effects. Clotrimazole (Canesten®) may be used during lactation. If used topically on the nipple area, wash breast before feeding child.

9. Undesirable Effects of this Medicine
Immune system disorders: angioedema, anaphylactic reaction, hypersensitivity.

Vascular disorders: hypotension, syncope

Respiratory, thoracic and mediastinal disorders: dyspnea

Skin and subcutaneous tissue disorders: blisters, contact dermatitis, erythema, paraesthesia, skin exfoliation, pruritus, rash, urticaria, stinging skin/burning sensation skin.

General disorders and administration site conditions: application site irritation, application site reaction, edema, pain.

10. What other medicine or food should be avoided while taking this Medicine?
None known.

11. What should you do if you miss a dose?
No available information.

12. Signs and Symptoms of Overdose
No risk of acute intoxication is seen as it is unlikely to occur following a single dermal application of an overdose (application over a large area under conditions favorable to absorption) or inadvertent oral ingestion.

13. What to do when you have taken more than the recommended dosage?
There is no specific antidote however patients should still consult their physician.

14. How should you keep this Medicine?
Cream: Store below 30°C
Topical Solution: Store below 25°C

15. When should you consult your doctor?
Patients should consult their physician if there is no improvement after 4 weeks of treatment.

Manufacturer:
Cream
Encube Ethicals Private Limited
Plot No. C-1, Madkaim Industrial Estate, Madkaim, Post: Mardol, Ponda, Goa 403404, India

Topical Solution
KVP Pharma + Veterinar Produkte GmbH Kiel, Germany
Projensdorfer Strasse 324 24106 Kiel, Germany

Marketing Authorization Holder:
Bayer Philippines, Inc.
8th Floor Science Hub Tower 1, Campus Ave.
Cor. Turin St., McKinley Hill Cyberpark,
Pinagsama, Taguig City

If you want to report a product complaint or side effect, please contact your health care professional or the Philippine FDA at adr@fda.gov.ph.
Inquiries can also be directed to:
Bayer Philippines, Inc.
Taguig City, Philippines
E-mail: MedInfoCH-PH@bayer.com
drugsafety.philippines@bayer.com

Registration Number:
Cream: DR-XY42700
Topical Solution: DR-8992

Date of First Authorization:
Cream: 30 September 2013
Topical Solution: 11 July 1973

Date of Revision of PIL:
November 2022 (Based from CCDS v5 dated 15 October 2020)

Bayer
CPH5/03

Canesten (Clotrimazole)

10 mg/g (1% w/w) Cream | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY42700

Canesten: Official FDA Philippines registration record

Brand name	Canesten
Generic name	Clotrimazole
Dosage strength and form	10 mg/g (1% w/w) Cream
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antifungal (Topical Imidazole)
Registration number	DR-XY42700
Manufacturer	Encube Ethicals Pvt. Ltd.

Canesten: OTC forms registered under the Canesten brand (capsule availability)

OTC capsule form registered under the Canesten brand: NO. 4 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Canesten	10 mg/g (1% w/w) Cream	Over-the-Counter (OTC) Drug	DR-XY42700
Canesten	10 mg/mL (1%) Topical Solution	Over-the-Counter (OTC) Drug	DR-8992
Canesten	100 mg Vaginal Tablet	Over-the-Counter (OTC) Drug	DR-XY32759
Canesten	100 mg Vaginal Tablet	Over-the-Counter (OTC) Drug	DR-XY47618

Canesten: Official leaflet (Package Insert / Patient Information Leaflet)

An official leaflet for this registration is published on the FDA Verification Portal: PIL_01 (https://drive.google.com/file/d/1T_bcnFHzVpzpNkhxgY8CkuIiiYx81AUo/view?usp=drivesdk). Its text (OCR) and the original leaflet pages follow.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Canesten"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Canesten: Official leaflet text (OCR transcription)

Machine transcription of the official FDA leaflet for Reg. No. DR-XY42700. OCR may contain errors; the original leaflet pages are attached after this section.

Canesten leaflet PIL_01

1. Name of the Product
Clotrimazole (Canesten®)
2. Description of the Product
Clotrimazole (Canesten®) appears as 10 mg/g (1% w/w) Cream and 10 mg/mL (1%) Topical Solution which is categorized as antifungal preparations for external use.
3. What is in the Medicine?
Cream:
Each gram of cream contains:
Clotrimazole.....10 mg
Topical Solution:
Each mL of topical solution contains:
4. Strength of the Medicine
Cream: 10 mg/g (1% w/w)
Topical Solution: 10 mg/mL (1%)
5. What is this Medicine used for?
For the treatment of dermatomycoses caused by fungi, yeasts and molds:
1. Athlete's foot (alipunga)
2. Hand fungus
3. Ringworm (buni)
4. Jock itch (hadhad)
5. Pityriasis versicolor (an-an)
6. Other skin fungal infections (cutaneous candidiasis, erythrasma)
6. How much and how often should you use this Medicine? How long is the treatment duration?
Cream:
Apply thinly 2 to 3 times a day and gently rub in. A strip of cream (1/2 cm long) is enough to treat an area about the size of the hand.
Topical Solution:
in. A few drops are enough to treat an area about the size of the hand.
To ensure complete healing depending on the indication, the treatment should be continued even if symptoms disappear according to the duration of treatment as specified below:
Pityriasis versicolor: 1 to 3 weeks
Candida vulvitis and
Candida balanitis: 1 to 2 weeks
Patients should consult their physician if there is no improvement after the recommended duration of treatment.
7. When should you not take this Medicine?
People who are hypersensitive to any ingredient in the preparation.
8. Care that should be taken when taking this Medicine?
Keep medicine out of the reach of children.
Avoid contact with eyes. Do not swallow.
Cream may reduce the effectiveness and safety of latex products such as condoms and diaphragms when applied on the genital area (women: labia and adjacent area of the vulva; men: prepuce and glans of the penis). The effect is temporary and occurs only during

treatment.
9. Undesirable Effects of this Medicine
Immune system disorders: angioedema, anaphylactic reaction, hypersensitivity.
10 mg/g (1% w/w) Cream
10 mg/mL (1%) Topical Solution
Antifungal
Respiratory, thoracic and mediastinal disorders: dyspnea
Skin and subcutaneous tissue disorders: blisters, contact dermatitis, erythema, paraesthesia, skin exfoliation, pruritus, rash, urticaria, stinging skin/burning sensation skin.
General disorders and administration site conditions: application site irritation, application site reaction, edema, pain.
10. What other medicine or food should be avoided while taking this Medicine?
None known.
11. What should you do if you miss a dose?
No available information.
12. Signs and Symptoms of Overdose
unlikely to occur following a single dermal application of an overdose (application over a large area under conditions favorable to absorption) or inadvertent oral ingestion.
13. What to do when you have taken more than the recommended dosage?
There is no specific antidote however patients should still consult their physician.
14. How should you keep this Medicine?
Cream: Store below 30°C
Topical Solution: Store below 25°C
15. When should you consult your doctor?
Patients should consult their physician if there is no improvement after 4 weeks of treatment.
Manufacturer:
Cream
Encube Ethicals Private Limited
Plot No. C-1, Madkaim Industrial Estate, Madkaim, Post: Mardol, Ponda, Goa 403404, India
Topical Solution
KVP Pharma + Veterinar Produkte GmbH Kiel, Germany
Projensdorfer Strasse 324 24106 Kiel,
Marketing Authorization Holder:
Bayer Philippines, Inc.
8th Floor Science Hub Tower 1, Campus Ave. Cor. Turin St., McKinley Hill Cyberpark, Pinagsama, Taguig City
Pregnancy
While there is a limited amount of clinical data in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity. Clotrimazole (Canesten®) can be used during pregnancy, but only under the direction of a health care professional.
Lactation
There are no data on the excretion of Clotrimazole (Canesten®) into human milk. However, systemic absorption is minimal after topical administration and is unlikely to lead to systemic effects. Clotrimazole

(Canesten®) may be used during lactation. If used topically on the nipple area, wash breast before feeding child.
If you want to report a product complaint or side effect, please contact your health care professional or the Philippine FDA at adr@fda.gov.ph.
Inquiries can also be directed to:
Bayer Philippines, Inc.
Taguig City, Philippines
E-mail: MedInfoCH-PH@bayer.com
drugsafety.philippines@bayer.com
Registration Number:
Cream: DR-XY42700
Topical Solution: DR-8992
Date of First Authorization:
Cream: 30 September 2013
Topical Solution: 11 July 1973
Date of Revision of PIL:
November 2022 (Based from CCDS v5 dated 15 October 2020)
Vascular disorders: hypotension, syncope
CPH5/03
N/A
86658291 86658313 86658348

148 mm

210 mm

Clotrimazole Canesten®

10 mg/g (1% w/w) Cream
10 mg/mL (1%) Topical Solution
Antifungal

1. Name of the Product
Clotrimazole (Canesten®)

2. Description of the Product
Clotrimazole (Canesten®) appears as 10 mg/g (1% w/w) Cream and 10 mg/mL (1%) Topical Solution which is categorized as antifungal preparations for external use.

3. What is in the Medicine?
Cream:
Each gram of cream contains:
Clotrimazole.....10 mg

Topical Solution:
Each mL of topical solution contains:
Clotrimazole.....10 mg

4. Strength of the Medicine
Cream: 10 mg/g (1% w/w)
Topical Solution: 10 mg/mL (1%)

5. What is this Medicine used for?
For the treatment of dermatomycoses caused by fungi, yeasts and molds:
1. Athlete's foot (alipunga)
2. Hand fungus
3. Ringworm (buni)
4. Jock itch (hadhad)
5. *Pityriasis versicolor* (an-an)
6. Other skin fungal infections (cutaneous candidiasis, erythrasma)

6. How much and how often should you use this Medicine? How long is the treatment duration?
Cream:
Apply thinly 2 to 3 times a day and gently rub in. A strip of cream (1/2 cm long) is enough to treat an area about the size of the hand.

Topical Solution:
Apply thinly 2 to 3 times a day and gently rub in. A few drops are enough to treat an area about the size of the hand.

To ensure complete healing depending on the indication, the treatment should be continued even if symptoms disappear according to the duration of treatment as specified below:

Dermatomycoses:	3 to 4 weeks
Erythrasma:	2 to 4 weeks
Pityriasis versicolor:	1 to 3 weeks
Candida vulvitis and Candida balanitis:	1 to 2 weeks

Patients should consult their physician if there is no improvement after the recommended duration of treatment.

7. When should you not take this Medicine?
People who are hypersensitive to any ingredient in the preparation.

8. Care that should be taken when taking this Medicine?
Keep medicine out of the reach of children. Avoid contact with eyes. Do not swallow.

Cream may reduce the effectiveness and safety of latex products such as condoms and diaphragms when applied on the genital area (women: labia and adjacent area of the vulva; men: prepuce and glans of the penis). The effect is temporary and occurs only during treatment.

Pregnancy
While there is a limited amount of clinical data in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity.

148 mm

210 mm

Clotrimazole (Canesten®) can be used during pregnancy, but only under the direction of a health care professional.

Lactation
There are no data on the excretion of Clotrimazole (Canesten®) into human milk. However, systemic absorption is minimal after topical administration and is unlikely to lead to systemic effects. Clotrimazole (Canesten®) may be used during lactation. If used topically on the nipple area, wash breast before feeding child.

9. Undesirable Effects of this Medicine
Immune system disorders: angioedema, anaphylactic reaction, hypersensitivity.

Vascular disorders: hypotension, syncope

Respiratory, thoracic and mediastinal disorders: dyspnea

Skin and subcutaneous tissue disorders: blisters, contact dermatitis, erythema, paraesthesia, skin exfoliation, pruritus, rash, urticaria, stinging skin/burning sensation skin.

General disorders and administration site conditions: application site irritation, application site reaction, edema, pain.

10. What other medicine or food should be avoided while taking this Medicine?
None known.

11. What should you do if you miss a dose?
No available information.

12. Signs and Symptoms of Overdose
No risk of acute intoxication is seen as it is unlikely to occur following a single dermal application of an overdose (application over a large area under conditions favorable to absorption) or inadvertent oral ingestion.

13. What to do when you have taken more than the recommended dosage?
There is no specific antidote however patients should still consult their physician.

14. How should you keep this Medicine?
Cream: Store below 30°C
Topical Solution: Store below 25°C

15. When should you consult your doctor?
Patients should consult their physician if there is no improvement after 4 weeks of treatment.

Manufacturer:
Cream
Encube Ethicals Private Limited
Plot No. C-1, Madkaim Industrial Estate, Madkaim, Post: Mardol, Ponda, Goa 403404, India

Topical Solution
KVP Pharma + Veterinar Produkte GmbH Kiel, Germany
Projensdorfer Strasse 324 24106 Kiel, Germany

Marketing Authorization Holder:
Bayer Philippines, Inc.
8th Floor Science Hub Tower 1, Campus Ave.
Cor. Turin St., McKinley Hill Cyberpark,
Pinagsama, Taguig City

If you want to report a product complaint or side effect, please contact your health care professional or the Philippine FDA at adr@fda.gov.ph.
Inquiries can also be directed to:
Bayer Philippines, Inc.
Taguig City, Philippines
E-mail: MedInfoCH-PH@bayer.com
drugsafety.philippines@bayer.com

Registration Number:
Cream: DR-XY42700
Topical Solution: DR-8992

Date of First Authorization:
Cream: 30 September 2013
Topical Solution: 11 July 1973

Date of Revision of PIL:
November 2022 (Based from CCDS v5 dated 15 October 2020)

Bayer
CPH5/03

Daktarin (Miconazole nitrate)

20 mg per 1 g cream | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY14359

Daktarin: Official FDA Philippines registration record

Brand name	Daktarin
Generic name	Miconazole nitrate
Dosage strength and form	20 mg per 1 g cream
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antifungal
Registration number	DR-XY14359
Manufacturer	Interphil Laboratories, Inc.

Daktarin: OTC forms registered under the Daktarin brand (capsule availability)

OTC capsule form registered under the Daktarin brand: NO. 2 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Daktarin	20 mg per 1 g cream	Over-the-Counter (OTC) Drug	DR-XY14359
Daktarin	20 mg/g Oral Gel	Over-the-Counter (OTC) Drug	DR-XY14361

Daktarin: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY14359 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Daktarin"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Daktarin (Miconazole)

20 mg/g Oral Gel | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY14361

Daktarin: Official FDA Philippines registration record

Brand name	Daktarin
Generic name	Miconazole
Dosage strength and form	20 mg/g Oral Gel
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antifungal
Registration number	DR-XY14361
Manufacturer	Interphil Laboratories, Inc

Daktarin: OTC forms registered under the Daktarin brand (capsule availability)

OTC capsule form registered under the Daktarin brand: NO. 2 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Daktarin	20 mg per 1 g cream	Over-the-Counter (OTC) Drug	DR-XY14359
Daktarin	20 mg/g Oral Gel	Over-the-Counter (OTC) Drug	DR-XY14361

Daktarin: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY14361 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Daktarin"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Diatabs (Loperamide Hydrochloride)

2 mg Capsule | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY24964

Diatabs: Official FDA Philippines registration record

Brand name	Diatabs
Generic name	Loperamide Hydrochloride
Dosage strength and form	2 mg Capsule
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antipropulsive
Registration number	DR-XY24964
Manufacturer	UNILAB, Inc.

Diatabs: OTC forms registered under the Diatabs brand (capsule availability)

OTC capsule form registered under the Diatabs brand: YES. 2 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Diatabs	2 mg Capsule	Over-the-Counter (OTC) Drug	DR-XY24964
Diatabs Advance	2 mg/ 125 mg Softgel Capsule	Over-the-Counter (OTC) Drug	DR-XY46907

Diatabs: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY24964 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Diatabs"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Diatabs Advance (Loperamide + Simeticone)

2 mg/ 125 mg Softgel Capsule | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY46907

Diatabs Advance: Official FDA Philippines registration record

Brand name	Diatabs Advance
Generic name	Loperamide + Simeticone
Dosage strength and form	2 mg/ 125 mg Softgel Capsule
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antidiarrheal/ Antiflatulent
Registration number	DR-XY46907
Manufacturer	Amherst Laboratories, Inc.

Diatabs Advance: OTC forms registered under the Diatabs brand (capsule availability)

OTC capsule form registered under the Diatabs brand: YES. 2 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Diatabs	2 mg Capsule	Over-the-Counter (OTC) Drug	DR-XY24964
Diatabs Advance	2 mg/ 125 mg Softgel Capsule	Over-the-Counter (OTC) Drug	DR-XY46907

Diatabs Advance: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY46907 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Diatabs"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Dulcolax (Bisacodyl)

5 mg Enteric-Coated Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY42393

Dulcolax: Official FDA Philippines registration record

Brand name	Dulcolax
Generic name	Bisacodyl
Dosage strength and form	5 mg Enteric-Coated Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Laxative
Registration number	DR-XY42393
Manufacturer	Sanofi Vietnam Shareholding Company

Dulcolax: OTC forms registered under the Dulcolax brand (capsule availability)

OTC capsule form registered under the Dulcolax brand: NO. 3 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Dulcolax	10 mg Rectal Suppository	Over-the-Counter (OTC) Drug	DR-XY44557
Dulcolax	5 mg Enteric-Coated Tablet	Over-the-Counter (OTC) Drug	DR-XY42393
Dulcolax	5 mg Suppository (Pediatric)	Over-the-Counter (OTC) Drug	DR-XY44940

Dulcolax: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY42393 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Dulcolax"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Dulcolax (Bisacodyl)

10 mg Rectal Suppository | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY44557

Dulcolax: Official FDA Philippines registration record

Brand name	Dulcolax
Generic name	Bisacodyl
Dosage strength and form	10 mg Rectal Suppository
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Laxative
Registration number	DR-XY44557
Manufacturer	Istituto De Angeli S.R.L.

Dulcolax: OTC forms registered under the Dulcolax brand (capsule availability)

OTC capsule form registered under the Dulcolax brand: NO. 3 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Dulcolax	10 mg Rectal Suppository	Over-the-Counter (OTC) Drug	DR-XY44557
Dulcolax	5 mg Enteric-Coated Tablet	Over-the-Counter (OTC) Drug	DR-XY42393
Dulcolax	5 mg Suppository (Pediatric)	Over-the-Counter (OTC) Drug	DR-XY44940

Dulcolax: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY44557 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Dulcolax"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Eye Mo Cool Red Eyes (Tetrahydrozoline Hydrochloride)

0.05% Solution (Eyedrops) | Household Remedy (HR) | FDA Reg. No. DRHR-2428

Eye Mo Cool Red Eyes: Official FDA Philippines registration record

Brand name	Eye Mo Cool Red Eyes
Generic name	Tetrahydrozoline Hydrochloride
Dosage strength and form	0.05% Solution (Eyedrops)
Classification	Household Remedy (HR)
Pharmacologic category	Vasoconstrictor
Registration number	DRHR-2428
Manufacturer	PT. Pharma Health Care

Eye Mo Cool Red Eyes: OTC forms registered under the Eye Mo brand (capsule availability)

OTC capsule form registered under the Eye Mo brand: NO. 2 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Eye Mo Cool Red Eyes	0.05% Solution (Eyedrops)	Household Remedy (HR)	DRHR-2428
Eye Mo Moisturizing Formula	3 mg/mL Ophthalmic Solution	Over-the-Counter (OTC) Drug	DR-XY35419

Eye Mo Cool Red Eyes: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-2428 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Eye Mo"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Eye Mo Moisturizing Formula (Hypromellose)

3 mg/mL Ophthalmic Solution | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY35419

Eye Mo Moisturizing Formula: Official FDA Philippines registration record

Brand name	Eye Mo Moisturizing Formula
Generic name	Hypromellose
Dosage strength and form	3 mg/mL Ophthalmic Solution
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Lubricant
Registration number	DR-XY35419
Manufacturer	PT. Pharma Health Care

Eye Mo Moisturizing Formula: OTC forms registered under the Eye Mo brand (capsule availability)

OTC capsule form registered under the Eye Mo brand: NO. 2 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Eye Mo Cool Red Eyes	0.05% Solution (Eyedrops)	Household Remedy (HR)	DRHR-2428
Eye Mo Moisturizing Formula	3 mg/mL Ophthalmic Solution	Over-the-Counter (OTC) Drug	DR-XY35419

Eye Mo Moisturizing Formula: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY35419 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Eye Mo"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Gaviscon (Sodium Alginate + Sodium Bicarbonate + Calcium Carbonate)

250 mg/ 133.5 mg/ 80 mg Chewable Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY35582

Gaviscon: Official FDA Philippines registration record

Brand name	Gaviscon
Generic name	Sodium Alginate + Sodium Bicarbonate + Calcium Carbonate
Dosage strength and form	250 mg/ 133.5 mg/ 80 mg Chewable Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antacid
Registration number	DR-XY35582
Manufacturer	Reckitt Benckiser Healthcare (UK) Limited

Gaviscon: OTC forms registered under the Gaviscon brand (capsule availability)

OTC capsule form registered under the Gaviscon brand: NO. 7 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Gaviscon Infant	225 mg/ 87.5 mg Powder for Oral Suspension	Over-the-Counter (OTC) Drug	DR-XY46801
Gaviscon	250 mg/ 133.5 mg/ 80 mg Chewable Tablet	Over-the-Counter (OTC) Drug	DR-XY35582
Gaviscon Double Action	250 mg/106.5 mg/187.5 mg Chewable Tablet	Over-the-Counter (OTC) Drug	DR-XY43658
Gaviscon Double Action	250 mg/106.5 mg/187.5 mg Chewable Tablet (Mixed Berries Flavor)	Over-the-Counter Drug (OTC)	DR-XY46902
Gaviscon	500 mg/ 267 mg/ 160 mg per 10 mL Oral Suspension	Over-the-Counter (OTC) Drug	DR-XY34746
Gaviscon	500 mg/ 267 mg/ 160 mg per 10 mL Oral suspension	Over-the-Counter (OTC) Drug	DR-XY34907
Gaviscon Double Action	500 mg/213 mg/325 mg per 10 mL Suspension	Over-the-Counter (OTC) Drug	DR-XY43337

Gaviscon: Official leaflet (Package Insert / Patient Information Leaflet)

An official leaflet for this registration is published on the FDA Verification Portal: PIL_01 (https://drive.google.com/file/d/1dY1NfdSS22QCCmE-kx6eVWrE8Z4yWmTD/view?usp=drivesdk). Its text (OCR) and the original leaflet pages follow.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Gaviscon"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Gaviscon: Official leaflet text (OCR transcription)

Machine transcription of the official FDA leaflet for Reg. No. DR-XY35582. OCR may contain errors; the original leaflet pages are attached after this section.

Gaviscon leaflet PIL_01

3. HOW TO TAKE SODIUM ALGINATE, SODIUM BICARBONATE, CALCIUM CARBONATE (GAVISCON) CHEWABLE TABLETS
For oral administration. Chew thoroughly before swallowing.
Adults, including the elderly and children 12 years and over: When symptoms occur take two to four tablets after meals and at bedtime, up to four times a day.
Children under 12 years: Should only be taken on medical advice.
If you forget a dose, it is not necessary to double the dose next time, just carry on taking as before.
If you take toomuch ofthis product youmay feel bloated and experience someabdominal discomfort. Itis unlikely to cause you any harm, but please consult your doctor or pharmacist.
After taking Sodium Alginate, Sodium Bicarbonate, Calcium Carbonate (Gaviscon) Chewable Tablets:
If symptoms persist after 7 days consult your doctor.
4. POSSIBLE SIDE EFFECTS
Very rare (lessthan 1 in 10,000) chance of an allergic reaction to the ingredients. Symptoms ofthismay include skin rash, itching, difficulty breathing, dizziness, orswelling of the face, lips,tongue or throat. If you experience these or any otherside-effects stop taking the product and consult your doctor immediately.
5. HOW TO STORE SODIUM ALGINATE, SODIUM BICARBONATE, CALCIUM CARBONATE (GAVISCON) CHEWABLE TABLETS
Keep all medicines out of the reach and sight of children.
Do not use after the expiry date shown on the pack. Do not store above 30oC.
Thismedicine contains 63.16 mg sodium(main component of cooking/table salt) in each tablet. Thisis equivalentto 3.2 % ofthe recommendedmaximumdailydietaryintakeofsodiumforan adult. Thismedicine also contains 3.2mmol of calcium per 4 tablet dose. The maximum recommended daily dose of this medicinal product contains 1012 mg sodium(found in table salt). Thisis equivalentto 50.6% of the adult recommendedmaximumdaily dietary intakefor sodium. Please also talk to your doctor regarding these salt contents if you suffer or have suffered from significant kidney or heart disease, as certain salts could interfere with these diseases. Talk to your pharmacist or doctor if you need this product on a daily basisfor a prolonged period of time, especiallyif you have been advised to have a low salt diet.
This medicine contains 3.75mg aspartame in each tablet. Aspartame is a source of phenylalanine. Itmay be harmful if you have phenylketonuria (PKU), a rare geneticdisorderinwhich phenylalanine builds up because the body cannot remove it properly.
Taking Other Medicines:
Do not take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines. This is especially important if you are taking antihistamines, antibiotics (tetracyclines and quinolones such as norfloxacin), iron preparation, antifungals such as ketoconazole, digoxin and beta blockers (for heart conditions), penicillamine (commonly for rheumatoid arthritis), glucocorticoid (for inflammatory and autoimmune disorders), neuroleptics(for mental illness), thyroid hormones, chloroquine (for malaria), estramustine (for prostate cancer) and bisphosphonates (for osteoporosis).
Pregnancy and breast-feeding:
You can take this productif you are pregnantor breast - feeding.Aswith all medicines,the treatment duration should be limited as much as possible.
6. FURTHER INFORMATION
SodiumAlginate, SodiumBicarbonate, CalciumCarbonate (Gaviscon)ChewableTablets are circulartabletswith an off white to cream, slightly motied appearance.
Each chewable tablet contains 250 mg sodiumalginate, 133.5 mg sodiumbicarbonate and 80 mg calciumcarbonate as the active ingredients. The otheringredients are mannitol,macrogol 20,000, copovidone, aspartame, peppermint flavouring, magnesium stearate and acesulfame potassium.
These tablets do not contain sugar or gluten.
Forsuspected adverse drug reaction,reportto the FDA: www.fda.gov.ph. Seekmedical atiention immediately atthe first sign of any adverse reaction.
Registration No.: DR-XY35582
Date of Last Revision: December 2023
Manufactured by: Reckiti Benckiser Healthcare (UK) Limited
Dansom Lane, Hull HU8 7DS UK
Imported by: Reckiti Benckiser Healthcare (Phils) Inc.
3/F Mead Johnson Bldg. 2309 Don Chino Roces Ave. Ext.,
Makati City, Philippines, 1231
GAVISCON
250 mg/ 133.50 mg/ 80 mg Chewable Tablet
1. WHAT IS SODIUM ALGINATE, SODIUM BICARBONATE, CALCIUM CARBONATE (GAVISCON) CHEWABLE TABLETS AND WHAT IS IT USED FOR?

This product belongsto a group of medicines called ‘reflux suppressants’, which forma protective layeron top ofthe stomach contentsto preventstomach acid escaping fromthe stomachwhere itworksinto the food pipe causingpain and discomfort. SodiumAlginate, SodiumBicarbonate, CalciumCarbonate(Gaviscon)ChewableTabletsis used forthe treatment ofsymptoms of gastro-oesophagealrefluxsuch as acid regurgitation, heartburn and indigestion (relatedto reflux), whichmay occur,forexample,followingmeals orduringpregnancy,and in patientswith symptomsrelatedto oesophagitis.
This medicine starts working after 3 to 4minutes of use.
This medicine is indicated for adults and children aged 12 years and above.
2. BEFORE USING SODIUM ALGINATE, SODIUM BICARBONATE, CALCIUM CARBONATE (GAVISCON) CHEWABLE TABLETS
Do not take this product if:
You know you are allergic to any of the ingredients as very rarely difficultyin breathing and skin rashes have occurred (see further information for full ingredients list).
Take special care before treatment with this product:
Talk to your doctor or pharmacist before taking this medicine if you:
• are on a low salt diet due to kidney problems or heart failure
• have high calcium levels in your blood
• have calcium deposition in the kidney
• have kidney stones
Sodium Alginate
Calcium Carbonate
Sodium Bicarbonate

Sodium Alginate
Sodium Bicarbonate
Calcium Carbonate

GAVISCON
250 mg/ 133.50 mg/ 80 mg Chewable Tablet

1. WHAT IS SODIUM ALGINATE, SODIUM BICARBONATE, CALCIUM CARBONATE (GAVISCON) CHEWABLE TABLETS AND WHAT IS IT USED FOR?

This product belongs to a group of medicines called 'reflux suppressants', which form a protective layer on top of the stomach contents to prevent stomach acid escaping from the stomach where it works into the food pipe causing pain and discomfort. Sodium Alginate, Sodium Bicarbonate, Calcium Carbonate (Gaviscon) Chewable Tablets is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion (related to reflux), which may occur, for example, following meals or during pregnancy, and in patients with symptoms related to oesophagitis.
This medicine starts working after 3 to 4 minutes of use.
This medicine is indicated for adults and children aged 12 years and above.

2. BEFORE USING SODIUM ALGINATE, SODIUM BICARBONATE, CALCIUM CARBONATE (GAVISCON) CHEWABLE TABLETS

Do not take this product if:
You know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see further information for full ingredients list).

Take special care before treatment with this product:
Talk to your doctor or pharmacist before taking this medicine if you:

- are on a low salt diet due to kidney problems or heart failure
- have high calcium levels in your blood
- have calcium deposition in the kidney
- have kidney stones

This medicine contains 63.16 mg sodium (main component of cooking/table salt) in each tablet. This is equivalent to 3.2 % of the recommended maximum daily dietary intake of sodium for an adult. This medicine also contains 3.2mmol of calcium per 4 tablet dose. The maximum recommended daily dose of this medicinal product contains 1012 mg sodium (found in table salt). This is equivalent to 50.6% of the adult recommended maximum daily dietary intake for sodium. Please also talk to your doctor regarding these salt contents if you suffer or have suffered from significant kidney or heart disease, as certain salts could interfere with these diseases. Talk to your pharmacist or doctor if you need this product on a daily basis for a prolonged period of time, especially if you have been advised to have a low salt diet.

This medicine contains 3.75 mg aspartame in each tablet. Aspartame is a source of phenylalanine. It may be harmful if you have phenylketonuria (PKU), a rare genetic disorder in which phenylalanine builds up because the body cannot remove it properly.

Taking Other Medicines:
Do not take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines. This is especially important if you are taking antihistamines, antibiotics (tetracyclines and quinolones such as norfloxacin), iron preparation, antifungals such as ketoconazole, digoxin and beta blockers (for heart conditions), penicillamine (commonly for rheumatoid arthritis), glucocorticoid (for inflammatory and autoimmune disorders), neuroleptics (for mental illness), thyroid hormones, chloroquine (for malaria), estramustine (for prostate cancer) and bisphosphonates (for osteoporosis).

Pregnancy and breast-feeding:
You can take this product if you are pregnant or breast - feeding. As with all medicines, the treatment duration should be limited as much as possible.

3. HOW TO TAKE SODIUM ALGINATE, SODIUM BICARBONATE, CALCIUM CARBONATE (GAVISCON) CHEWABLE TABLETS

For oral administration. Chew thoroughly before swallowing.
Adults, including the elderly and children 12 years and over: When symptoms occur take two to four tablets after meals and at bedtime, up to four times a day.
Children under 12 years: Should only be taken on medical advice.
If you forget a dose, it is not necessary to double the dose next time, just carry on taking as before.
If you take too much of this product you may feel bloated and experience some abdominal discomfort. It is unlikely to cause you any harm, but please consult your doctor or pharmacist.

After taking Sodium Alginate, Sodium Bicarbonate, Calcium Carbonate (Gaviscon) Chewable Tablets:
If symptoms persist after 7 days consult your doctor.

4. POSSIBLE SIDE EFFECTS

Very rare (less than 1 in 10,000) chance of an allergic reaction to the ingredients. Symptoms of this may include skin rash, itching, difficulty breathing, dizziness, or swelling of the face, lips, tongue or throat. If you experience these or any other side-effects stop taking the product and consult your doctor immediately.

5. HOW TO STORE SODIUM ALGINATE, SODIUM BICARBONATE, CALCIUM CARBONATE (GAVISCON) CHEWABLE TABLETS

Keep all medicines out of the reach and sight of children.
Do not use after the expiry date shown on the pack. Do not store above 30°C.

6. FURTHER INFORMATION

Sodium Alginate, Sodium Bicarbonate, Calcium Carbonate (Gaviscon) Chewable Tablets are circular tablets with an off white to cream, slightly motted appearance.

Each chewable tablet contains 250 mg sodium alginate, 133.5 mg sodium bicarbonate and 80 mg calcium carbonate as the active ingredients. The other ingredients are mannitol, macrogol 20,000, copovidone, aspartame, peppermint flavouring, magnesium stearate and acesulfame potassium.
These tablets do not contain sugar or gluten.

For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph. Seek medical attention immediately at the first sign of any adverse reaction.

Registration No.: DR-XY35582

Date of Last Revision: December 2023

Manufactured by: Reckitt Benckiser Healthcare (UK) Limited
Dansom Lane, Hull HU8 7DS UK

Imported by: Reckitt Benckiser Healthcare (Phils) Inc.
3/F Mead Johnson Bldg. 2309 Don Chino Roces Ave. Ext.,
Makati City, Philippines, 1231

Gaviscon Double Action (Sodium Alginate + Sodium Bicarbonate + Calcium Carbonate)

250 mg/106.5 mg/187.5 mg Chewable Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY43658

Gaviscon Double Action: Official FDA Philippines registration record

Brand name	Gaviscon Double Action
Generic name	Sodium Alginate + Sodium Bicarbonate + Calcium Carbonate
Dosage strength and form	250 mg/106.5 mg/187.5 mg Chewable Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antacid/Antireflux Agent
Registration number	DR-XY43658
Manufacturer	RECKITT BENCKISER HEALTHCARE (UK) LIMITED

Gaviscon Double Action: OTC forms registered under the Gaviscon brand (capsule availability)

OTC capsule form registered under the Gaviscon brand: NO. 7 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Gaviscon Infant	225 mg/ 87.5 mg Powder for Oral Suspension	Over-the-Counter (OTC) Drug	DR-XY46801
Gaviscon	250 mg/ 133.5 mg/ 80 mg Chewable Tablet	Over-the-Counter (OTC) Drug	DR-XY35582
Gaviscon Double Action	250 mg/106.5 mg/187.5 mg Chewable Tablet	Over-the-Counter (OTC) Drug	DR-XY43658
Gaviscon Double Action	250 mg/106.5 mg/187.5 mg Chewable Tablet (Mixed Berries Flavor)	Over-the-Counter Drug (OTC)	DR-XY46902
Gaviscon	500 mg/ 267 mg/ 160 mg per 10 mL Oral Suspension	Over-the-Counter (OTC) Drug	DR-XY34746
Gaviscon	500 mg/ 267 mg/ 160 mg per 10 mL Oral suspension	Over-the-Counter (OTC) Drug	DR-XY34907
Gaviscon Double Action	500 mg/213 mg/325 mg per 10 mL Suspension	Over-the-Counter (OTC) Drug	DR-XY43337

Gaviscon Double Action: Official leaflet (Package Insert / Patient Information Leaflet)

An official leaflet for this registration is published on the FDA Verification Portal: PIL_01 (https://drive.google.com/file/d/1tEu7YpwhWL4v_7SZ02ViCqNKMVZsyGgO/view?usp=drivesdk). Its text (OCR) and the original leaflet pages follow.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Gaviscon"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Gaviscon Double Action: Official leaflet text (OCR transcription)

Machine transcription of the official FDA leaflet for Reg. No. DR-XY43658. OCR may contain errors; the original leaflet pages are attached after this section.

Gaviscon Double Action leaflet PIL_01

CUSTOMER INFO
Minimum Point Size =6.28pt
28
Barcode Type: Pharmacode Standard
Code Value:
Magnification:
BARCODE INFO
Overall Height:
BWR Applied:
0 Micrometre
100 %
Bar Height: 13 mm
RB Artwork and
Print Specification
Trident Reference No:
ZEN Ref:
Action:
Brand:
Category:
Segment Group:
Segment:
Pack Size:
Market/Country:
Date:
RB547058
TR2685297
B
Gaviscon
Adult
Double Action Tablets
Tablets
12x6x8 Tablets
Philippines
03/12/2022
RBH Contact: Safina Liang
Artwork Type: BW Commercial
3253940
Edgemark Position: N/A
2nd Component Code (if applicable): N/A
3rd Party Code:
Pharmacode No/NE:
Component Code (2D if applicable):
Supply Point:
Finished Goods Code:
Parent Technical Packaging Specification:
1101 (28)
N/A
RB Hull
3026991
D0064065
CAD Cam Ref:
Printer:
Substrate:
G-Lflt-D0064065-175x200mm
Medica Packaging Limited (Crewe, UK)
Paper White
Technical & Non Printing Items
Black

1
LTH
Colours
Dieline
Guides
Dieline 2 (if applicable)
Guides 2 (if applicable)
UNDER NO CIRCUMSTANCES SHOULD THIS ARTWORK BE ALTERED WITHOUT PRIOR PERMISSION FROM TRIDENT.
Please note that any low resolution paper Canon colour copies associated with this job should be referred to for content, layout and colour separation only.
STUDIO USE ONLY Angelo Elfa v2.0
Connaught House, Connaught Road,
Kingswood Business Park, Hull, HU7 3AP, England.
T:+44 (0) 01482 828100

1. WHAT ARE SODIUM ALGINATE, SODIUM BICARBONATE, CALCIUM CARBONATE (GAVISCON DOUBLE ACTION) CHEWABLE TABLETS AND WHAT ARE THEY USED FOR?

This product is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate. It works in two ways:
1. Neutralizing excess stomach acid to relieve the pain and discomfort.
2. Forming a protective barrier over the stomach contents to soothe the burning pain in your chest.
This medicine is used for the treatment of acid related symptoms of gastro-esophageal reflux such as acid regurgitation, heartburn and indigestion which may occur, for example, following meals or during pregnancy.

2. BEFORE YOU TAKE SODIUM ALGINATE, SODIUM BICARBONATE, CALCIUM CARBONATE (GAVISCON DOUBLE ACTION) CHEWABLE TABLETS:

Do not take this product:
If you know you are allergic (hypersensitive) to any of the ingredients of this product as very rarely difficulty in breathing and skin rashes have occurred.
Take special care with this product:
This medicinal product contains 221.5mg sodium per 4 tablet dose, equivalent to 11% of the WHO recommended maximum daily intake for sodium.

3.HOW TO TAKE SODIUM ALGINATE, SODIUM BICARBONATE, CALCIUM CARBONATE (GAVISCON DOUBLE ACTION) CHEWABLE TABLETS:

For oral administration, chew thoroughly before swallowing. Adults and children 12 years and over: Two to four tablets after meals and at bedtime, up to four times per day. Children under 12 years: Should be given only on medical advice. Elderly: No dose modifications necessary for this age group.
If you take the product more than you should:
If you take too much of this product, it is unlikely to cause you any harm. However, you may feel bloated. Consult your doctor if this doesn’t go away.
If you forget to take the product:
If you forget a dose, it is not necessary to double the dose next time, just carry on taking as before.

4.POSSIBLE ADVERSE/UNDESIRABLE EFFECTS:

Like all medicines, the product can cause adverse/undesirable effects, although not everybody gets them. If you experience these adverse/ undesirable effects, stop taking the product and consult your doctor immediately. There is a very rare (less than 1 in 1000) chance of an allergic reaction to the ingredients. Symptom of this may include skin rash, itching, difficulty in breathing, dizziness, or swelling of the face, lips, tongue or throat. If any of the adverse/ undesirable effects gets serious, or you notice any adverse effects not listed in this leaflet, please consult your doctor.

5.HOW TO STORE SODIUM ALGINATE, SODIUM BICARBONATE,

CALCIUM CARBONATE (GAVISCON DOUBLE ACTION) CHEWABLE TABLETS:
Do not use this product after the expiry date (EXP:month/year) shown.
Keep out of the reach and sight of children. Store below 30°C and store in the original package to protect from moisture.
The maximum daily dose of this product is equivalent to 44% of the WHO recommended daily intake for sodium. This product is considered high in sodium. This should be particularly taken into account for those on a low salt diet, e.g. in some cases of congestive cardiac failure and renal impairment.
Please also talk to your doctor regarding these salt contents if you suffer or have suffered from kidney or heart disease, as certain salts could interfere with these diseases. Please consult your doctor if you know you have reduced amounts of gastric acid in your stomach, as this product may be less effective. The product should not be used in patients with severely impaired renal function and in patients with hypophosphatemia; Prolonged use should be avoided.
As with other antacid products, taking the product can mask the symptoms of other more serious, underlying medical conditions.
If symptoms persist after 7 days, consult your doctor.
Taking other medicines:
Do not take this medicine within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines.
Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines obtained without prescription.
Pregnancy and breast-feeding:
You can take this product if you are pregnant or breast-feeding.
As with all medicines, the treatment duration should be limited much as possible.
Important information about some of the ingredients of Sodium Alginate, Sodium Bicarbonate, Calcium Carbonate (Gaviscon Double Action) Chewable Tablets:
If you have phenylketonuria, note that the product is sweetened with aspartame (E951), a source of phenylalanine.
These tablets contain carmoisine lake (E122) which may cause an allergic reaction.
Medicines should not be disposed of via wastewater of household waste. Ask your pharmacist how to dispose of medicines that are no longer required.
These measures will help to protect the environment.
6.FURTHER INFORMATION
What Sodium Alginate, Sodium Bicarbonate, Calcium Carbonate (Gaviscon Double Action) Chewable Tablets contain:
The active substances in each chewable tablet are 250mg sodium alginate, 106.5mg sodium bicarbonate and 187.5mg calcium carbonate.
The other ingredients are xylitol, carmellose sodium, magnesium stearate, macrogol 20,000, mannitol (E421), copovidone, acesulfame-K, aspartame(E951), mint flavor and carmolsine lake (E122). This product does not contain sugar or gluten.
For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph. Seek medical attention immediately at the first sign of any adverse reaction.
Registration No.: DR-XY43658
Date of Last Revision: June 2021
250mg/106.5mg/187.5mg chewable tablet
DOUBLE ACTION
Mfd. by: Reckitt Benckiser Healthcare (UK) Limited
Dansom Lane, Hull HU8 7DS UK
Imported by:
Reckitt Benckiser Healthcare (Phils.) Inc.
3/F Mead Johnson Bldg.,
2309 Don Chino Roces Ave. Ext.,
Makati City, Philippines, 1231
3253940
N/A
PG1

Sodium Alginate Sodium Bicarbonate Calcium Carbonate

GAVISCON DOUBLE ACTION

250mg/106.5mg/187.5mg chewable tablet

1. WHAT ARE SODIUM ALGINATE, SODIUM BICARBONATE, CALCIUM CARBONATE (GAVISCON DOUBLE ACTION) CHEWABLE TABLETS AND WHAT ARE THEY USED FOR?

This product is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate. It works in two ways:

1. Neutralizing excess stomach acid to relieve the pain and discomfort.
2. Forming a protective barrier over the stomach contents to soothe the burning pain in your chest.

This medicine is used for the treatment of acid related symptoms of gastro-esophageal reflux such as acid regurgitation, heartburn and indigestion which may occur, for example, following meals or during pregnancy.

2. BEFORE YOU TAKE SODIUM ALGINATE, SODIUM BICARBONATE, CALCIUM CARBONATE (GAVISCON DOUBLE ACTION) CHEWABLE TABLETS:

Do not take this product:

If you know you are allergic (hypersensitive) to any of the ingredients of this product as very rarely difficulty in breathing and skin rashes have occurred.

Take special care with this product:

This medicinal product contains 221.5mg sodium per 4 tablet dose, equivalent to 11% of the WHO recommended maximum daily intake for sodium.

The maximum daily dose of this product is equivalent to 44% of the WHO recommended daily intake for sodium. This product is considered high in sodium. This should be particularly taken into account for those on a low salt diet, e.g. in some cases of congestive cardiac failure and renal impairment.

Please also talk to your doctor regarding these salt contents if you suffer or have suffered from kidney or heart disease, as certain salts could interfere with these diseases. Please consult your doctor if you know you have reduced amounts of gastric acid in your stomach, as this product may be less effective. The product should not be used in patients with severely impaired renal function and in patients with hypophosphatemia; Prolonged use should be avoided.

As with other antacid products, taking the product can mask the symptoms of other more serious, underlying medical conditions.

If symptoms persist after 7 days, consult your doctor.

Taking other medicines:

Do not take this medicine within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines.

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines obtained without prescription.

Pregnancy and breast-feeding:

You can take this product if you are pregnant or breast-feeding.
As with all medicines, the treatment duration should be limited much as possible.

Important information about some of the ingredients of Sodium Alginate, Sodium Bicarbonate, Calcium Carbonate (Gaviscon Double Action) Chewable Tablets:

If you have phenylketonuria, note that the product is sweetened with aspartame (E951), a source of phenylalanine.
These tablets contain carmoisine lake (E122) which may cause an allergic reaction.

3253940

FIRST FOLD

25mm 25mm 25mm 25mm 200mm

25mm 25mm 25mm 25mm 200mm

200mm

3.HOW TO TAKE SODIUM ALGINATE, SODIUM BICARBONATE, CALCIUM CARBONATE (GAVISCON DOUBLE ACTION) CHEWABLE TABLETS:

For oral administration, chew thoroughly before swallowing. Adults and children 12 years and over: Two to four tablets after meals and at bedtime, up to four times per day. Children under 12 years: Should be given only on medical advice. Elderly: No dose modifications necessary for this age group.

If you take the product more than you should:
If you take too much of this product, it is unlikely to cause you any harm. However, you may feel bloated. Consult your doctor if this doesn't go away.

If you forget to take the product:

If you forget a dose, it is not necessary to double the dose next time, just carry on taking as before.

4.POSSIBLE ADVERSE/UNDESIRABLE EFFECTS:

Like all medicines, the product can cause adverse/undesirable effects, although not everybody gets them. If you experience these adverse/ undesirable effects, stop taking the product and consult your doctor immediately. There is a very rare (less than 1 in 1000) chance of an allergic reaction to the ingredients. Symptom of this may include skin rash, itching, difficulty in breathing, dizziness, or swelling of the face, lips, tongue or throat. If any of the adverse/ undesirable effects gets serious, or you notice any adverse effects not listed in this leaflet, please consult your doctor.

5.HOW TO STORE SODIUM ALGINATE, SODIUM BICARBONATE, CALCIUM CARBONATE (GAVISCON DOUBLE ACTION) CHEWABLE TABLETS:

Do not use this product after the expiry date (EXP:month/year) shown. Keep out of the reach and sight of children. Store below 30°C and store in the original package to protect from moisture.

Medicines should not be disposed of via wastewater of household waste. Ask your pharmacist how to dispose of medicines that are no longer required.
These measures will help to protect the environment.

6.FURTHER INFORMATION

What Sodium Alginate, Sodium Bicarbonate, Calcium Carbonate (Gaviscon Double Action) Chewable Tablets contain:

The active substances in each chewable tablet are 250mg sodium alginate, 106.5mg sodium bicarbonate and 187.5mg calcium carbonate. The other ingredients are xylitol, carmellose sodium, magnesium stearate, macrogol 20,000, mannitol (E421), copovidone, acesulfame-K, aspartame(E951), mint flavor and carmolsine lake (E122). This product does not contain sugar or gluten.

For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph. Seek medical attention immediately at the first sign of any adverse reaction.

Registration No.: DR-XY43658

Date of Last Revision: June 2021

HEALTH ▸ HYGIENE ▸ HOME

Mfd. by: Reckitt Benckiser Healthcare (UK) Limited
Dansom Lane, Hull HU8 7DS UK

Imported by:
Reckitt Benckiser Healthcare (Phils.) Inc.
3/F Mead Johnson Bldg.,
2309 Don Chino Roces Ave. Ext.,
Makati City, Philippines, 1231

3253940

Gaviscon Infant (Sodium Alginate + Magnesium Alginate)

225 mg/ 87.5 mg Powder for Oral Suspension | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY46801

Gaviscon Infant: Official FDA Philippines registration record

Brand name	Gaviscon Infant
Generic name	Sodium Alginate + Magnesium Alginate
Dosage strength and form	225 mg/ 87.5 mg Powder for Oral Suspension
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antacid/Anti-Reflux
Registration number	DR-XY46801
Manufacturer	Reckitt Benckiser Healthcare (UK) Limited

Gaviscon Infant: OTC forms registered under the Gaviscon brand (capsule availability)

OTC capsule form registered under the Gaviscon brand: NO. 7 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Gaviscon Infant	225 mg/ 87.5 mg Powder for Oral Suspension	Over-the-Counter (OTC) Drug	DR-XY46801
Gaviscon	250 mg/ 133.5 mg/ 80 mg Chewable Tablet	Over-the-Counter (OTC) Drug	DR-XY35582
Gaviscon Double Action	250 mg/106.5 mg/187.5 mg Chewable Tablet	Over-the-Counter (OTC) Drug	DR-XY43658
Gaviscon Double Action	250 mg/106.5 mg/187.5 mg Chewable Tablet (Mixed Berries Flavor)	Over-the-Counter Drug (OTC)	DR-XY46902
Gaviscon	500 mg/ 267 mg/ 160 mg per 10 mL Oral Suspension	Over-the-Counter (OTC) Drug	DR-XY34746
Gaviscon	500 mg/ 267 mg/ 160 mg per 10 mL Oral suspension	Over-the-Counter (OTC) Drug	DR-XY34907
Gaviscon Double Action	500 mg/213 mg/325 mg per 10 mL Suspension	Over-the-Counter (OTC) Drug	DR-XY43337

Gaviscon Infant: Official leaflet (Package Insert / Patient Information Leaflet)

An official leaflet for this registration is published on the FDA Verification Portal: PIL_01 (https://drive.google.com/file/d/1ns1_Wcnljzutg0Bxa25hdolvxEhAsdQT/view?usp=drivesdk). Its text (OCR) and the original leaflet pages follow.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Gaviscon"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Gaviscon Infant: Official leaflet text (OCR transcription)

Machine transcription of the official FDA leaflet for Reg. No. DR-XY46801. OCR may contain errors; the original leaflet pages are attached after this section.

Gaviscon Infant leaflet PIL_01

CUSTOMER INFO:
Minimum Point Size = 7.00pt
N/A
BARCODE INFO
Barcode Type:
Barcode Number:
Magnification:
Truncated By:
Full Height:
Bar Height (Smallest Bar):
BWR:
Encoded Data:
Trident Reference No:
ZEN Ref:
Action:
Brand:
Category:
Segment Group:
Segment:
Pack Size:
Market/Country:
Date:
Component Code (2D if applicable)
:
2nd Component Code (if applicable)
Parent Technical Packaging Specification:
Finished Goods Code:
Supply Point:
3rd Party Code
Pharmacode No/NE:
Edgemark Position:
CAD Cam Ref:
Printer:
Substrate:
01000(39)
N/A
RB Hull
3164116
Gaviscon
Infant
Powder
Sachet
16x12x15x2Pack
RB443728
Please note that any low resolution paper Canon colour copies associated with this job should be referred to for content, layout and colour separation only.
STUDIO USE ONLY v2.0
Sm•Art check results:
Cutter
Guides
Technical & Non Printing Items
Cutter 2 (if applicable)
Guides 2 (if applicable)
Multi Packaging Solutions (Tewkesbury, UK)
G-Lflt-D0303972-256x130mm
TR2022205
Paper White
Philippines
27/11/20 F

Jamie Tan
3173800
N/A
G=2; O=4; R=0; - JT - 27/11/20 09:04:19
RB Artwork and
Print Specification
D0303972
Artwork Type: BW Commercial
10 mm 20 30 40 50 60 70 80 90 100 110 120 130 140 150 160 170 180 190 200 210 220 230 240 250
1
LTH
Process Black
Colours (Leaflet)
RBH Contact: Melvin Cruz
Connaught House, Connaught Road,
Kingswood Business Park, Hull, HU7 3AP, England.
T: +44 (0) 1482 828100
UNDER NO CIRCUMSTANCES SHOULD THIS
ARTWORK BE ALTERED WITHOUT PRIOR
PERMISSION FROM TRIDENT.
• Keep this leaflet. You may need to read it again.
• If you have any further questions, ask your doctor or pharmacist.
• This medicine has been prescribed for your child only. Do not pass it on to
others. It may harm them, even if their signs of illness are the same as your child's.
• If your child has any side effects, talk to your doctor or pharmacist. This
includes any possible side effects not listed in this leaflet. See section 5.
• Except on a doctor's or other healthcare professional's recommendation
• If your child is allergic to any of the ingredients of Sodium Alginate/Magnesium Alginate
(Gaviscon Infant) (listed in section 1)
• If your child is being sick, has a fever or diarrhoea
• If you have been told or it is suspected that your child has damage to their kidneys or
they have been put on a low salt diet
• If you have been told that your child has a blockage in their gut (an intestinal obstruction)
The active substances in each Sodium Alginate/Magnesium Alginate (Gaviscon
Infant) sachet are 225 mg sodium alginate and 87.5 mg magnesium alginate.
The other ingredients are mannitol (E421) and colloidal silica.
Sodium Alginate/Magnesium Alginate (Gaviscon Infant) is a powder for oral suspension. 2. WHAT SODIUM
ALGINATE/MAGNESIUM ALGINATE (GAVISCON
INFANT) IS AND WHAT IT IS USED FOR?
Sodium Alginate/Magnesium Alginate (Gaviscon Infant) contains the active
substances sodium alginate and magnesium alginate which belong to a group of
medicines called "reflux suppressants".
Reflux is the process by which the stomach contents flow back up the tube that
leads from the mouth to the stomach (known as the gullet or oesophagus). This can
cause an infant to be sick, spit up more than normal and become distressed. Sodium
Alginate/Magnesium Alginate (Gaviscon Infant) prevents reflux by stabilising the
stomach contents, so reducing the incidence of reflux and bringing up of feeds
(posseting) in infants and young children aged 1 to 24 months.
Always use Sodium Alginate/Magnesium Alginate (Gaviscon Infant) exactly as
described in this leaflet or as your doctor or pharmacist has told you. Check with
your doctor or pharmacist if you are not sure.
For infants aged 1 to 24 months. Sodium Alginate/Magnesium Alginate (Gaviscon
Infant) should not be used in premature infants or those under 1 month of age
unless advised by your doctor. Not suitable for children over 24 months (2 years),
adults or the elderly.
The dosage and mixing instructions depend on the weight of your child and the
feeding method you use:
Infants under 4.5 kg (10 lb) – one sachet should be used
Infants over 4.5 kg (10 lb) – two sachets should be used
Open the sachet(s) and mix the contents immediately before use as directed below:
PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER
Sodium Alginate
Magnesium Alginate
225mg/87.5mg Powder for Oral Suspension
Antacid/ Anti-Reux
1
. NAME, DESCRIPTION AND STRENGTH OF THE MEDICINE.

WHAT IS IN THE MEDICINE? 3. HOW MUCH AND HOW OFTEN SHOULD YOU USE SODIUM ALGINATE/MAGNESIUM ALGINATE (GAVISCON INFANT)?
WHEN SHOULD YOU NOT TAKE SODIUM ALGINATE/MAGNESIUM ALGINATE (GAVISCON INFANT)? CARE THAT SHOULD BE TAKEN WHEN TAKING SODIUM ALGINATE/MAGNESIUM ALGINATE (GAVISCON INFANT)?
Bottle fed infants
Breast fed infants
and other infants
(2 years) of age
• Mix each sachet into 115 ml (4 fluid oz) of milk feed in the bottle
• Shake well
• Feed as normal
• Mix each sachet with 5 ml (1 teaspoon) of cooled, boiled water until a smooth paste is formed
• Add another 10 ml (2 teaspoons) of cooled, boiled water and mix
• For breast fed infants give Sodium Alginate/Magnesium Alginate (Gaviscon Infant) part way through each feed or meal using a spoon or feeding bottle
• For all other infants give Sodium Alginate/Magnesium Alginate (Gaviscon Infant) at the end of each meal using a spoon or feeding bottle
Do not give more than six times (maximum of 6 or 12 sachets depending on the weight of your child) in 24 hours.
If your child's condition persists or becomes worse ask your doctor or other healthcare professional for advice immediately.
You should not use Sodium Alginate/Magnesium Alginate (Gaviscon Infant) for more than seven days without checking with your doctor or other healthcare professional. 4. Do not use Sodium Alginate/Magnesium Alginate (Gaviscon Infant):
5
. UNDESIRABLE EFFECTS OF SODIUM ALGINATE/MAGNESIUM ALGINATE (GAVISCON INFANT)?
WHAT OTHER MEDICINE OR FOOD SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?
Like all medicines, Sodium Alginate/Magnesium Alginate (Gaviscon Infant) can cause side effects, although not everybody gets them.
Very rarely (less than 1 in 10,000 patients) patients may experience constipation and diarrhoea. Side effects for which the frequency cannot be estimated from available data include a rash, bloating, discomfort of the stomach (gastric distention) and intestinal obstruction.
If the child has any side effects, stop giving Sodium Alginate/Magnesium Alginate (Gaviscon Infant) and tell your doctor or other healthcare professional.
Reporting of side effects
If you get any side effects, talk to your doctor or pharmacist.This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Gaviscon Infoline 1800-10-8896292/ MMla: 8889-6292 or
Email at: FeedbackPH@reckittbenckiser.com. By reporting side effects you can help provide more information on the safety of this medicine.
Other medicines and Sodium Alginate/Magnesium Alginate (Gaviscon Infant)
Tell your doctor or pharmacist if your child is taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.
Sodium Alginate/Magnesium Alginate (Gaviscon Infant) with food and drink
Do not use Sodium Alginate/Magnesium Alginate (Gaviscon Infant) with other feed thickening agents or infant milk preparations containing a thickening agent.
You should ask your doctor or pharmacist before you give Sodium Alginate/ Magnesium Alginate (Gaviscon Infant) to your child if you are not sure.
Sodium Alginate/Magnesium Alginate (Gaviscon Infant) contains sodium
This medicine contains 23.9 mg sodium (main component of cooking/table salt) in each sachet. This is equivalent to 1% of the recommended maximum daily dietary intake of sodium for an adult.
If you forget to give Sodium Alginate/Magnesium Alginate (Gaviscon Infant):
If a dose is missed do not double dose with the next feed. Just carry on with the recommended dose.
If your child takes more Sodium Alginate/Magnesium Alginate (Gaviscon Infant) than they should: If your child takes too much Sodium Alginate/Magnesium Alginate (Gaviscon Infant) they may suffer from intestinal obstructions, be sick, lose their appetite, become irritable or tired and weak.
If you accidently give too much Sodium Alginate/Magnesium Alginate (Gaviscon

Infant) ask your doctor or other healthcare professional for advice immediately. 6. 7. WHAT SHOULD YOU DO IF YOU MISS A DOSE? 8. SIGNS AND SYMPTOMS OF OVERDOSE
WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN THE
RECOMMENDED DOSAGE? 9. DOSAGE?
10
. HOW SHOULD YOU KEEP THIS MEDICINE?
Keep this medicine out of the sight and reach of children.
Do not use this medicine after the expiry date which is stated on the sachet and carton after EXP(month/year).
Store at temperatures not exceeding 30°C.
Do not use if the sachet is damaged or the security seal is broken.
Medicines should not be disposed of via wastewater or household waste.
Ask your pharmacist how to dispose of medicines no longer required. These measures will help protect the environment.
If your child's symptoms persist after 7 days consult your doctor.
Manufacturer and Marketing Authorization Holder:
Manufactured by: Reckitt Benckiser Healthcare (UK)
Limited Dansom Lane, Hull HU8 7DS, United Kingdom
Imported by: Reckitt Benckiser Healthcare (Phils.) Inc.
3/F Mead Johnson Bldg., 2309 Don Chino Roces Ave. Ext.,
Makati City, Philippines, 1231
Questions or concerns:
Gaviscon Infoline 1800-10-8896292/ MMla: 8889-6292
Email at: FeedbackPH@reckittbenckiser.com
For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph.
Seek medical attention immediately at the first sign of any adverse drug reaction.
Registration Number: DR-XY46801
Date of First Authorization: 21 January 2020
Date of Revision of Patient Information Leaet: October 2020
11
. WHEN SHOULD YOU CONSULT YOUR DOCTOR?
Read all of this leaflet carefully before you start using this medicine because it contains important information for you and your child.
Warnings and Precautions
Talk to your doctor or pharmacist before using Sodium Alginate/Magnesium Alginate (Gaviscon Infant) if your baby is premature or under one month.
There have been extremely rare reports of altered stool consistency (e.g. constipation and diarrhoea) in relation to use of this product. If you are concerned or you notice significant or sustained changes in bowel habit, consult a healthcare professional immediately.
N/A
3173800
3173800 3173800
Pharmacode Reads This Way
Last Bar First Bar

PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER

Sodium Alginate Magnesium Alginate

GAVISCON Infant

225mg/87.5mg Powder for Oral Suspension
Antacid/ Anti-Reflux

Read all of this leaflet carefully before you start using this medicine because it contains important information for you and your child.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for your child only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as your child's.
- If your child has any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 5.

1. NAME, DESCRIPTION AND STRENGTH OF THE MEDICINE. WHAT IS IN THE MEDICINE?

The active substances in each Sodium Alginate/Magnesium Alginate (Gaviscon Infant) sachet are 225 mg sodium alginate and 87.5 mg magnesium alginate. The other ingredients are mannitol (E421) and colloidal silica.

Sodium Alginate/Magnesium Alginate (Gaviscon Infant) is a powder for oral suspension.

2. WHAT SODIUM ALGINATE/MAGNESIUM ALGINATE (GAVISCON INFANT) IS AND WHAT IT IS USED FOR?

Sodium Alginate/Magnesium Alginate (Gaviscon Infant) contains the active substances sodium alginate and magnesium alginate which belong to a group of medicines called "reflux suppressants".

Reflux is the process by which the stomach contents flow back up the tube that leads from the mouth to the stomach (known as the gullet or oesophagus). This can cause an infant to be sick, spit up more than normal and become distressed. Sodium Alginate/Magnesium Alginate (Gaviscon Infant) prevents reflux by stabilising the stomach contents, so reducing the incidence of reflux and bringing up of feeds (posseting) in infants and young children aged 1 to 24 months.

3. HOW MUCH AND HOW OFTEN SHOULD YOU USE SODIUM ALGINATE/MAGNESIUM ALGINATE (GAVISCON INFANT)?

Always use Sodium Alginate/Magnesium Alginate (Gaviscon Infant) exactly as described in this leaflet or as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

For infants aged 1 to 24 months. Sodium Alginate/Magnesium Alginate (Gaviscon Infant) should **not** be used in premature infants or those under 1 month of age unless advised by your doctor. Not suitable for children over 24 months (2 years), adults or the elderly.

The dosage and mixing instructions depend on the weight of your child and the feeding method you use:
Infants under 4.5 kg (10 lb) – one sachet should be used
Infants over 4.5 kg (10 lb) – two sachets should be used

Open the sachet(s) and mix the contents immediately before use as directed below:

Bottle fed infants	• Mix each sachet into 115 ml (4 fluid oz) of milk feed in the bottle • Shake well • Feed as normal
Breast fed infants and other infants up to 24 months (2 years) of age	• Mix each sachet with 5 ml (1 teaspoon) of cooled, boiled water until a smooth paste is formed • Add another 10 ml (2 teaspoons) of cooled, boiled water and mix • For breast fed infants give Sodium Alginate/Magnesium Alginate (Gaviscon Infant) part way through each feed or meal using a spoon or feeding bottle • For all other infants give Sodium Alginate/Magnesium Alginate (Gaviscon Infant) at the end of each meal using a spoon or feeding bottle

Do not give more than six times (maximum of 6 or 12 sachets depending on the weight of your child) in 24 hours.

If your child's condition persists or becomes worse ask your doctor or other healthcare professional for advice immediately.

You **should not** use Sodium Alginate/Magnesium Alginate (Gaviscon Infant) for more than seven days without checking with your doctor or other healthcare professional.

4. WHEN SHOULD YOU NOT TAKE SODIUM ALGINATE/MAGNESIUM ALGINATE (GAVISCON INFANT)? CARE THAT SHOULD BE TAKEN WHEN TAKING SODIUM ALGINATE/MAGNESIUM ALGINATE (GAVISCON INFANT)?

Do not use Sodium Alginate/Magnesium Alginate (Gaviscon Infant):

- Except on a doctor's or other healthcare professional's recommendation
- If your child is allergic to any of the ingredients of Sodium Alginate/Magnesium Alginate (Gaviscon Infant) (listed in section 1)
- If your child is being sick, has a fever or diarrhoea
- If you have been told or it is suspected that your child has damage to their kidneys or they have been put on a low salt diet
- If you have been told that your child has a blockage in their gut (an intestinal obstruction)

Warnings and Precautions
Talk to your doctor or pharmacist before using Sodium Alginate/Magnesium Alginate (Gaviscon Infant) if your baby is premature or under one month.

There have been extremely rare reports of altered stool consistency (e.g. constipation and diarrhoea) in relation to use of this product. If you are concerned or you notice significant or sustained changes in bowel habit, consult a healthcare professional immediately.

5. UNDESIRABLE EFFECTS OF SODIUM ALGINATE/MAGNESIUM ALGINATE (GAVISCON INFANT)?

Like all medicines, Sodium Alginate/Magnesium Alginate (Gaviscon Infant) can cause side effects, although not everybody gets them.

Very rarely (less than 1 in 10,000 patients) patients may experience constipation and diarrhoea. Side effects for which the frequency cannot be estimated from available data include a rash, bloating, discomfort of the stomach (gastric distention) and intestinal obstruction.

If the child has any side effects, stop giving Sodium Alginate/Magnesium Alginate (Gaviscon Infant) and tell your doctor or other healthcare professional.

Reporting of side effects
If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Gaviscon Infoline 1800-10-8896292/ MMIa: 8889-6292 or
Email at: FeedbackPH@reckittbenckiser.com. By reporting side effects you can help provide more information on the safety of this medicine.

6. WHAT OTHER MEDICINE OR FOOD SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?

Other medicines and Sodium Alginate/Magnesium Alginate (Gaviscon Infant)
Tell your doctor or pharmacist if your child is taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

Sodium Alginate/Magnesium Alginate (Gaviscon Infant) with food and drink
Do not use Sodium Alginate/Magnesium Alginate (Gaviscon Infant) with other feed thickening agents or infant milk preparations containing a thickening agent. You should ask your doctor or pharmacist before you give Sodium Alginate/ Magnesium Alginate (Gaviscon Infant) to your child if you are not sure.

Sodium Alginate/Magnesium Alginate (Gaviscon Infant) contains sodium
This medicine contains 23.9 mg sodium (main component of cooking/table salt) in each sachet. This is equivalent to 1% of the recommended maximum daily dietary intake of sodium for an adult.

7. WHAT SHOULD YOU DO IF YOU MISS A DOSE?

If you forget to give Sodium Alginate/Magnesium Alginate (Gaviscon Infant):
If a dose is missed do not double dose with the next feed. Just carry on with the recommended dose.

8. SIGNS AND SYMPTOMS OF OVERDOSE

If your child takes more Sodium Alginate/Magnesium Alginate (Gaviscon Infant) than they should: If your child takes too much Sodium Alginate/Magnesium Alginate (Gaviscon Infant) they may suffer from intestinal obstructions, be sick, lose their appetite, become irritable or tired and weak.

9. WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN THE RECOMMENDED DOSAGE?

If you accidently give too much Sodium Alginate/Magnesium Alginate (Gaviscon Infant) ask your doctor or other healthcare professional for advice **immediately.**

10. HOW SHOULD YOU KEEP THIS MEDICINE?

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the sachet and carton after EXP(month/year).

Store at temperatures not exceeding 30°C.

Do not use if the sachet is damaged or the security seal is broken.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help protect the environment.

11. WHEN SHOULD YOU CONSULT YOUR DOCTOR?

If your child's symptoms persist after 7 days consult your doctor.

HEALTH ▸ HYGIENE ▸ HOME

Manufacturer and Marketing Authorization Holder:
Manufactured by: Reckitt Benckiser Healthcare (UK) Limited Dansom Lane, Hull HU8 7DS, United Kingdom
Imported by: Reckitt Benckiser Healthcare (Phils.) Inc. 3/F Mead Johnson Bldg., 2309 Don Chino Roces Ave. Ext., Makati City, Philippines, 1231

Questions or concerns:
Gaviscon Infoline 1800-10-8896292/ MMIa: 8889-6292
Email at: FeedbackPH@reckittbenckiser.com
For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph.
Seek medical attention immediately at the first sign of any adverse drug reaction.

Registration Number: DR-XY46801
Date of First Authorization: 21 January 2020
Date of Revision of Patient Information Leaflet: October 2020

3173800

Gaviscon (Sodium Alginate + Sodium Bicarbonate + Calcium Carbonate)

500 mg/ 267 mg/ 160 mg per 10 mL Oral Suspension | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY34746

Gaviscon: Official FDA Philippines registration record

Brand name	Gaviscon
Generic name	Sodium Alginate + Sodium Bicarbonate + Calcium Carbonate
Dosage strength and form	500 mg/ 267 mg/ 160 mg per 10 mL Oral Suspension
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antacid
Registration number	DR-XY34746
Manufacturer	Reckitt Benckiser Healthcare (UK) Limited

Gaviscon: OTC forms registered under the Gaviscon brand (capsule availability)

OTC capsule form registered under the Gaviscon brand: NO. 7 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Gaviscon Infant	225 mg/ 87.5 mg Powder for Oral Suspension	Over-the-Counter (OTC) Drug	DR-XY46801
Gaviscon	250 mg/ 133.5 mg/ 80 mg Chewable Tablet	Over-the-Counter (OTC) Drug	DR-XY35582
Gaviscon Double Action	250 mg/106.5 mg/187.5 mg Chewable Tablet	Over-the-Counter (OTC) Drug	DR-XY43658
Gaviscon Double Action	250 mg/106.5 mg/187.5 mg Chewable Tablet (Mixed Berries Flavor)	Over-the-Counter Drug (OTC)	DR-XY46902
Gaviscon	500 mg/ 267 mg/ 160 mg per 10 mL Oral Suspension	Over-the-Counter (OTC) Drug	DR-XY34746
Gaviscon	500 mg/ 267 mg/ 160 mg per 10 mL Oral suspension	Over-the-Counter (OTC) Drug	DR-XY34907
Gaviscon Double Action	500 mg/213 mg/325 mg per 10 mL Suspension	Over-the-Counter (OTC) Drug	DR-XY43337

Gaviscon: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY34746 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Gaviscon"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Gaviscon Double Action (Sodium Alginate + Sodium Bicarbonate + Calcium Carbonate)

500 mg/213 mg/325 mg per 10 mL Suspension | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY43337

Gaviscon Double Action: Official FDA Philippines registration record

Brand name	Gaviscon Double Action
Generic name	Sodium Alginate + Sodium Bicarbonate + Calcium Carbonate
Dosage strength and form	500 mg/213 mg/325 mg per 10 mL Suspension
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antacid
Registration number	DR-XY43337
Manufacturer	RECKITT BENCKISER HEALTHCARE (UK) LIMITED

Gaviscon Double Action: OTC forms registered under the Gaviscon brand (capsule availability)

OTC capsule form registered under the Gaviscon brand: NO. 7 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Gaviscon Infant	225 mg/ 87.5 mg Powder for Oral Suspension	Over-the-Counter (OTC) Drug	DR-XY46801
Gaviscon	250 mg/ 133.5 mg/ 80 mg Chewable Tablet	Over-the-Counter (OTC) Drug	DR-XY35582
Gaviscon Double Action	250 mg/106.5 mg/187.5 mg Chewable Tablet	Over-the-Counter (OTC) Drug	DR-XY43658
Gaviscon Double Action	250 mg/106.5 mg/187.5 mg Chewable Tablet (Mixed Berries Flavor)	Over-the-Counter Drug (OTC)	DR-XY46902
Gaviscon	500 mg/ 267 mg/ 160 mg per 10 mL Oral Suspension	Over-the-Counter (OTC) Drug	DR-XY34746
Gaviscon	500 mg/ 267 mg/ 160 mg per 10 mL Oral suspension	Over-the-Counter (OTC) Drug	DR-XY34907
Gaviscon Double Action	500 mg/213 mg/325 mg per 10 mL Suspension	Over-the-Counter (OTC) Drug	DR-XY43337

Gaviscon Double Action: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY43337 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Gaviscon"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Hydrite (Oral Rehydration Salts)

Granules for Solution (Melon Flavor) | Household Remedy (HR) | FDA Reg. No. DRHR-1810

Hydrite: Official FDA Philippines registration record

Brand name	Hydrite
Generic name	Oral Rehydration Salts
Dosage strength and form	Granules for Solution (Melon Flavor)
Classification	Household Remedy (HR)
Pharmacologic category	Oral Rehydration Salts
Registration number	DRHR-1810
Manufacturer	Amherst Laboratories, Inc.

Hydrite: OTC forms registered under the Hydrite brand (capsule availability)

OTC capsule form registered under the Hydrite brand: NO. 4 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Hydrite	(see reverse for formulation) Granules for Solution (Apple Flavor)	Household Remedy (HR)	HRP-443
Hydrite	(see reverse for formulation) Granules for Solution (Banana Flavor)	Household Remedy (HR)	HRP-442
Hydrite	(see reverse for formulation) Granules for Solution (Unflavored)	Household Remedy (HR)	HRP-441
Hydrite	Granules for Solution (Melon Flavor)	Household Remedy (HR)	DRHR-1810

Hydrite: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-1810 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Hydrite"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Imodium (Loperamide hydrochloride)

2 mg Capsule | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY14916

Imodium: Official FDA Philippines registration record

Brand name	Imodium
Generic name	Loperamide hydrochloride
Dosage strength and form	2 mg Capsule
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antipropulsive
Registration number	DR-XY14916
Manufacturer	Interphil Laboratories, Inc.

Imodium: OTC forms registered under the Imodium brand (capsule availability)

OTC capsule form registered under the Imodium brand: YES. 2 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Imodium	2 mg Capsule	Over-the-Counter (OTC) Drug	DR-XY14916
Imodium Complete Relief	2 mg/125 mg Tablet	Over-the-Counter (OTC) Drug	DR-XY49096

Imodium: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY14916 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Imodium"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Imodium Complete Relief (Loperamide + Simethicone)

2 mg/125 mg Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY49096

Imodium Complete Relief: Official FDA Philippines registration record

Brand name	Imodium Complete Relief
Generic name	Loperamide + Simethicone
Dosage strength and form	2 mg/125 mg Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antipropulsive
Registration number	DR-XY49096
Manufacturer	JNTL Consumer Health (France) S.A.S.

Imodium Complete Relief: OTC forms registered under the Imodium brand (capsule availability)

OTC capsule form registered under the Imodium brand: YES. 2 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Imodium	2 mg Capsule	Over-the-Counter (OTC) Drug	DR-XY14916
Imodium Complete Relief	2 mg/125 mg Tablet	Over-the-Counter (OTC) Drug	DR-XY49096

Imodium Complete Relief: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY49096 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Imodium"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Kremil-S (Aluminum Hydroxide + Magnesium Hydroxide + Simethicone)

178 mg/233 mg/30 mg Chewable Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY29748

Kremil-S: Official FDA Philippines registration record

Brand name	Kremil-S
Generic name	Aluminum Hydroxide + Magnesium Hydroxide + Simethicone
Dosage strength and form	178 mg/233 mg/30 mg Chewable Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Antacid/Antiflatulent
Registration number	DR-XY29748
Manufacturer	UNILAB, Inc.

Kremil-S: OTC forms registered under the Kremil-S brand (capsule availability)

OTC capsule form registered under the Kremil-S brand: NO. 4 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Kremil-S Spasm Relief	10 mg Film-Coated Tablet	Over-the-Counter (OTC) Drug	DRP-16071-01
Kremil-S	178 mg/233 mg/30 mg Chewable Tablet	Over-the-Counter (OTC) Drug	DR-XY29748
Kremil-S Cool Relief	356 mg/466 mg/20 mg per 20 mL Suspension (Menthol Flavor)	Over-the-Counter (OTC) Drug	DR-XY47731
Kremil-S Cool Relief	500 mg / 213 mg / 325 mg per 10 mL Oral Suspension	Over-the-Counter (OTC) Drug	DRP-7065-01

Kremil-S: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY29748 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Kremil-S"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Kremil-S Spasm Relief (Hyoscine-N-Butylbromide)

10 mg Film-Coated Tablet | Over-the-Counter (OTC) Drug | FDA Reg. No. DRP-16071-01

Kremil-S Spasm Relief: Official FDA Philippines registration record

Brand name	Kremil-S Spasm Relief
Generic name	Hyoscine-N-Butylbromide
Dosage strength and form	10 mg Film-Coated Tablet
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Anticholinergic
Registration number	DRP-16071-01
Manufacturer	Amherst Laboratories, Inc.

Kremil-S Spasm Relief: OTC forms registered under the Kremil-S brand (capsule availability)

OTC capsule form registered under the Kremil-S brand: NO. 4 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Kremil-S Spasm Relief	10 mg Film-Coated Tablet	Over-the-Counter (OTC) Drug	DRP-16071-01
Kremil-S	178 mg/233 mg/30 mg Chewable Tablet	Over-the-Counter (OTC) Drug	DR-XY29748
Kremil-S Cool Relief	356 mg/466 mg/20 mg per 20 mL Suspension (Menthol Flavor)	Over-the-Counter (OTC) Drug	DR-XY47731
Kremil-S Cool Relief	500 mg / 213 mg / 325 mg per 10 mL Oral Suspension	Over-the-Counter (OTC) Drug	DRP-7065-01

Kremil-S Spasm Relief: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRP-16071-01 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Kremil-S"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Maalox No. 1 (Aluminum hydroxide + Magnesium hydroxide)

200 mg/200 mg Chewable Tablet | Household Remedy (HR) | FDA Reg. No. DRHR-697

Maalox No. 1: Official FDA Philippines registration record

Brand name	Maalox No. 1
Generic name	Aluminum hydroxide + Magnesium hydroxide
Dosage strength and form	200 mg/200 mg Chewable Tablet
Classification	Household Remedy (HR)
Pharmacologic category	Antacid
Registration number	DRHR-697
Manufacturer	Sanofi S.r.l.

Maalox No. 1: OTC forms registered under the Maalox brand (capsule availability)

OTC capsule form registered under the Maalox brand: NO. 3 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Maalox No. 1	200 mg/200 mg Chewable Tablet	Household Remedy (HR)	DRHR-697
Maalox Plus	225 mg/ 200 mg/ 25 mg per 5 mL Suspension	Over-the-Counter (OTC) Drug	DR-XY8389
Maalox	225mg/ 200mg per 5mL Suspension	Household Remedy (HR)	DRHR-698

Maalox No. 1: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-697 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Maalox"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Maalox (Aluminum hydroxide/ Magnesium hydroxide)

225mg/ 200mg per 5mL Suspension | Household Remedy (HR) | FDA Reg. No. DRHR-698

Maalox: Official FDA Philippines registration record

Brand name	Maalox
Generic name	Aluminum hydroxide/ Magnesium hydroxide
Dosage strength and form	225mg/ 200mg per 5mL Suspension
Classification	Household Remedy (HR)
Pharmacologic category	Antacid
Registration number	DRHR-698
Manufacturer	Sanofi S.p.A

Maalox: OTC forms registered under the Maalox brand (capsule availability)

OTC capsule form registered under the Maalox brand: NO. 3 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Maalox No. 1	200 mg/200 mg Chewable Tablet	Household Remedy (HR)	DRHR-697
Maalox Plus	225 mg/ 200 mg/ 25 mg per 5 mL Suspension	Over-the-Counter (OTC) Drug	DR-XY8389
Maalox	225mg/ 200mg per 5mL Suspension	Household Remedy (HR)	DRHR-698

Maalox: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-698 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Maalox"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Pedialyte 45 (Oral Rehydration Salts (Maintenance))

(see reverse for formulation) Oral Solution (Fruit Flavor) | Over-the-Counter (OTC) Drug | FDA Reg. No. DRHR-202

Pedialyte 45: Official FDA Philippines registration record

Brand name	Pedialyte 45
Generic name	Oral Rehydration Salts (Maintenance)
Dosage strength and form	(see reverse for formulation) Oral Solution (Fruit Flavor)
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Electrolyte Replenisher
Registration number	DRHR-202
Manufacturer	Euro-Med Laboratories Phil., Inc.

Pedialyte 45: OTC forms registered under the Pedialyte brand (capsule availability)

OTC capsule form registered under the Pedialyte brand: NO. 1 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Pedialyte 45	(see reverse for formulation) Oral Solution (Fruit Flavor)	Over-the-Counter (OTC) Drug	DRHR-202

Pedialyte 45: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DRHR-202 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Pedialyte"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.

Visine (Tetrahydrozoline Hydrochloride)

500 mcg/mL (0.05% w/v) Ophthalmic Solution (Eye Drops) | Over-the-Counter (OTC) Drug | FDA Reg. No. DR-XY37227

Visine: Official FDA Philippines registration record

Brand name	Visine
Generic name	Tetrahydrozoline Hydrochloride
Dosage strength and form	500 mcg/mL (0.05% w/v) Ophthalmic Solution (Eye Drops)
Classification	Over-the-Counter (OTC) Drug
Pharmacologic category	Vasoconstrictor
Registration number	DR-XY37227
Manufacturer	PT Pfizer Indonesia

Visine: OTC forms registered under the Visine brand (capsule availability)

OTC capsule form registered under the Visine brand: NO. 3 OTC / Household Remedy product(s) registered for this brand on the FDA Verification Portal:

Brand name	Dosage strength and form	Classification	Reg. No.
Visine	500 mcg/mL (0.05% w/v) Ophthalmic Solution (Eye Drops)	Over-the-Counter (OTC) Drug	DR-XY37227
Visine	500 mcg/mL (0.05% w/v) Ophthalmic Solution (Eye Drops)	Over-the-Counter (OTC) Drug	DR-XY46844
Visine	500mcg/mL (0.05% w/v) Ophthalmic Solution (Eye Drops)	Over-the-Counter (OTC) Drug	DR-XY47700

Visine: Official leaflet (Package Insert / Patient Information Leaflet)

No PI/PIL is published on the FDA Verification Portal for registration DR-XY37227 as of 2026-10-22. Indication and dosage are not covered by this record.

Source

FDA Philippines Verification Portal, https://verification.fda.gov.ph (search: "Visine"), retrieved 2026-10-22. Fields above are reproduced from the official registration record without modification.

Compiled for an academic RAG case study. Not medical advice. The portal states its information is for verification purposes. Verify current status on the portal before relying on it.