

# Matrix Population Models 1

Hal Caswell

Institute for Biodiversity and Ecosystem Dynamics  
University of Amsterdam

IDEM 128 May 2023



European Research Council  
Established by the European Commission



## Starting simple

Age classes 1,2,3

$$n_1(t+1) = F_1 n_1(t) + F_2 n_2(t) + F_3 n_3(t)$$

$$n_2(t+1) = P_1 n_1(t)$$

$$n_3(t+1) = P_2 n_2(t)$$

or

$$\begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix} (t+1) = \begin{pmatrix} F_1 & F_2 & F_3 \\ P_1 & 0 & 0 \\ 0 & P_2 & 0 \end{pmatrix} \begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix} (t)$$

or

$$\mathbf{n}(t+1) = \mathbf{A} \mathbf{n}(t)$$



"Matrices unify and simplify demography. ... For many of us, including myself, multidimensionality is esoteric ... But we can be sure that what is strange and difficult to us will be natural and easy to our children. What is a narrow and almost closed sect within the profession today is going to become standard and obvious demographic technique in the next decade. A table of working life made the way Dublin et al (1949) made theirs, or a table of marriage similarly calculated, will be regarded as quaint; it will compare with the matrix method discussed in this issue as a cumbersome and inflexible desk calculator of the 1940s would compare with a hand-programmable calculator of the late 1970s."

Nathan Keyfitz (1984)



## So, what is this doing?<sup>1</sup>

$$\begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix} (t+1) = \begin{pmatrix} F_1 & F_2 & F_3 \\ P_1 & 0 & 0 \\ 0 & P_2 & 0 \end{pmatrix} \begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix} (t)$$

<sup>1</sup>transformation, distribution, projection, age, time, ...



## P. H. Leslie (1900–1972)<sup>2</sup>

VOLUME XXXIII, PART III

NOVEMBER 1945

### ON THE USE OF MATRICES IN CERTAIN POPULATION MATHEMATICS

By P. H. LESLIE, *Bureau of Animal Population, Oxford University*

#### CONTENTS

	PAGE
1. Introduction . . . . .	183
2. Derivation of the matrix elements . . . . .	184
3. Numerical examples . . . . .	185
4. Properties of the basic matrix . . . . .	187
5. Transformation of the co-ordinates . . . . .	188
6. Relation between the canonical form . . . . .	188
B and the $L_m$ column . . . . .	190
7. The stable age distribution . . . . .	191
8. Properties of the stable vectors . . . . .	192
9. The spectral set of operators . . . . .	193
10. Relation of $\psi$ to classical canonical form . . . . .	194
11. The relation between $\phi$ and $\psi$ vectors . . . . .	195
12. Case of repeated latent roots . . . . .	197
13. The approach to the stable age distribution . . . . .	199
14. Stability of the matrix with only a single non-zero $F_{ij}$ element . . . . .	200
15. Numerical comparison with the usual methods of computation . . . . .	201
16. Further practical applications . . . . .	207
Appendix: (1) Calculation of mortality and fertility rates . . . . .	209
(2) Calculation of the rate of increase . . . . .	210
(3) Numerical values of the matrix elements . . . . .	212
References . . . . .	212

$$A = \begin{bmatrix} F_0 & F_1 & F_2 & F_3 & \dots & F_{k-1} & F_k \\ P_0 & \cdot & \cdot & \cdot & \dots & \cdot & \cdot \\ \cdot & P_1 & \cdot & \cdot & \dots & \cdot & \cdot \\ \cdot & \cdot & P_2 & \cdot & \dots & \cdot & \cdot \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \cdot & \cdot & \cdot & \cdot & \dots & P_{k-1} & \cdot \end{bmatrix}$$



<sup>2</sup>1945, Biometrika



## Leonard Lefkovitch (1929–2010)<sup>4</sup>

### THE STUDY OF POPULATION GROWTH IN ORGANISMS GROUPED BY STAGES

L. P. LEFKOVITCH

Agricultural Research Council, Pest Infestation Laboratory, Slough, Bucks, England

#### SUMMARY

In this extension to the use of matrices in population mathematics (Lewis [1942] and Leslie [1945]), the division of a population into equal age groups is replaced by one of unequal stage groups, no assumptions being made about the variation of the duration of the stage that different individuals may show. This extension has application in ecological studies where the age of an individual is rarely known. The model is briefly applied to three experimental situations.



$$M_{I'} = \begin{bmatrix} \cdot & 1.978 & \cdot & \cdot \\ 1.203 & 0.998 & 27.776 & 15.309 \\ 0.362 & 0.191 & \cdot & \cdot \\ \cdot & 0.835 & \cdot & \cdot \end{bmatrix}$$

<sup>4</sup>Biometrics 21:1–18, 1965



## Nathan Keyfitz (1913–2010)<sup>3</sup>

### THE POPULATION PROJECTION AS A MATRIX OPERATOR

N. KEYFITZ  
University of Chicago

#### RESUMEN

El método de "componentes" para proyectar la población, uno de los métodos demográficos más valiosos, ha sido tratado casi enteramente como método empírico, desprovisto de una teoría básica o una estructura matemática. Este artículo considera la proyección como una matriz, con énfasis en su naturaleza en la dimensión temporal y en la relación de una función que convierte la

$$M = \begin{bmatrix} 0 & 0 & .0570 & .2316 & .3338 & .2622 & .1565 & .0692 & .0158 \\ .99108 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & .99781 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & .99546 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & .99329 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & .99140 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & .98954 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & .98664 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & .98237 & 0 \end{bmatrix}$$



<sup>3</sup>Demography (1964). Volume 1. Issue 1!



## Why did it take 20 years?

## State variables for individuals and populations

individual state (i-state):



- information needed to determine the probabilities of survival, reproduction, and transition from  $t$  to  $t + 1$
- state + environment at time  $t$  determines state at time  $+1$
- demographers' favorite i-state variable is age

population state (p-state):


- distribution of individuals over i-state categories<sup>5</sup>
  - age → age distribution
  - size → size distribution
  - health status → health status distribution
  - you get the idea

<sup>5</sup>Subject to some conditions


## i-state variables and life cycle graphs




## Life cycle graphs



## Wandering albatross




## Snow petrel



mean = 0.98  
 SD = 3.8  
 CV = 3.9  
 skew = 5.8  
 Crow's  $\mathcal{I}$  = 15.2




# Greater snow goose



mean	=	2.1
SD	=	4.8
CV	=	2.3
skew	=	3.5
Crow's $\mathcal{I}$	=	5.3






A set of small, light-blue navigation icons typically found in presentation software like Beamer. They include symbols for back, forward, search, and other document-related functions.




**Figure 3** The illness-death or disability model.

A set of small, light-blue navigation icons typically found in Beamer presentations, including symbols for back, forward, search, and table of contents.









<sup>7</sup>Tyas et al. 2007



<sup>8</sup>Ragnel-Frausto et al. 1998




## Union status<sup>9</sup>



<sup>9</sup>Mills 2000

## From the life cycle graph to the matrix



$$\mathbf{A} = \begin{pmatrix} 0 & F_2 & F_3 & F_4 \\ S_1 & 0 & 0 & 0 \\ 0 & S_a & 0 & 0 \\ 0 & 0 & S_a & S_a \end{pmatrix}$$

Important

$a_{i,j}$  = coefficient from  $j$  to  $i$

## A useful decomposition

$$\mathbf{A} = \mathbf{U} + \mathbf{F}$$

$$\mathbf{A} = \underbrace{\begin{pmatrix} 0 & 0 & 0 & 0 \\ S_1 & 0 & 0 & 0 \\ 0 & S_a & 0 & 0 \\ 0 & 0 & S_a & S_a \end{pmatrix}}_{\mathbf{U}} + \underbrace{\begin{pmatrix} 0 & F_2 & F_3 & F_4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}}_{\mathbf{F}}$$

## Population projection

$$\mathbf{n}(t+1) = \mathbf{An}(t)$$

projection initial population  $\mathbf{n}(0)$

$$\mathbf{n}(1) = \mathbf{An}(0)$$

$$\begin{aligned} \mathbf{n}(2) &= \mathbf{An}(1) \\ &= \mathbf{A}^2 \mathbf{n}(0) \end{aligned}$$

$$\begin{aligned} \mathbf{n}(3) &= \mathbf{An}(2) \\ &= \mathbf{A}^3 \mathbf{n}(0) \end{aligned}$$

⋮

$$\mathbf{n}(t) = \mathbf{A}^t \mathbf{n}(0)$$

## Transient and asymptotic dynamics

$$\mathbf{A} = \begin{pmatrix} 0 & 1 & 5 \\ .3 & 0 & 0 \\ 0 & .5 & 0 \end{pmatrix} \quad \mathbf{n}(0) = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

projecting

$$\mathbf{n}(1) = \begin{pmatrix} 0 \\ 0.3 \\ 0 \end{pmatrix} \quad \mathbf{n}(2) = \begin{pmatrix} 0.3 \\ 0 \\ 0.15 \end{pmatrix} \quad \dots$$


## Some types of matrix models

- linear, time-invariant

$$\mathbf{n}(t+1) = \mathbf{An}(t)$$

- what else?

## Transient and asymptotic dynamics



## Next

- solve the projection equation
- explain transient dynamics
- characterize asymptotic dynamics