

# Report on comparison between analyses for semi- and dileptonic channel HH production at HL-LHC

Izaak Neutelings (University of Zurich)

November 29, 2015

In this reports, I tempt to reproduce selection level significances from the analysis note on the  $HH \rightarrow bbWW \rightarrow bb\ell\nu\ell\nu$  channel by C. Delaere *et al.* [1].

## 1 Samples

The Monte Carlo samples of the analysis note were generated by MADGRAPH\_AMC@NLO, and the parton shower and hadronization was done in PYTHIA6. Our samples were fully produced in PYTHIA6. All samples were finally reconstructed with Delphes for the CMS Phase II technical proposal. For our analysis, the samples only contain  $bbWW \rightarrow bb\ell\nu\ell\nu$  or  $bbWW \rightarrow bbqq\ell\nu$  at generator level, where taus coming from a W are excluded.

## 2 Event selections & clean-up

To reproduce the results by C. Delaere *et al.* the event selections and clean-up as described in their analysis note are applied to our samples. These cuts and the ones we use for the semileptonic final state are compared in Table 1.

For the semileptonic sample, the  $\Delta R_{\ell\ell}$  cut at generator level is replaced with  $\Delta R_{q\ell}$ , where  $q$  is the closest quark to the lepton. Please note that there are no equivalent cuts to  $\Delta R_{\ell\ell}$ ,  $m_{\ell\ell}$  or  $\Delta\phi_{bb,\ell\ell}$  used in the clean-up, leading to bigger differences between the two cases.

## 3 Cross section & branching ratios

For our analysis, we have been using the cross sections at  $\sqrt{s} = 14$  TeV given in Table 2. The analysis note lists next leading order  $\sigma_{LO}$  with k-factor  $k_{NNLO}$  of the samples, all listed in Table 3.

The branching ratios are found using

$$\mathcal{B}(HH \rightarrow bbWW) = 2\mathcal{B}(H \rightarrow bb)\mathcal{B}(H \rightarrow WW) \simeq 0.248$$

$$\mathcal{B}(t\bar{t} \rightarrow bWbW) = \mathcal{B}(t \rightarrow bW)^2 \simeq 0.997$$

$$\mathcal{B}(WW \rightarrow \ell\nu\ell\nu) = \mathcal{B}(W \rightarrow \ell\nu)^2 \simeq 0.046$$

$$\mathcal{B}(WW \rightarrow qq\ell\nu) = 2\mathcal{B}(W \rightarrow qq)\mathcal{B}(W \rightarrow \ell\nu) \simeq 0.288$$

with numbers from [5], [6] and [7]:

$$\mathcal{B}(HH \rightarrow bbWW \rightarrow bb\ell\nu\ell\nu) \simeq 0.011$$

$$\mathcal{B}(HH \rightarrow bbWW \rightarrow bbqq\ell\nu) \simeq 0.072$$

$$\mathcal{B}(t\bar{t} \rightarrow bbWW \rightarrow bbqq\ell\nu\ell\nu) \simeq 0.045$$

$$\mathcal{B}(t\bar{t} \rightarrow bbWW \rightarrow bbqq\ell\nu) \simeq 0.287$$

## 4 Significance & yield

Using the yield  $N = \sigma L$  with an integrated luminosity  $L = 3000 \text{ fb}^{-1}$ , the Punzi significance is calculated as:

$$P = \frac{N(HH)}{\sqrt{1 + N(tt)}}. \quad (1)$$

**Table 1:** Event selection and clean-up: comparison between the dileptonic and semileptonic final state.

dileptonic final state	semileptonic final state
<b>Generator level filter on background</b>	
b-quarks: $p_T > 15$ GeV	b-quarks: $p_T > 15$ GeV
leptons: $p_T > 15$ GeV, $ \eta  < 2.5$	lepton: $p_T > 15$ GeV, $ \eta  < 2.5$
$\Delta R_{\ell\ell} < 2.5$	$\Delta R_{q\ell} < 2.5$
<b>Selection</b>	
two b-jets: $p_T > 30$ GeV, $ \eta  < 2.5$	two b-jets: $p_T > 30$ GeV, $ \eta  < 2.5$
two oppositely charged leptons with:	two non b-jets: $p_T > 20$ GeV, $ \eta  < 2.5$
muons: $p_T > 20$ GeV, $ \eta  < 2.5$	one lepton with:
electrons: $p_T > 25$ GeV, $ \eta  < 2.5$	muon: $p_T > 20$ GeV, $ \eta  < 2.5$
MET $> 20$ GeV	electron: $p_T > 25$ GeV, $ \eta  < 2.5$
	MET $> 20$ GeV
<b>Clean-up</b>	
$60 \text{ GeV} < m_{bb} < 160 \text{ GeV}$	$60 \text{ GeV} < m_{bb} < 160 \text{ GeV}$
$\Delta R_{bb} < 3.1 \text{ GeV}$	$\Delta R_{bb} < 3.1 \text{ GeV}$
$m_{ll} < 85 \text{ GeV}$	
$\Delta R_{\ell\ell} < 2$	
$\Delta\phi_{bb,\ell\ell} < 1.7$	

**Table 2:** Cross sections at NNLO and  $\sqrt{s} = 14$  TeV [2][3], branching ratios (excluding  $W \rightarrow \tau\bar{\tau}$ ) [5][7] and number of Monte Carlo events per process in our analysis sample.

process	$\sigma\mathcal{B}$ [fb]	branching ratio $\mathcal{B}$	number of MC events
<b>HH</b>	<b>40</b>		
$HH \rightarrow bbWW \rightarrow bbqq\ell\nu$	2.86	0.072	166 483
$HH \rightarrow bbWW \rightarrow bb\ell\nu\ell\nu$	0.452	0.011	22 812
<b>t<math>\bar{t}</math></b>	<b>984 500</b>		
$t\bar{t} \rightarrow bbWW \rightarrow bbqq\ell\nu$	282 846	0.287	164 661
$t\bar{t} \rightarrow bbWW \rightarrow bb\ell\nu\ell\nu$	44 598	0.045	22 546

**Table 3:** Cross sections for their Monte Carlo samples listed in the analysis note by C. Deleare *et al.* [1].

process	$\sigma_{LO}$ [fb]	$k_{NNLO}$	number of MC events
$HH \rightarrow bbWW \rightarrow bb\ell\nu\ell\nu$	0.163	2.3	1.1M
t $\bar{t}$ full leptonic	9030	1.85	4.8M

**Table 4:** Number of MC events in our sample at each level of selection.

selection level	dileptonic final state		semileptonic final state	
	signal	background	signal	background
$bbWW \rightarrow bb\ell\nu\ell\nu$ or $bbWW \rightarrow bbqq\ell\nu$	22812	22546	166483	164661
Generator level filter	22812	8339	166483	68011
Selection	2748	1755	28203	19401
Clean-up	2149	296	22189	8748

**Table 5:** Comparison of the significance  $P$  and yields  $N$  between the semileptonic and dileptonic final state for our results using the cross sections at NNLO from Table 2 and the results by C. Delaere *et al.* at  $\sqrt{s} = 14$  TeV and with integrated luminosity  $L = 3000 \text{ fb}^{-1}$ . The factor  $k_B = \sqrt{\mathcal{B}(\text{WW} \rightarrow \ell\nu\ell\nu)/\mathcal{B}(\text{WW} \rightarrow \text{qq}\ell\nu)} \simeq 0.397$  allows for comparison.

	dileptonic final state			semileptonic final state			
	$P$	$N(\text{HH})$	$N(\text{t}\bar{\text{t}})$	$P$	$k_B P$	$N(\text{HH})$	$N(\text{t}\bar{\text{t}})$
<b>Initial significance, before any cuts</b>							
Assuming NNLO	0.117	1356	133 793 550	0.295	0.117	8580	848 540 550
<b>Gen level cuts on background</b>							
Assuming NNLO	0.193	1356	49 485 692	0.458	0.182	8580	350 478 202
<b>Selection</b>							
Assuming NNLO	0.051	163	10 414 605	0.145	0.058	1454	99 978 350
C. Delaere <i>et al.</i>	0.043	113	6 759 579				
<b>Clean-up</b>							
Assuming NNLO	0.096	127	1 756 537	0.170	0.068	1144	45 080 698
C. Delaere <i>et al.</i>	0.075	90	1 437 144				
<b>Neural network</b>							
C. Delaere <i>et al.</i>	0.60	37	3875				

To obtain the significance at after each round of cuts, we multiply each yield with the selection efficiency. The total number of Monte Carlo events per sample in our analysis is listed in Table 2.

## 5 Results

Results are summarized in Table 5. To compare the dileptonic to the semileptonic case, the significance of the latter is also scaled by  $k_B = \sqrt{\mathcal{B}(\text{WW} \rightarrow \ell\nu\ell\nu)/\mathcal{B}(\text{WW} \rightarrow \text{qq}\ell\nu)} \simeq 0.397$ .


## References

- [1] C. Delaere *et al.*, *Study of HH production with  $H \rightarrow b\bar{b}$ ,  $H \rightarrow WW \rightarrow \ell\nu\ell\nu$  for an upgraded CMS detector at the HL-LHC*, CMS draft analysis note 2014/141.
- [2] D. de Florian & J. Mazzitelli, *Higgs Boson Pair Production at Next-to-Next-to-Leading Order in QCD*. Phys. Rev. Lett. **111** (Nov, 2013) 201801, doi:10.1103/PhysRevLett.111.201801, arXiv:1309.6594.
- [3] *NNLO+NNLL top-quark-pair cross sections - ATLAS-CMS recommended predictions for top-quark-pair cross sections using the Top++v2.0 program (M. Czakon, A. Mitov, 2013)*, [https://twiki.cern.ch/twiki/bin/view/LHCPhysics/TtbarNNLO#Top\\_quark\\_pair\\_cross\\_sections\\_at](https://twiki.cern.ch/twiki/bin/view/LHCPhysics/TtbarNNLO#Top_quark_pair_cross_sections_at).
- [4] R. Frederix *et al.*, *Higgs pair production at the LHC with NLO and parton-shower effects*, Phys. Rev. Lett. **B723** (May, 2014) 142, doi:10.1016/j.physletb.2014.03.026, arXiv:1401.7340.
- [5] *Higgs cross sections for European Strategy studies in 2012*, [https://twiki.cern.ch/twiki/bin/view/LHCPhysics/HiggsEuropeanStrategy2012#SM\\_Higgs\\_decay\\_branching\\_ratio\\_M](https://twiki.cern.ch/twiki/bin/view/LHCPhysics/HiggsEuropeanStrategy2012#SM_Higgs_decay_branching_ratio_M).
- [6] T. Aaltonen *et al.* (CDF Collaboration), *Measurement of  $\mathcal{B}(t \rightarrow Wb)/\mathcal{B}(t \rightarrow Wq)$  in Top-Quark-Pair Decays Using Dilepton Events and the Full CDF Run II Data Set*, Phys. Rev. Lett. **112**, 221801 (June, 2014), doi:10.1103/PhysRevLett.112.221801, arXiv:1404.3392.
- [7] J. Beringer *et al.* (Particle Data Group), PR **D86**, 010001 (2012) and 2013 partial update for the 2014 edition (<http://pdg.lbl.gov/2013/listings/rpp2013-list-w-boson.pdf>).


## Appendix A: Preliminary results on multivariate analysis of $HH \rightarrow bbWW \rightarrow bbqq\ell\nu$

The multilayer perceptron (MLP) from the TMVA class in ROOT is used as neural network. The following variables are input for the NN:  $p_T$  of the two leading b-jets,  $p_T$  of the two leading light jets,  $p_T$  of the leading lepton, missing transverse momentum  $\cancel{E}_T$ ,  $\Delta R_{j_1\ell}$ ,  $\Delta R_{j_2\ell}$ ,  $\Delta R_{b_1\ell}$ ,  $\Delta R_{b_2\ell}$ ,  $\Delta R_{bb}$ ,  $M_{bb}$ ,  $M_{jj,b}$  and  $M_{b,\ell\nu}$ . Here  $j_1$  denotes the light jet closest to the lepton, and  $j_2$  the second closest, while  $b_1$  denotes the b-tagged jet farthest to the lepton and  $b_2$  the second farthest. In case of more than 2 b-jets, the b-jets pair closest in  $\Delta R_{bb}$  is used for  $M_{bb}$  and other b-tagged jets are then regarded as light jets. The other two invariant masses reconstruct a leptonic and hadronic top as follows: the two leading jets and closest b-jet second closest to the lepton form  $M_{jj,b}$  and the lepton, reconstructed neutrino and b-jet closest to the lepton make  $M_{b,\ell\nu}$ . The neutrino is reconstructed assuming its transverse momentum  $p_T^\nu$  is given by the missing transverse and its longitudinal component  $p_z^\nu$  is the real solution of  $M_W^2 = (p_\ell + p_\nu)^2$ .


All variables are shown Figs. 1-3.



**Figure 1:** Variables distribution of  $HH$  (red) and  $t\bar{t}$  (blue) for the neural network: transverse momentum  $p_T$  for the two leading jets and two leading b-jets.



**Figure 2:** Variables distribution of HH (red) and  $t\bar{t}$  (blue) for the neural network:  $p_T$  of the leading lepton,  $\cancel{E}_T$ ,  $M_{bb}$  and  $M_{jj,b}$ .



**Figure 3:** Variables distribution of HH (red) and  $t\bar{t}$  (blue) for the neural network:  $\Delta R$  between the lepton and the two closest light jets and between the lepton and two farthest b-jets.