EPSR-test-ex77 sharm DMSO-DMSO with high l (l1, l2 <= 6, l <= 12), mapgr, mapgrentropy

Program(s): sharm, mapgr, mapgrentropy | chained passes: 1 | verification: ALL IDENTICAL (1 pass)

What this case tests

A precision test of the special functions of `sharm` [1]: high orders lead to large log-factorial arguments in the Clebsch-Gordan coefficients (`cgcoef`) and Wigner rotation matrices (`dlm`), where single-precision rounding must match the Fortran exactly.

Code exercised

Fortran (original, fortran_src/):

- `sharm_rev2.f`: `run_sharm` (keywords, `nsumt` rule), `sharm` (centres, axes, pair loop, averaged molecules)
- `sharm_routines_rev1.f`: `get_l_and_n_values`, `form_cgcoeffs`, `form_eulerangles`, `form_dlmn`, `form_sum_prod_dlmn_cgcoeff`
- `auxiliary_routines.f`: `dlm`, `cgcoef`, `rotateeuler`; `read_and_save_files.f`: `outgofzs_sharm` (`.h0n` files, averaging); `process_symbols.f`: `check_symbol`; `lngamma_mod.f90`: `fctrln`
- `mapgrquad.f`: `run_mapgr`; `plotsharm_routines.f`: `smoosharm`, `sumoverm2`; `plot_routines.f`: `find_plot_par`
- `mapgrentropy.f`: `run_mapgrentropy` (entropy sums); `auxiliary_routines.f`: `initran1`, `saveran1`

Rust (port, rust/epsr-core/src/):

- `tools/sharm.rs`: `run`, `sharm`, `read_axis`, `read_l_values`, `average_molecule`
- `tools/sharm_routines.rs`: `Sharm` methods with the Fortran names (incl. `outgofzs_sharm`), `dlm`, `cgcoef`, `form_eulerangles`, `rotateeuler`; `LnGamma::fctrln`
- `tools/mapgr.rs`: `run`; `tools/sharm_routines.rs`: `Coeffs::read`, `Coeffs::angular`, `smoosharm`, `sumoverm2`, `check_symbol_mapgr`; `output.rs`: `w_int`
- `tools/mapgrentropy.rs`: `run`, `initran1`; `rng.rs`: `Ran1::ran1`

Physical system

The EPSR-test configuration `200DMSO400Water.ato`: 200 DMSO (atoms S, C, C, O, 6 H) and 400 water molecules (O1, H1, H1; SPC/E parameters [2]), 3200 atoms at 300 K in a cubic box of side 33.12 Å, a model of the DMSO-water mixtures of Soper and Luzar [3]. `sharm` expands the orientational pair correlation function of a central and a second molecule type in rotation matrices coupled by Clebsch-Gordan coefficients; the coefficients g(l1,l2,l;n1,n2;r) are averages over all molecule pairs [4,5]. Here identical DMSO molecules (centre = mass centre of S, C, C, O; axes -z towards O, y from C 2), l1, l2 = 0 2 4 6, l = 0-12 (even), n1step = n2step = 3: 181 coefficients in `.h01` and `.h02`. `mapgr` rebuilds g(r) along chosen directions and relative orientations from the coefficients (grtype 1 radial, 2 spatial, 3 full orientational). `mapgrentropy` adds to the ideal-gas entropy the two-body excess term, an integral of $r^2 \cdot (g - 1 - g \ln g)$ [6], for the radial, spatial and full g, sampling random orientations with `ran1` [7].

Inputs

Input files written by hand from the keyword tables (`setup_symbols_*`), one `keyword value` per line, ended by `q`; `200DMSO400Water.ato` copied from EPSR-test. `dmso.SHARM.dat` (nr 120, rmax 12); `dmso.mapgr.txt`: radial and two orientational views, smoothing `3.0 2 1 1`; `dmso.mapgrentropy.dat`: 40 trials, nfreq 10.

How the accuracy was verified

`commands.txt` (3 program calls, run once) was run with the Fortran reference build (`rust/tools/build_fortran_reference.sh`: the original sources, gfortran -O2 -fno-tree-vectorize, single thread) and with the Rust `epsr-rs` on identical copies of the inputs, and every file written was compared byte for byte (`rust/tools/run_ex_test.sh`, `compare_outputs.py`; no masking except where noted below). The Fortran output is stored in `fortran_reference.tar.gz`. A run-time FORMAT in the original source (`sharm_rev2.f`/`sdf_rev2.f`, and `sharm_diffuse` for `epsr`) lacks a comma after the Hollerith `1h(` and is rejected by current gfortran, so the reference build inserts that comma (originals kept as `*.orig`); the port behaves like the patched program. Result of the final regression run of the complete port: every file identical in 1 pass.
One pass. The reference shows `ncoeffs 181`, titles up to `(6,6,12;...)` in `dmso.SHARM.h02`, and an entropy output after 40 trials; all files are identical, i.e. every coefficient agrees to the last printed digit. Reference build: `sharm`/`sdf` were rebuilt with one comma inserted in the run-time title format (`'(1h(,'//il1fmt`), since gfortran 15.2 otherwise stops with "Missing comma between descriptors" before writing the coefficients; all other programs are the unmodified reference binaries. Also masked: characters 31-45 of the `*.mapgr.g01` data lines, the error column of `g_centres(r)`, which `mapgrquad.f` never sets (uninitialised in the Fortran, 0 in the port).

References

[1] A. K. Soper, "Empirical Potential Structure Refinement - EPSRshell: a user's guide", Technical Report RAL-TR-2011-012, STFC Rutherford Appleton Laboratory (2011).
[2] H. J. C. Berendsen, J. R. Grigera and T. P. Straatsma, "The missing term in effective pair potentials", J. Phys. Chem. 91, 6269-6271 (1987).
[3] A. K. Soper and A. Luzar, "A neutron diffraction study of dimethyl sulphoxide-water mixtures", J. Chem. Phys. 97, 1320-1331 (1992).
[4] A. K. Soper, "Orientational correlation function for molecular liquids: The case of liquid water", J. Chem. Phys. 101, 6888-6901 (1994).
[5] C. G. Gray and K. E. Gubbins, Theory of Molecular Fluids, Vol. 1: Fundamentals, Clarendon Press, Oxford (1984).
[6] A. Baranyai and D. J. Evans, "Direct entropy calculation from computer simulation of liquids", Phys. Rev. A 40, 3817-3822 (1989).
[7] S. K. Park and K. W. Miller, "Random number generators: good ones are hard to find", Commun. ACM 31, 1192-1201 (1988).
[8] GNU Fortran (gfortran) 15.2, GCC, https://gcc.gnu.org/fortran/ - reference build with -O2 -fno-tree-vectorize.