EPSR-test-ex90 Harmonic repulsive core (reppottype harmonic)

Program(s): epsr | chained passes: 3 | verification: ALL IDENTICAL (all 3 passes)

What this case tests

Exercises the alternative, harmonic form of the short-range repulsive core of the `epsr` engine of EPSR [1,2], selected by `reppottype harmonic` in the `.pcof` file instead of the default exponential core. The core keeps molecules apart at distances below `rminex` and its amplitude `pcoefex` is adjusted from the coordination number `coordex` between iterations [3].

Code exercised

Fortran (original, fortran_src/):

- `epsr_standalone_rev3.f`: `inita` (`reppottype` -> `harmonic=.true.`), `generate_rp` (harmonic core, quadratic branch), `repulsivept` (feedback of `pcoefex`)
- `read_and_save_files.f`: `read_pcof`, `save_files_others` (`.pcof` read and written back)
- `ato_manipulation_rev4.f`: `fmolec1`, `atogen`; `molfit.f`: `en_change` (energies from the tabulated potential)

Rust (port, rust/epsr-core/src/):

- `structure.rs`: `inita`; `potential.rs`: `generate_rp`, `repulsivept`
- `pcof.rs`: `read_pcof`, `write_pcof_pairs`
- `energy.rs`: `fmolec1`, `en_change`; `structure.rs`: `atogen`

Physical system

The EPSR-test system: a liquid mixture of 200 dimethyl sulphoxide (DMSO, (CH3)2SO) and 400 water molecules (DMSO mole fraction 1/3; 3200 atoms) at 300 K in a cubic box of side 33.12 Å (atomic number density 0.0881 atoms/Å^3). DMSO uses EPSR's Lennard-Jones + Coulomb reference potential with methyl rotational groups; water uses SPC/E Lennard-Jones and charge parameters [4] with EPSR's flexible intramolecular geometry (O-H 0.976 Å). The empirical potential is refined against three neutron diffraction data sets of deuterated DMSO in D2O, in null-scattering H2O/D2O and in H2O (ISIS SANDALS runs SLS03863, SLS03871, SLS03850), the isotope-substitution approach of Soper and Luzar [5]. For this case the repulsive core is harmonic: 0.5·pcoefex·ddiff^2 with ddiff = (rminex + expecfuse - r)/expecfuse and expecfuse = expecf·sqrt(rminex), applied on three pairs (`rminex pcoefex coordex`: S-S `4.0 0 0.1`, O1-O1 `2.6 0.5 0`, O1-H1 `1.5 0.2 0.05`).

Inputs

Made from EPSR-test with `scripts/make_case.py` (keyword and line edits): `.pcof` keyword `reppottype harmonic` and the repulsive-core lines of EPSR-test-ex1 (`rminex pcoefex coordex`: S-S `4.0 0 0.1`, O1-O1 `2.6 0.5 0`, O1-H1 `1.5 0.2 0.05`). All other files are those of EPSR-test.

How the accuracy was verified

`commands.txt` (1 program call, repeated 3 times, each pass reading the files written by the previous one) was run with the Fortran reference build (`rust/tools/build_fortran_reference.sh`: the original sources, gfortran -O2 -fno-tree-vectorize, single thread) and with the Rust `epsr-rs` on identical copies of the inputs, and every file written was compared byte for byte (`rust/tools/run_ex_test.sh`, `compare_outputs.py`; only the date line of `.out` and the wall-clock columns of `fort.35` are masked). The Fortran output is stored in `fortran_reference.tar.gz`. Result of the final regression run of the complete port: every file identical in all 3 passes.
Each pass reads back the `.ato`, `.pcof`, `.EPSR.inp`, `.inpa` and accumulators of the previous one, so the S-S `pcoefex` driven from 0 by `repulsivept` (-4.05E-03 after pass 3 in the stored Fortran `.pcof`) feeds back into `generate_rp`. Only the quadratic branch is active (second `expecf` value 0, no gradient limit); the results differ from EPSR-test-ex1, which uses the same `.pcof` lines with the exponential core. The symbol value `harmonic` is written back unchanged.

References

[1] A. K. Soper, "Empirical potential Monte Carlo simulation of fluid structure", Chem. Phys. 202, 295-306 (1996).
[2] A. K. Soper, "Partial structure factors from disordered materials diffraction data: An approach using empirical potential structure refinement", Phys. Rev. B 72, 104204 (2005).
[3] A. K. Soper, "Empirical Potential Structure Refinement - EPSRshell: a user's guide", Technical Report RAL-TR-2011-012, STFC Rutherford Appleton Laboratory (2011).
[4] H. J. C. Berendsen, J. R. Grigera and T. P. Straatsma, "The missing term in effective pair potentials", J. Phys. Chem. 91, 6269-6271 (1987).
[5] A. K. Soper and A. Luzar, "A neutron diffraction study of dimethyl sulphoxide-water mixtures", J. Chem. Phys. 97, 1320-1331 (1992).
[6] GNU Fortran (gfortran) 15.2, GCC, https://gcc.gnu.org/fortran/ - reference build with -O2 -fno-tree-vectorize.