

Ryan Chasse

LHP · PRO

MOVER TYPE · LOW CONFIDENCE

Quiet-Trunk (expression, not capacity) / drifting toward Arm Thrower; the post has to be fixed first

Elite engine, bottom-of-cohort trunk speed, and a front leg that catches hard instead of posting.

WEIGHT

122 kg

TEST DATE

2026-09-15

CAPTURES

4 captures ·
2024-09-10,
2025-01-27,
2025-10-24,
2026-09-15

VELOCITY

87.5 mph

MEAN VELOCITY BY CAPTURE

89.4

2024-09-10

91.46

2025-01-27

91.875

2025-10-24

87.5

2026-09-15

What Kind of Mover This Is

Chasse is down 4.4 mph from his best capture, not the 1.9 the coach page reports — the headline compares his first capture to his last while labelling it as the last-two comparison, and the report's own appendix carries the right number (R3 to R4, delta velo -4.375). What sits under the drop is consistent across two independent witnesses: the rotational engine got stronger and the trunk got slower. Proteus shotput power, velocity and acceleration all climbed 20-38% through 2026; peak thorax speed on the mound is 4th percentile in a 23-man PRO cohort. That divergence is an expression problem, not a capacity problem, and the front leg is why. He now catches the ground violently and early — RFD around 111 BW/s (elite is 80), peak at 54% of foot-contact-to-release, which is the sweet spot — but the vertical and braking peaks are about 50 ms apart on three of four pitches (over 40 ms is clearly desynchronised), the lead knee extends roughly 4 degrees through the block against a 15-25 degree band, and total front-leg force impulse is 0.176 BW-s against a cohort median of 0.202 — 17th percentile on the single most reliable force metric on the page (ICC 0.99). Big collision, small delivery.

Why This Type

Peak thorax angular velocity 919 deg/s vs cohort median 1,100 — 4th percentile of 23 PRO pitchers (ICC 0.78).

Peak pelvis angular velocity 663 deg/s vs 710 — 22nd percentile (ICC 0.91). Pelvis-to-thorax amplification 1.39x against a cohort ratio near 1.55x: the trunk is the link that fails to amplify.

Proteus rotational output is rising hard over the same window — shotput best power 434 W (Jan 2026) to 531 W (Sep 2026), velocity 5.42 to 6.63 m/s, acceleration 21.4 to 29.5 m/s²; D2 extension power 100.5 to 143.4 W. Capacity is present and is not reaching the ball.

Front-knee extension footstrike to release is roughly 4 deg (51.7 to 47.6, X) against a 15-25 deg elite band, and the knee is MORE flexed at MER (57.2 deg, 91st pct) than at footstrike. He collapses then partially recovers.

Drop jump over the same year: RSI 3.23 to 1.93, ground contact 0.427 to 0.540 s, jump height 24.4 to 20.1 in. The landing-side leg (right, LHP) develops power at 42,990 W/s against 76,290 on the drive side while producing slightly MORE peak force. Absorbs instead of returns, on and off the mound.

Shoulder max internal-rotation torque 0.081 vs cohort median 0.066 (91st pct, ICC 0.92) while throwing 87.5 mph. High stress per mph — the arm is paying the upstream tax.

Does the Testing Agree With the Delivery?

Each row asks one question of the delivery and the same question of the testing that happens away from the mound. Rows that agree are the ones to act on; rows that diverge are where the capture and the athlete disagree, and that gap is usually the finding.

QUESTION	ON THE MOUND	OFF THE MOUND	READ
Can he post the front side? Both witnesses say the same thing on the same leg: it absorbs and does not return. This is the finding of the report.	Lead knee 51.7 deg at footstrike, 57.2 at MER, 47.6 at release — ~4 deg of net extension against a 15-25 deg band; knee flexion at release 87th-91st percentile (more bent than the group)	Drop jump RSI 3.23 to 1.93, ground contact 0.427 to 0.540 s, height 24.4 to 20.1 in; landing-side (right) single-leg RPD 42,990 W/s vs 76,290 on the drive side	AGREE
Is rotational power available? The engine is bigger than it was and less of it is reaching the ball. Expression problem. More gym work on rotation will not move this.	Peak thorax 919 deg/s, 4th percentile; pelvis-to-thorax amplification 1.39x	Proteus shotput best power 434 to 531 W, velocity 5.42 to 6.63 m/s, acceleration 21.4 to 29.5 m/s ² ; D2 extension power 100.5 to 143.4 W across 2026	DIVERGE
Is the ground half strong enough? Raw force production is not the constraint. Do not spend the block here.	Drive-leg peak vertical 1.73 BW (typical band 1.3-1.6), 68th percentile at ICC 0.99; lead peak resultant 2.21 BW vs cohort 2.23	CMJ height flat across five dates (17.9 to 18.7 in); CMJ force at peak power flat 2,686 to 2,606 N	AGREE
Does the force get into the ball? Strong, reliable on-mound signal with no testing counterpart. It is the highest-value row on the page and the appendix is where it lives.	Vertical-to-braking peak lag -50, -53, -50, -3 ms across four pitches (over 40 ms = desynchronised); resultant impulse 0.176 BW-s vs 0.202 median, 17th percentile, ICC 0.99; force-into-ball at release 0th percentile (ICC 0.95)	No off-mound proxy for peak synchrony — this one only the capture can see	UNKNOWN
Is the arm restricted, or just late? Not a mobility problem. The arm is late and overloaded, not restricted. Mobility is a single date eleven months old — context, not a trend.	Shoulder max int-rot torque 91st percentile at 87.5 mph; elbow angle at footstrike 112 deg, dead median	Mobility 2025-10-24: shoulder ER 105 deg, IR 57 deg, total arc 162 deg — a normal arc; hip IR 33 left / 32 right, ER 38/38	DIVERGE
Is the distal end carrying the chain?	Hand rotation speed at release 4,150 deg/s vs 3,510 median (83rd pct) while thorax sits at 4th percentile	Plyo push-up climbing on every stable column — height 5.05 to	AGREE

QUESTION	ON THE MOUND	OFF THE MOUND	READ
Upper body is getting stronger and is being asked to rescue the chain. This is the drift toward Arm Thrower and it is the injury-cost line.		6.15 in, force at peak power 1,062 to 1,949 N	

Who He Should Be Measured Against

PRO cohort, ranked on ICC-clean metrics only · n=23

23-24 athletes with one row each — a percentile here is a rank with a percentage sign. Only metrics with ICC above ~0.5 are quoted; the front-leg fm_ pipeline rows on the coach page carry reliability 0.32-0.41 and are excluded.

Two force pipelines disagree with each other on the same athlete. The fm_ rows put his front-leg peak vertical at 91st percentile (1.89 BW vs a 1.13 median, reliability 0.36); the kin_PROCESSED rows put the same leg at 48th percentile (1.95 BW vs a 1.95 median, ICC 0.99). Same for the drive leg: 100th percentile in one table, 68th in the other, from the same 1.73 BW value. The fm_ cohort medians are implausible against the framework's own thresholds, so the high-ICC table is the one to believe.

METRIC	HIS	GROUP MEDIAN	%ILE	NOTE
Peak thorax angular velocity	919 deg/s	1,100	4.3	ICC 0.78 — the weak link in the chain
Peak pelvis angular velocity	663 deg/s	710	21.7	ICC 0.91
Front-leg resultant impulse	0.176 BW-s	0.202	17.4	ICC 0.99 — most reliable force row on the page
Front-leg peak resultant	2.21 BW	2.23	47.8	ICC 1.00 — solid (2.0+), not elite (2.5+), exactly average here
Front-leg peak vertical	1.95 BW	1.95	47.8	ICC 0.99
Front-leg peak braking	1.10 BW	1.24	69.6	ICC 0.99; percentile is on a signed negative value — the magnitude is below the median
Drive-leg peak vertical	1.73 BW	1.54	68.2	ICC 0.99; the coach page's 100th-percentile version comes from the low-reliability table
Shoulder max internal-rotation torque	0.081	0.066	91.3	ICC 0.92 — high stress at 87.5 mph
Lead knee angle at release (X)	47.6 deg	34.7	87.0	ICC 0.86 — more flexed at release than the group
Force into ball at release	0.166	0.222	0.0	ICC 0.95 — bottom of the cohort
Body weight	122 kg	97.5	100.0	Heaviest in the group — this is why every raw-newton headline reads as a strength

What This Mover Should Chase

The post on the landing (right) leg — everything else is behind this

Iso holds in the split/landing position and plyo drops onto the right leg, loaded for return rather than absorption. The number that has to move is drop-jump ground contact time: 0.540 s back under 0.45 s, with RSI back above 2.5. On the mound the read-across is lead-knee net extension from footstrike to release, currently ~4 deg; 12 deg would be a real change. Re-test the drop jump and single-leg vertical at six weeks.

Peak synchrony on the front leg

Vertical and braking peaks are ~50 ms apart on three of four pitches. Target is under 20 ms. This is a landing-intent and foot-placement problem more than a strength one — sprint mechanics and controlled decel work into the same position. The metric to watch is the vertical-to-braking peak lag row in the pitch-to-pitch table, per pitch, not the session average.

Trunk against the post — but only once the post exists

Med-ball rotational throws and connection ball, staged after the block holds. Doing this first is the classic error with this profile: there is nothing to rotate around yet. The target is peak thorax angular velocity 919 to 1,050+ deg/s with pelvis unchanged, i.e. amplification 1.39x toward 1.55x.

What To Leave Alone

Coaching these would move him toward a type he is not, and cost the thing he is good at.

Do not add rotational power work

Proteus shotput power, velocity and acceleration are all up 20-38% across 2026 and D2 extension power is up 43%. The engine is not the limiter. Adding load here makes the divergence worse and eats training time that the post needs.

Do not touch the drive leg

1.73 BW peak vertical against a 1.3-1.6 typical band, 68th percentile on the reliable table. It is doing its job. A well-meaning read of the '100th percentile, best in the group' headline into 'this is his weapon, build on it' spends a block on the one thing that is already fine.

Do not chase velocity intent yet

Shoulder internal-rotation torque is 91st percentile at 87.5 mph. Stress per mph is already high because the arm is covering for the trunk. Velo work before the block closes buys the tax, not the mph.

Do not coach the front-leg force down

RFD near 111 BW/s and peak at 54% of the foot-contact-to-release window are both in the elite range. He produces force early and fast. The problem is that the peaks do not stack and the knee does not hold, not that the force is wrong.

What To Watch

Body weight against the BW-normalised numbers

122 kg is the heaviest in the cohort and it is the divisor on every BW metric on the page. If the stored weight is stale, every force read moves with it. Confirm it before the next capture.

Pitch 4 of the focus session

P4 shows front-leg vertical force rate of 9.1 BW/s against 111, 79 and 122 on the other three, a doubled impulse, and force peak count 2 where the others are 1. Either a collapse-and-reload or an extraction failure. It drags every session-level aggregate on the page, including the peak-lag average that flatters the desynchrony from -50 ms to -39 ms.

The force pipeline break

12 metrics shrank by a factor of 1,094 between 2025-01-27 and 2025-10-24, and the 2025-10-24 front-leg values imply under 1.0 BW for a PRO pitcher, which is not physically credible. The 2026 force numbers are usable in absolute terms and not usable as a change from 2025. Do not read 'front-leg force went up 2.5x' as improvement.

CMJ and single-leg work/power columns

CMJ work done runs 657 J, 231 J, 2,106 J, 3,715 J, 2,414 J while jump height stays flat at 18-19 in. That is the known pipeline discontinuity at 2026-03-09, not a training effect. Same break sits in single-leg work, W/kg and RPD. Jump heights, RSI, ground contact time and force values are the stable columns — read only those.

How Much of This to Believe

C The direction of this profile is well supported; the magnitude is not settled. Four pitches, one of them probably corrupt, across a 326-day gap that contains a season and a training block, with every 'usual variation' band estimated from within-session spread.

Before acting on this:

- 4 pitches in the focus session; the report itself says 8 is where a ranked scan starts to mean anything.
- 740 of 740 measures have provisional bands built from pitch-to-pitch spread, so every change reads larger than it is.
- 12 metrics crossed a 1,094x pipeline break between 2025-01-27 and 2025-10-24 and cannot be differenced.
- Two force pipelines disagree on this athlete's cohort position; the low-ICC one drives the coach-page headlines.
- The velocity headline (-1.9 mph) is a first-to-last comparison labelled as a last-two comparison; the real last-two number is -4.375 mph and sits in the appendix.
- Mobility is a single date from 2025-10-24 — context, not a trend.
- 18 assessment dates fell outside every capture's +/-60-day window and were excluded, including 8 Proteus dates.

In One Paragraph

Primarily a Quiet-Trunk mover with an expression problem rather than a capacity problem, drifting toward Arm Thrower because the front side is not posting. The engine is measurably bigger than it was a year ago and the trunk is slower; between them sits a lead leg that catches the ground hard and early but does not extend against it, with vertical and braking peaks 50 ms apart and bottom-quintile force impulse. Fix the post first, then the peak synchrony, then trunk-against-post. Leave the drive leg, the rotational gym work and the velocity intent alone until the block holds. One full-intent capture of 8-10 pitches, plus a drop jump and single-leg vertical on the same day, settles almost all of this.