

## Part IA: Mathematics for Natural Sciences A

### Examples Sheet 11: Linear ordinary differential equations

### Model Solutions

*Please send all comments and corrections to [jmm232@cam.ac.uk](mailto:jmm232@cam.ac.uk).*

#### Basic definitions

1. Consider the general linear  $n$ th-order ordinary differential equation:

$$\alpha_n(x) \frac{d^n y}{dx^n} + \alpha_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + \alpha_1(x) \frac{dy}{dx} + \alpha_0(x)y = f(x).$$

where  $\alpha_n(x) \neq 0$ .

- (a) Give the definitions of the following terms: (i) homogeneous equation; (ii) coefficient functions; (iii) forcing.
- (b) Define a *complementary function* for this equation. How many arbitrary constants feature in the complementary function for this equation?
- (c) Define a *particular integral* for this equation. Is a particular integral for this equation unique?
- (d) Show that if  $y_{CF}$  is the complementary function for this equation, and  $y_{PI}$  is a particular integral, then the sum  $y = y_{CF} + y_{PI}$  solves the equation.
- (e) Suppose that we now seek a particular solution of this equation satisfying certain boundary conditions. How many boundary conditions are needed to fully specify a particular solution?

#### ◆ Solution:

- (a) The *homogeneous equation* is the same differential equation, but with  $f(x) = 0$  on the right hand side. The *coefficient functions* are the functions  $\alpha_i(x)$  for  $i = 0, \dots, n$ . The *forcing* is the term on the right hand side,  $f(x)$ .
- (b) A complementary function  $y_{CF}$  is a solution of the homogeneous equation. The general complementary function contains two arbitrary constants of integration, corresponding to the fact we are dealing with a second-order ODE.
- (c) A particular integral  $y_{PI}$  is a solution of the inhomogeneous equation (i.e. including the forcing term). It is not unique, because we can add on parts of the complementary function and still solve the inhomogeneous equation.
- (d) If  $y_{CF}$  is the general complementary function, and  $y_{PI}$  is a particular integral, the sum satisfies:

$$\begin{aligned} & \alpha_n(x) \frac{d^n}{dx^n} (y_{CF} + y_{PI}) + \alpha_{n-1}(x) \frac{d^{n-1}}{dx^{n-1}} (y_{CF} + y_{PI}) + \cdots + \alpha_1(x) \frac{d}{dx} (y_{CF} + y_{PI}) + \alpha_0(x)(y_{CF} + y_{PI}) \\ &= \left( \alpha_n(x) \frac{d^n y_{CF}}{dx^n} + \alpha_{n-1}(x) \frac{d^{n-1} y_{CF}}{dx^{n-1}} + \cdots + \alpha_1(x) \frac{dy_{CF}}{dx} + y_{CF} \right) \\ & \quad + \left( \alpha_n(x) \frac{d^n y_{PI}}{dx^n} + \alpha_{n-1}(x) \frac{d^{n-1} y_{PI}}{dx^{n-1}} + \cdots + \alpha_1(x) \frac{dy_{PI}}{dx} + y_{PI} \right) \\ &= f(x), \end{aligned}$$

since the first term goes to zero as  $y_{CF}$  is the general complementary function, and the second term gives  $f(x)$  since  $y_{PI}$  is a solution of the inhomogeneous equation.

- (e) We need  $n$  boundary conditions to fix  $n$  arbitrary constants.

2. By direct differentiation, verify that the following ordinary differential equations have the given complementary functions:

- (a)  $y_{\text{CF}} = Ax + Be^x$  is the complementary function for  $(x - 1)y'' - xy' + y = 0$ ;
  - (b)  $y_{\text{CF}} = A + B \log(x)$  is the complementary function for  $xy'' + y' = \cos(x)e^{x^2}$ ;
  - (c)  $y_{\text{CF}} = Ax + B \sin(x)$  is the complementary function for  $(1 - x \cot(x))y'' - xy' + y = x$ ;
  - (d)  $y_{\text{CF}} = A + Bx + Ce^x$  is the complementary function for  $y''' - y'' = x$ .
- 

◆ Solution:

- (a) Computing the derivatives, we have  $y'_{\text{CF}} = A + Be^x$ ,  $y''_{\text{CF}} = Be^x$ . Substituting, we have:

$$(x - 1)Be^x - Ax - Bxe^x + Ax + Be^x = 0,$$

as required.

- (b) Computing the derivatives, we have  $y'_{\text{CF}} = B/x$ ,  $y''_{\text{CF}} = -B/x^2$ . Substituting into the homogeneous version of the equation, we have:

$$x \left( -\frac{B}{x^2} \right) + \frac{B}{x} = 0,$$

as required.

- (c) Computing the derivatives, we have  $y'_{\text{CF}} = A + B \cos(x)$ ,  $y''_{\text{CF}} = -B \sin(x)$ . Substituting into the homogeneous version of the equation, we have:

$$(1 - x \cot(x))(-B \sin(x)) - x(A + B \cos(x)) + Ax + B \sin(x) = 0,$$

as required.

- (d) Computing the derivatives, we have  $y'_{\text{CF}} = B + Ce^x$ ,  $y''_{\text{CF}} = Ce^x$ ,  $y'''_{\text{CF}} = Ce^x$ . Substituting into the homogeneous version of the equation, we have:

$$Ce^x - Ce^x = 0,$$

as required.

3. By direct differentiation, verify that the following ordinary differential equations have the given particular integrals:

- (a)  $y_{\text{PI}} = \cos(x)$  is a particular integral for  $-y'' + y = 2 \cos(x)$ ;
  - (b)  $y_{\text{PI}} = x^2$  is a particular integral for  $xy'' + y' = 4x$ ;
  - (c)  $y_{\text{PI}} = e^{x^2}$  is a particular integral for  $y''' - 2xy'' - 2y' - y = (4x - 1)e^{x^2}$ ;
  - (d)  $y_{\text{PI}} = \sin(x)/x$  is a particular integral for  $xy^{(4)} + 4y^{(3)} + xy^{(2)} + 2y^{(1)} + xy = \sin(x)$ .
- 

◆ Solution:

- (a) We have  $y'_{\text{PI}} = -\sin(x)$ ,  $y''_{\text{PI}} = -\cos(x)$ , hence  $-(-\cos(x)) + \cos(x) = 2\cos(x)$ , showing that  $y_{\text{PI}}$  is indeed a particular integral.
- (b) We have  $y'_{\text{PI}} = 2x$ ,  $y''_{\text{PI}} = 2$ , hence  $2x + 2x = 4x$ , showing that  $y_{\text{PI}}$  is indeed a particular integral.
- (c) We have  $y'_{\text{PI}} = 2xe^{x^2}$ ,  $y''_{\text{PI}} = (2 + 4x^2)e^{x^2}$ ,  $y'''_{\text{PI}} = (12x + 8x^3)e^{x^2}$ . Substituting into the equation, we have:

$$(12x + 8x^3)e^{x^2} - 2x(2 + 4x^2)e^{x^2} - 2(2xe^{x^2}) - e^{x^2} = (4x - 1)e^{x^2},$$

showing that  $y_{\text{PI}}$  is indeed a particular integral.

- (d) To help, observe that:

$$\frac{d}{dx} \left( \frac{\sin(x)}{x^n} \right) = \frac{\cos(x)}{x^n} - \frac{n \sin(x)}{x^{n+1}}, \quad \frac{d}{dx} \left( \frac{\cos(x)}{x^n} \right) = -\frac{\sin(x)}{x^n} - \frac{n \cos(x)}{x^{n+1}}.$$

Then, we have:

$$\begin{aligned} y_{\text{PI}}^{(1)} &= \frac{\cos(x)}{x} - \frac{\sin(x)}{x^2} \\ y_{\text{PI}}^{(2)} &= -\frac{\sin(x)}{x} - \frac{2 \cos(x)}{x^2} + \frac{2 \sin(x)}{x^3} \\ y_{\text{PI}}^{(3)} &= -\frac{\cos(x)}{x} + \frac{3 \sin(x)}{x^2} + \frac{6 \cos(x)}{x^3} - \frac{6 \sin(x)}{x^4} \\ y_{\text{PI}}^{(4)} &= \frac{\sin(x)}{x} + \frac{4 \cos(x)}{x^2} - \frac{12 \sin(x)}{x^3} - \frac{24 \cos(x)}{x^4} + \frac{24 \sin(x)}{x^5}. \end{aligned}$$

Summing with the required coefficients, we obtain the result.

4. Verify that the equation:

$$(3+x)y'' + (2+x)y' - y = x^2 + 6x + 6$$

has complementary function  $y_{\text{CF}}(x) = Ae^{-x} + B(x+2)$ . Hence, by finding a particular integral of the form

$$y_{\text{PI}}(x) = \alpha x^2 + \beta x + \gamma,$$

determine the full solution to the equation subject to the boundary conditions  $y(0) = 0$  and  $y'(0) = 1$ .

---

◆ **Solution:** Differentiating, we have  $y'_{\text{CF}} = -Ae^{-x} + B$ ,  $y''_{\text{CF}} = Ae^{-x}$ . Substituting into the equation, we have:

$$(3+x)Ae^{-x} + (2+x)(-Ae^{-x} + B) - (Ae^{-x} + B(x+2)) = 0,$$

so indeed this is the complementary function. Trialling  $y_{\text{PI}} = \alpha x^2 + \beta x + \gamma$  as a particular integral, we have:

$$(3+x)(2\alpha) + (2+x)(2\alpha x + \beta) - (\alpha x^2 + \beta x + \gamma) = x^2 + 6x + 6.$$

Comparing coefficients, we see that  $\alpha = 1$ ,  $6\alpha = 6$ , and  $6\alpha + 2\beta - \gamma = 6$ . So provided we choose  $2\beta - \gamma = 0$ , we will have found a particular integral. This corresponds to the freedom we have to add on any multiple of  $x+2$  to the particular integral, because  $x+2$  is in the complementary function:

$$y_{\text{PI}} = x^2 + \beta(x+2),$$

for arbitrary  $\beta$ . Taking  $\beta = 0$ , we have the complete solution:

$$y(x) = Ae^{-x} + B(x+2) + x^2.$$

The solution satisfying  $y(0) = 0$  and  $y'(0) = 1$  must obey  $A + 2B = 0$  and  $-A + B = 1$ . Hence  $A = -2/3$ ,  $B = 1/3$ . Thus the particular solution is:

$$y(x) = x^2 + \frac{1}{3}x + \frac{2}{3} - \frac{2}{3}e^{-x}.$$

**Constant coefficient equations**

5. Consider the linear second-order ordinary differential equation with *constant coefficients*:

$$\alpha \frac{d^2 y}{dx^2} + \beta \frac{dy}{dx} + \gamma y = f(x),$$

where  $\alpha, \beta, \gamma$  are *constants*, with  $\alpha \neq 0$ .

- (a) Show that  $y = e^{\mu x}$  solves the homogeneous equation if and only if  $\mu$  satisfies the *auxiliary equation*:

$$\alpha \mu^2 + \beta \mu + \gamma = 0.$$

Deduce the complementary function of the equation in the case that the auxiliary equation does not have repeated roots.

- (b) Show that if the auxiliary equation has a repeated root  $\omega$ , then the complementary function is given by  $y_{\text{CF}} = (Ax + B)e^{\omega x}$ .
- (c) How does these results generalise to  $n$ th order linear differential equations with constant coefficients?
- 

**• Solution:**

- (a) Substituting directly into the homogeneous version of the equation ( $f(x) = 0$ ), we have:

$$\alpha \mu^2 e^{\mu x} + \beta \mu e^{\mu x} + \gamma e^{\mu x} = 0,$$

which gives the auxiliary equation as required. The complementary function is therefore  $y_{\text{CF}} = Ae^{\omega_1 x} + Be^{\omega_2 x}$  where  $\omega_1, \omega_2$  are the distinct roots of the auxiliary equation.

- (b) In the case of a repeated root, we can check that  $y_{\text{CF}} = (Ax + B)e^{\omega x}$  solves the equation. We note that if there is a repeated root  $\omega$ , then the auxiliary equation takes the form  $\alpha \mu^2 + \beta \mu + \gamma = \alpha(\mu - \omega)^2 = \alpha \mu^2 - 2\omega \alpha \mu + \alpha \omega^2$ . Hence we substitute into the equation:

$$\frac{d^2 y}{dx^2} - 2\omega \frac{dy}{dx} + \omega^2 y = 0.$$

Substituting in to the equation, we require:

$$2A\omega e^{\omega x} + (Ax + B)\omega^2 e^{\omega x} - 2\omega A\omega e^{\omega x} - 2\omega^2 A x e^{\omega x} + \omega^2 (Ax + B)e^{\omega x} = 0.$$

This holds identically, and thus indeed this is the complementary function.

- (c) If there is an  $m$ -fold repeated root  $\omega$ , the corresponding term in the complementary function is

$$(A_0 + A_1 x + \dots + A_{m-1} x^{m-1})e^{\omega x},$$

which is an  $(m - 1)$ th degree polynomial multiplied by an exponential factor.

6. Determine the solutions of the following differential equations:

(a)  $y'' + 6y' + 5y = 0;$

(b)  $y'' + 3y' + 4y = 0;$

(c)  $y'' + 4y = x;$

(d)  $y'' - 2y' + 2y = 2x^2;$

(e)  $y'' + y = |x|;$

(f)  $y'' + 3y' + 2y = e^{-x};$

(g)  $y'' - 2y' + 5y = e^x \cos(2x);$

(h)  $y'' + 2y' + y = 2xe^{-x}.$ 

---

◆ Solution:

(a) The auxiliary equation is  $\mu^2 + 6\mu + 5 = 0$ , with roots  $-5, -1$ . Hence the solution is  $y = Ae^{-5x} + Be^{-x}$ .

(b) The auxiliary equation is  $\mu^2 + 3\mu + 4 = 0$ , which can be solved by the quadratic equation to give:

$$\mu = \frac{-3 \pm i\sqrt{7}}{2}.$$

Hence the solution is:

$$y = Ae^{-(3+i\sqrt{7})x/2} + Be^{-(3-i\sqrt{7})x/2}.$$

This can be simplified as:

$$\begin{aligned} y &= Ae^{-3x/2}e^{-i\sqrt{7}x/2} + Be^{-3x/2}e^{i\sqrt{7}x/2} \\ &= Ae^{-3x/2} \left( \cos(\sqrt{7}x/2) - i \sin(\sqrt{7}x/2) \right) + Be^{-3x/2} \left( \cos(\sqrt{7}x/2) + i \sin(\sqrt{7}x/2) \right) \\ &= A'e^{-3x/2} \cos(\sqrt{7}x/2) + B'e^{-3x/2} \sin(\sqrt{7}x/2), \end{aligned}$$

where we have defined  $A' = A + B$  and  $B' = -iA + iB$  as new constants. This is the more standard way of writing the solution.

(c) The auxiliary equation is  $\mu^2 + 4 = 0$ , with roots  $\pm 2i$ . Hence the complementary function is  $y_{\text{CF}} = Ae^{2ix} + Be^{-2ix} = A' \cos(2x) + B' \sin(2x)$ .

We now seek a particular integral, which we guess is of the form  $y_{\text{PI}} = \alpha x + \beta$ . We then have:

$$4(\alpha x + \beta) = x,$$

giving us  $\alpha = 1/4, \beta = 0$ . Thus the complete solution is:

$$y = A \cos(2x) + B \sin(2x) + \frac{1}{4}x.$$

(d) The auxiliary equation is  $\mu^2 - 2\mu + 2 = 0$ , which has roots  $\mu = 2 \pm i$ . Hence the complementary function is  $y_{\text{CF}} = e^{2x} (A \cos(x) + B \sin(x))$ .

We now seek a particular integral, which we guess is of the form  $y_{\text{PI}} = \alpha x^2 + \beta x + \gamma$ . We then have:

$$2\alpha - 2(2\alpha x + \beta) + 2(\alpha x^2 + \beta x + \gamma) = 2x^2.$$

Comparing coefficients, we see that  $\alpha = 1, \beta = 2, \gamma = 1$ . Thus the complete solution is:

$$y = e^{2x} (A \cos(x) + B \sin(x)) + x^2 + 2x + 1.$$

---

- (e) The auxiliary equation is  $\mu^2 + 1 = 0$ , which has roots  $\mu = \pm i$ . Hence the complementary function is  $y_{\text{CF}} = A \cos(x) + B \sin(x)$ .

We now seek a particular integral. Since the equation involves a modulus sign, we consider the regions  $x > 0$  and  $x < 0$  separately.

- In the region  $x > 0$ , guess  $y_{\text{PI}} = \alpha x + \beta$ . Then:

$$\alpha x + \beta = x,$$

giving  $\alpha = 1, \beta = 0$ .

- In the region  $x < 0$ , guess  $y_{\text{PI}} = \alpha x + \beta$ . Then:

$$\alpha x + \beta = -x,$$

giving  $\alpha = -1, \beta = 0$ .

So the particular integral is  $x$  when  $x > 0$  and  $-x$  when  $x < 0$ . This gives  $y_{\text{PI}} = |x|$ . Hence the complete solution is:

$$y = A \cos(x) + B \sin(x) + |x|.$$

- (f) The auxiliary equation is  $\mu^2 + 3\mu + 2 = 0$ , which has roots  $\mu = -2, -1$ . Thus the complementary function is  $y_{\text{CF}} = Ae^{-x} + Be^{-2x}$ . Since  $e^{-x}$  is already contained in the complementary function, as a particular integral we trial  $y_{\text{PI}} = \alpha xe^{-x}$ . This gives:

$$\alpha(-2e^{-x} + xe^{-x}) + 3\alpha(e^{-x} - xe^{-x}) + 2\alpha xe^{-x} = e^{-x}.$$

Comparing coefficients, we see that  $3\alpha - 2\alpha = 1$ , so that  $\alpha = 1$ . Thus we have complete solution:

$$y = Ae^{-x} + Be^{-2x} + xe^{-x}.$$

- (g) The auxiliary equation is  $\mu^2 - 2\mu + 5 = 0$ , which has roots  $\mu = 1 \pm 2i$ . Thus the complementary function is  $y_{\text{CF}} = e^x(A \cos(2x) + B \sin(2x))$ . Since the forcing is contained in the complementary function, for the particular integral we guess  $y_{\text{PI}} = xe^x(\alpha \cos(2x) + \beta \sin(2x))$ . Then:

$$y'_{\text{PI}} = e^x(\alpha \cos(2x) + \beta \sin(2x)) + xe^x((\alpha + 2\beta) \cos(2x) + (\beta - 2\alpha) \sin(2x))$$

$$y''_{\text{PI}} = e^x((2\alpha + 4\beta) \cos(2x) + (2\beta - 4\alpha) \sin(2x)) + xe^x((-3\alpha + 4\beta) \cos(2x) + (-4\alpha - 3\beta) \sin(2x)).$$

Combining these according to the equation, we see that:

$$4e^x(\beta \cos(2x) - \alpha \sin(2x)) = e^x \cos(2x).$$

Comparing coefficients, we see that  $4\beta = 1$  and  $-4\alpha = 0$ , giving  $\beta = 1/4$ . Thus the general solution is:

$$y = e^x(A \cos(2x) + B \sin(2x)) + \frac{1}{4}xe^x \sin(2x).$$

- (h) The auxiliary equation is  $\mu^2 + 2\mu + 1 = (\mu + 1)^2 = 0$ , which has a repeated root  $\mu = -1$ . This gives the complementary function  $y_{\text{CF}} = (A + Bx)e^{-x}$ .

Since  $xe^{-x}$  is already contained in the complementary function, we now consider  $x^2$  times the complementary function as a particular integral. This gives  $y_{\text{PI}} = (\alpha x^2 + \beta x^3)e^{-x}$ , which on differentiation yields:

$$y'_{\text{PI}} = (2\alpha x + 3\beta x^2)e^{-x} - (\alpha x^2 + \beta x^3)e^{-x}$$

$$y''_{\text{PI}} = (2\alpha + 6\beta x)e^{-x} - (4\alpha x + 6\beta x^2)e^{-x} + (\alpha x^2 + \beta x^3)e^{-x}.$$

Combining these according to the equation, we see that:

$$\alpha = 0, \quad \beta = \frac{1}{3}.$$

Hence the complete solution is:

$$y_{\text{CF}} = (A + Bx)e^{-x} + \frac{1}{3}x^3e^{-x}.$$

7. Determine the solutions of the following differential equations subject to the given constraints:

- (a)  $y'' - 4y' + 13y = 0$ , subject to  $y(0) = \pi$  and  $y(-\pi/2) = 1$ ;
  - (b)  $y'' - 4y' + 5y = 125x^2$ , subject to  $y(0) = 1$  and  $y(\pi/2) = \frac{25\pi^2}{4} + 20\pi + 22$ ;
  - (c)  $y'' + 7y' + 12y = 6$ , subject to  $y(0) = 0$  and  $y(1/3) = \frac{1-e^{-1}}{2}$ ;
  - (d)  $y'' + 7y' + 12y = 2e^{-3x}$ , subject to  $y(0) = 1$  and  $y'(0) = 0$ .
- 

◆ Solution:

- (a) The auxiliary equation is  $\mu^2 - 4\mu + 13 = 0$ , with roots  $\mu = 2 \pm 3i$ , which gives the complementary function  $y_{CF} = e^{2x}(A \cos(3x) + B \sin(3x))$ . Imposing the boundary data, we have  $A = \pi$  and:

$$1 = -Be^{-\pi} \quad \Rightarrow \quad B = -e^{\pi}.$$

Hence the particular solution is  $y(x) = e^{2x}(\pi \cos(3x) - e^{\pi} \sin(3x))$ .

- (b) The auxiliary equation is  $\mu^2 - 4\mu + 5 = 0$ , with roots  $\mu = 2 \pm i$ , hence the complementary function is  $y_{CF} = e^{2x}(A \cos(x) + B \sin(x))$ . Trialling  $y_{PI} = \alpha x^2 + \beta x + \gamma$  as a particular integral, we have:

$$2\alpha - 4(2\alpha x + \beta) + 5(\alpha x^2 + \beta x + \gamma) = 125x^2.$$

Comparing coefficients, we see that  $\alpha = 25$ ,  $\beta = 40$  and  $\gamma = 22$ . Hence we have the complete solution:

$$y(x) = e^{2x}(A \cos(x) + B \sin(x)) + 25x^2 + 40x + 22.$$

Imposing  $y(0) = 1$ , we have:

$$1 = A + 22 \quad \Rightarrow \quad A = -21.$$

Imposing  $y(\pi/2) = 25\pi^2/4 + 20\pi + 22$ , we have:

$$\frac{25\pi^2}{4} + 20\pi + 22 = e^{\pi}B + \frac{25\pi^2}{4} + 20\pi + 22 \quad \Rightarrow \quad B = 0.$$

Hence the particular solution is:

$$y(x) = 25x^2 + 40x + 22 - 21e^{2x} \cos(x).$$

- (c) The auxiliary equation is  $\mu^2 + 7\mu + 12 = 0$ , which has roots  $\mu = -3, -4$ . Hence the complementary function is  $y_{CF} = Ae^{-3x} + Be^{-4x}$ . The particular integral is obviously  $y_{PI} = 1/2$ . This gives the complete solution:

$$y(x) = Ae^{-3x} + Be^{-4x} + \frac{1}{2}.$$

Imposing the boundary conditions, we have  $y(0) = 0$  implying  $0 = A + B + 1/2$ . We also have  $y(1/3) = (1 - e^{-1})/2$  implying:

$$\frac{1 - e^{-1}}{2} = Ae^{-1} + Be^{-4/3} + \frac{1}{2}.$$

Comparing, we see that  $B = 0$ ,  $A = -1/2$ . This gives the solution:

$$y(x) = \frac{1}{2}(1 - e^{-3x}).$$



- (d) This part has the same complementary function as the previous part,  $y_{\text{CF}} = Ae^{-3x} + Be^{-4x}$ , but the particular integral should now be  $y_{\text{PI}} = \alpha xe^{-3x}$ . Inserting this into the equation, we have:

$$-6\alpha e^{-3x} + 9\alpha xe^{-3x} + 7\alpha e^{-3x} - 21\alpha xe^{-3x} + 12\alpha xe^{-3x} = 2e^{-3x}$$

we see that  $\alpha = 2$ , giving  $\alpha = 2$ , hence the complete solution is:

$$y(x) = Ae^{-3x} + Be^{-4x} + 2xe^{-3x}.$$

Imposing the boundary conditions, we have  $1 = A + B$  and  $0 = -3A - 4B + 2$ . These simultaneous equations have solution  $A = 2, B = -1$ . Hence the solution is:

$$y(x) = 2e^{-3x} - e^{-4x} + 2xe^{-3x}.$$

- 
8. Find the value of  $a$  for which the complementary function of the ODE:

$$\frac{d^2y}{dx^2} + 4\frac{dy}{dx} + ay = 0,$$

is given by  $y_{\text{CF}} = Axe^{-2x} + Be^{-2x}$ .

---

◆ **Solution:** The auxiliary equation is  $\mu^2 + 4\mu + a = 0$ , which has roots:

$$\mu = \frac{-4 \pm \sqrt{16 - 4a}}{2} = -2 \pm \sqrt{4 - a}.$$

There is a repeated root when  $a = 4$ , which gives rise to the solution  $y_{\text{CF}} = (Ax + B)e^{-2x}$ .

9. Find the general solution of the differential equation:

$$\frac{d^2y}{dx^2} + y = \cos(kx),$$

where  $k$  is a real number.

---

◆ **Solution:** The auxiliary equation is  $\mu^2 + 1 = 0$ , with roots  $\mu = \pm i$ . This gives complementary function  $y_{\text{CF}} = A \cos(x) + B \sin(x)$ .

In the case where  $k \neq \pm 1$ , a particular integral is obviously  $\cos(kx)/(1 - k^2)$ . Hence the complete solution is:

$$y = A \cos(x) + B \sin(x) + \frac{\cos(kx)}{1 - k^2}.$$

When  $k = \pm 1$ , we need a different particular integral. We guess  $y_{\text{PI}} = \alpha x \cos(x) + \beta x \sin(x)$ . Then:

$$y'_{\text{PI}} = \alpha \cos(x) + \beta \sin(x) - \alpha x \sin(x) + \beta x \cos(x),$$

$$y''_{\text{PI}} = -2\alpha \sin(x) + 2\beta \cos(x) - \alpha x \cos(x) - \beta x \sin(x).$$

Comparing coefficients, we see that  $\alpha = 0$  and  $\beta = 1/2$ . Thus the complete solution is:

$$y = A \cos(x) + B \sin(x) + \frac{1}{2}x \sin(x).$$

For those who are interested, this can be considered the limit as  $k \rightarrow 1$  of the previous case. Rewriting the constants in the form:

$$y = A' \cos(x) + B \sin(x) + \frac{\cos(kx) - \cos(x)}{1 - k^2}.$$

Taking the limit as  $k \rightarrow 1$ , we have:

$$\lim_{k \rightarrow 1} \frac{\cos(kx) - \cos(x)}{1 - k^2} = \lim_{k \rightarrow 1} \frac{-x \sin(kx)}{-2k} = \frac{1}{2}x \sin(x),$$

using L'Hôpital's rule in the final step. Nice that we can do this, huh!

10. The differential operator  $\mathcal{L}$  is defined by:

$$\mathcal{L} = \frac{d^2}{dx^2} + \sqrt{3} \frac{d}{dx} + 3.$$

Solve the equation  $\mathcal{L}y = 0$ , and hence solve the equations:

(a)  $\mathcal{L}y = e^{-\sqrt{3}x}$ ;

(b)  $\mathcal{L}y = x$ .

Without further calculation, state the general solution of  $\mathcal{L}y = 2x + e^{-\sqrt{3}x}$ . Find also the solution of this equation satisfying the boundary conditions:

$$y(0) = 0, \quad y(\pi) = \frac{e^{-\sqrt{3}\pi}}{3} - \frac{2}{3\sqrt{3}}.$$

---

◆ **Solution:** The solution to the equation  $\mathcal{L}y = 0$  is just the complementary function of the equation  $y'' + \sqrt{3}y' + 3y = 0$ . The auxiliary equation is  $\mu^2 + \sqrt{3}\mu + 3 = 0$ , which has roots  $\mu = (-\sqrt{3} \pm 3i)/2$ , hence the solution to  $\mathcal{L}y = 0$  is:

$$y(x) = e^{-\sqrt{3}x/2} (A \cos(3x/2) + B \sin(3x/2)).$$

In the given cases, we have:

(a) A particular integral for  $\mathcal{L}y = e^{-\sqrt{3}x}$  is  $y_{\text{pl}} = \alpha e^{-\sqrt{3}x}$ . Substituting into the equation, we have:

$$3\alpha e^{-\sqrt{3}x} - 3\alpha e^{-\sqrt{3}x} + 3\alpha e^{-\sqrt{3}x} = e^{-\sqrt{3}x},$$

so that  $\alpha = 1/3$ . Thus the general solution is:

$$y(x) = e^{-\sqrt{3}x/2} (A \cos(3x/2) + B \sin(3x/2)) + \frac{1}{3}e^{-\sqrt{3}x}.$$

(b) A particular integral for  $\mathcal{L}y = x$  is  $y_{\text{pl}} = \alpha x + \beta$ , which gives:

$$\sqrt{3}\alpha + 3\alpha x + 3\beta = x.$$

This implies  $\alpha = 1/3$ , and  $\sqrt{3}/3 + 3\beta = 0$ , so that  $\beta = -1/3\sqrt{3}$ . Thus the general solution is:

$$y(x) = e^{-\sqrt{3}x/2} (A \cos(3x/2) + B \sin(3x/2)) + \frac{1}{3}x - \frac{1}{3\sqrt{3}}.$$

The general solution of the equation  $\mathcal{L}y = 2x + e^{-\sqrt{3}x}$  is now given by linearity as:

$$y(x) = e^{-\sqrt{3}x/2} (A \cos(3x/2) + B \sin(3x/2)) + \frac{1}{3}e^{-\sqrt{3}x} + \frac{2}{3}x - \frac{2}{3\sqrt{3}}.$$

Imposing the boundary condition  $y(0) = 0$ , we have:

$$0 = A + \frac{1}{3} - \frac{2}{3\sqrt{3}} \quad \Rightarrow \quad A = \frac{2}{3\sqrt{3}} - \frac{1}{3}.$$

Imposing the boundary condition at  $x = \pi$ , we have:

$$\frac{e^{-\sqrt{3}\pi}}{3} - \frac{2}{3\sqrt{3}} = -Be^{-\sqrt{3}\pi/2} + \frac{1}{3}e^{-\sqrt{3}\pi} + \frac{2}{3}\pi - \frac{2}{3\sqrt{3}} \quad \Rightarrow \quad B = -\frac{2}{3}\pi e^{\sqrt{3}\pi/2}.$$

The particular solution is therefore:

$$y(x) = e^{-\sqrt{3}x/2} \left( \left( \frac{2}{3\sqrt{3}} - \frac{1}{3} \right) \cos(3x/2) - \frac{2}{3}\pi e^{\sqrt{3}\pi/2} \sin(3x/2) \right) + \frac{1}{3}e^{-\sqrt{3}x} + \frac{2}{3}x - \frac{2}{3\sqrt{3}}$$

## Harmonic oscillators

11. Consider the constant coefficient linear second-order ordinary differential equation:

$$\frac{d^2 y}{dt^2} + 2\gamma \frac{dy}{dt} + \omega_0^2 y = f(t),$$

modelling an oscillating system which depends on time  $t$ . The coefficients  $\gamma, \omega_0$  are positive.

- What is the physical interpretation of the constant  $\gamma$ ? What is the physical interpretation of the function  $f(t)$ ?
- Find the complementary function of this equation. Discuss the different forms the complementary function can take (in particular, defining the terms *underdamping*, *critical damping*, and *overdamping*), and how this relates to the *transient* behaviour of the oscillator.
- In the underdamped case, find the long-term behaviour of the oscillator in the case of resonant forcing:

$$f(t) = e^{-\gamma t} \sin \left( t \sqrt{\omega_0^2 - \gamma^2} \right).$$

### ◆ Solution:

- $\gamma$  is the strength of the damping of the oscillator. The function  $f(t)$  is the external driving force exerted on the oscillator per unit mass.
- The auxiliary equation is:

$$\mu^2 + 2\gamma\mu + \omega_0^2 = 0 \quad \Rightarrow \quad \mu = \frac{-2\gamma \pm \sqrt{4\gamma^2 - 4\omega_0^2}}{2} = -\gamma \pm \sqrt{\gamma^2 - \omega_0^2}.$$

- If  $\gamma^2 > \omega_0^2$ , then the complementary function takes the form:

$$y_{CF} = Ae^{(-\gamma + \sqrt{\gamma^2 - \omega_0^2})t} + Be^{(-\gamma - \sqrt{\gamma^2 - \omega_0^2})t}.$$

This is the case of *overdamping*. The displacement  $y$  tends to zero exponentially without oscillation. In the general case, there are two modes of decay, with the slower decay happening at a rate  $-\gamma + \sqrt{\gamma^2 - \omega_0^2}$  per second.

- If  $\gamma^2 = \omega_0^2$ , then the complementary function takes the form:

$$y_{CF} = (A + Bt)e^{-\gamma t}.$$

This is the case of *critical damping*. The displacement  $y$  tends to zero exponentially without oscillation. There is only one mode of decay with rate  $\gamma$ , so that for general initial conditions, the decay is faster than the case of overdamping.

- If  $\gamma^2 < \omega_0^2$ , then the complementary function takes the form:

$$y_{CF} = e^{-\gamma t} \left( A \cos \left( t \sqrt{\omega_0^2 - \gamma^2} \right) + B \sin \left( t \sqrt{\omega_0^2 - \gamma^2} \right) \right).$$

This is the case of *underdamping*. The displacement  $y$  tends to zero exponentially with oscillation.

- In this case, the complementary function contains the forcing, so we should trial a solution of the form

$$y_{PI} = te^{-\gamma t} \left( \alpha \cos \left( t \sqrt{\omega_0^2 - \gamma^2} \right) + \beta \sin \left( t \sqrt{\omega_0^2 - \gamma^2} \right) \right)$$

Differentiating, we have:

$$y'_{\text{PI}} = e^{-\gamma t} \left[ (1 - \gamma t) \left( \alpha \cos \left( t \sqrt{\omega_0^2 - \gamma^2} \right) + \beta \sin \left( t \sqrt{\omega_0^2 - \gamma^2} \right) \right) + t \sqrt{\omega_0^2 - \gamma^2} \left( -\alpha \sin \left( t \sqrt{\omega_0^2 - \gamma^2} \right) + \beta \cos \left( t \sqrt{\omega_0^2 - \gamma^2} \right) \right) \right].$$

and:

$$y''_{\text{PI}} = e^{-\gamma t} \left\{ [-2\gamma + t(2\gamma^2 - \omega_0^2)] \left( \alpha \cos \left( t \sqrt{\omega_0^2 - \gamma^2} \right) + \beta \sin \left( t \sqrt{\omega_0^2 - \gamma^2} \right) \right) + 2(1 - \gamma t) \sqrt{\omega_0^2 - \gamma^2} \left( -\alpha \sin \left( t \sqrt{\omega_0^2 - \gamma^2} \right) + \beta \cos \left( t \sqrt{\omega_0^2 - \gamma^2} \right) \right) \right\}.$$

Putting everything together, we have:

$$2\sqrt{\omega_0^2 - \gamma^2} e^{-\gamma t} \left( -\alpha \sin \left( t \sqrt{\omega_0^2 - \gamma^2} \right) + \beta \cos \left( t \sqrt{\omega_0^2 - \gamma^2} \right) \right) = e^{-\gamma t} \sin \left( t \sqrt{\omega_0^2 - \gamma^2} \right).$$

Comparing coefficients, we see that  $\beta = 0$  and:

$$\alpha = -\frac{1}{2\sqrt{\omega_0^2 - \gamma^2}}.$$

Hence the dominant long-term behaviour of the oscillator in this case is:

$$y_{\text{PI}} = -\frac{1}{2\sqrt{\omega_0^2 - \gamma^2}} t e^{-\gamma t} \cos \left( t \sqrt{\omega_0^2 - \gamma^2} \right),$$

which is an exponentially decaying curve, but one that is enhanced by the linear polynomial term. Note also that it is a cosine response to a sine forcing - there is a  $\pi/2$  phase shift between the forcing and response of the oscillator.

**Coupled systems of differential equations**

12.

(a) Consider the system of differential equations:

$$\frac{dx}{dt} = ax + by + p, \quad \frac{dy}{dt} = cx + dy + q,$$

for the variables  $x(t), y(t)$ , where  $a, b, c, d, p, q$  are constants. Show that:

$$\frac{d^2x}{dt^2} = (a + d)\frac{dx}{dt} + (bc - ad)x + bq - pd.$$

(b) Hence:

(i) Find the general solution of the system:

$$\frac{dx}{dt} = 4y + 2, \quad \frac{dy}{dt} = x.$$

(ii) Solve the system:

$$\frac{dx}{dt} = 3x - y, \quad \frac{dy}{dt} = x + y,$$

subject to the initial conditions  $x(0) = 0$  and  $y(0) = 1$ .

(iii) Solve the system:

$$\frac{dx}{dt} = -3x + y, \quad \frac{dy}{dt} = -5x + y,$$

subject to the initial conditions  $x(0) = 1, y(0) = 1$ .

---

**◆ Solution:**

(a) Differentiating the first equation, we have:

$$\frac{d^2x}{dt^2} = a\frac{dx}{dt} + b\frac{dy}{dt}.$$

Multiplying the second equation through by  $b$ , and inserting it into this equation, we have:

$$\frac{d^2x}{dt^2} = a\frac{dx}{dt} + bcx + bdy + bq.$$

Finally, note from the original first equation that  $by = -dx/dt - ax - p$ . So substituting for  $by$ , we have:

$$\frac{d^2x}{dt^2} = a\frac{dx}{dt} + bcx - d\frac{dx}{dt} - adx - dp + bq.$$

Grouping terms, we have:

$$\frac{d^2x}{dt^2} = (a + d)\frac{dx}{dt} + (bc - ad)x + bq - pd,$$

as required.

- (b) (i) Here, we have  $a = 0, b = 4, p = 2$  and  $c = 1, d = 0, q = 0$ . Thus the second-order equation we derived for  $x$  becomes:

$$\frac{d^2x}{dt^2} = 4x.$$

This has auxiliary equation  $\mu^2 - 4 = 0$  with roots  $\mu = \pm 2$ , hence the general solution of this equation for  $x$  is:

$$x(t) = Ae^{-2t} + Be^{2t}.$$

We must now insert this back into the original system for two reasons: (i) to check that we actually do solve the system for all  $A, B$  and that they are not fixed by some other conditions; (ii) to find the expression for  $y$ . Inserting back into the first original equation, we can derive an expression for  $y$ :

$$-2Ae^{-2t} + 2Be^{2t} = 4y + 2 \quad \Rightarrow \quad y(t) = -\frac{1}{2}Ae^{-2t} + \frac{1}{2}Be^{2t} - \frac{1}{2}.$$

It remains to check the consistency with the original second equation. Inserting, we have:

$$Ae^{-2t} + Be^{2t} = Ae^{-2t} + Be^{2t}.$$

So, all is well. Thus the general solution is:

$$x(t) = Ae^{-2t} + Be^{2t}, \quad y(t) = -\frac{1}{2}Ae^{-2t} + \frac{1}{2}Be^{2t} - \frac{1}{2},$$

for arbitrary constants  $A, B$ .

- (ii) We employ a similar method. In this case, we have  $a = 3, b = -1, p = 0$  and  $c = 1, d = 1, q = 0$ . Thus the second order equation for  $x$  becomes:

$$\frac{d^2x}{dt^2} = 4\frac{dx}{dt} - 4x.$$

The auxiliary equation is  $\mu^2 - 4\mu + 4 = 0$ , which has a repeated roots  $\mu = 2$ . Thus the solution for  $x(t)$  is  $x(t) = (A + Bt)e^{2t}$ . Imposing the boundary data  $x(0) = 0$ , we see that  $x(t) = Bte^{2t}$ .

Inserting into the first original equation, we can determine  $y(t)$ . We have:

$$2Bte^{2t} + Be^{2t} = 3Bte^{2t} - y \quad \Rightarrow \quad y(t) = Bte^{2t} - Be^{2t}.$$

Imposing  $y(0) = 1$ , we see that  $B = -1$ , so that  $y(t) = (1 - t)e^{2t}$ . Finally, checking the consistency with the second original equation, we see that:

$$-e^{2t} + 2(1 - t)e^{2t} = -te^{2t} + (1 - t)e^{2t},$$

as anticipated. Hence the solution is:

$$x(t) = -te^{2t}, \quad y(t) = (1 - t)e^{2t}.$$

- (iii) For our final system, we have  $a = -3, b = 1, p = 0$  and  $c = -5, d = 1, q = 0$ . Thus the second order equation for  $x$  is:

$$\frac{d^2x}{dt^2} = -2\frac{dx}{dt} - 2x.$$

The auxiliary equation is  $\mu^2 + 2\mu + 2 = 0$ , which has roots  $\mu = -1 \pm i$ . Hence, we have:

$$x(t) = e^{-t} (A \cos(t) + B \sin(t)).$$

Imposing the boundary condition  $x(0) = 1$ , we see that  $A = 1$ . Differentiating  $x(t)$  to obtain  $y(t)$  from the first original equation, we have:

$$-e^{-t} (\cos(t) + B \sin(t)) + e^{-t} (-\sin(t) + B \cos(t)) = -3e^{-t} (\cos(t) + B \sin(t)) + y(t).$$

Rearranging, we see that:

$$y(t) = e^{-t} ((B + 2) \cos(t) + (2B - 1) \sin(t)).$$

Imposing  $y(0) = 1$ , we see that  $B = -1$ . Thus the complete solution is:

$$\begin{aligned} x(t) &= e^{-t} (\cos(t) - \sin(t)), \\ y(t) &= e^{-t} (\cos(t) - 3 \sin(t)). \end{aligned}$$