

Tucatinib in Human Breast Cancer

Drug Summary: Tucatinib (trade name *Tukysa*) is an orally active, highly selective small-molecule tyrosine kinase inhibitor of the human epidermal growth factor receptor-2 (HER2/ERBB2). By binding to HER2's intracellular kinase domain, tucatinib inhibits HER2 phosphorylation and downstream signaling (MAPK/ERK and PI3K/AKT pathways), blocking tumor cell proliferation ¹ ². Tucatinib also inhibits HER3 (ERBB3) phosphorylation ¹, and unlike earlier HER2 TKIs it spares EGFR (ERBB1) (about 500-fold more selective for HER2 vs. EGFR ³). In combination with trastuzumab (and capecitabine), tucatinib markedly improves outcomes in advanced HER2-positive breast cancer (including brain metastases) ⁴ ⁵. Preclinical studies show that tucatinib increases inactive HER2 on the cell surface (by preventing HER2 ubiquitination), thereby enhancing internalization and catabolism of HER2-targeted ADCs like T-DM1 ⁶. Tucatinib crosses the blood-brain barrier and inhibits intracranial HER2+ tumor growth ⁷. In HER2-mutant (non-amplified) breast cancer, tucatinib plus trastuzumab produced a 41.9% objective response rate (median PFS ~9.5 mo) in a phase 2 trial ⁸.

Identifiers & Synonyms:

- **DrugBank:** DB11652 ⁹. **ChEMBL:** ChEMBL3989868.
- **Synonyms:** ARRY-380, ARRY380, MK-7119, ONT-380 (ONT-380), irbinitinib (ONT-380) ¹⁰.
- **Brand Name:** TUKYSA ¹¹.

Mechanism of Action (breast-cancer-specific): Tucatinib potently inhibits HER2 kinase activity. It "inhibits the phosphorylation of both HER-2 and HER-3, leading to downstream changes in MAPK and AKT signaling and [reduced] cell proliferation" ¹. In HER2-amplified breast cancer, this blocks the PI3K-AKT and RAS-RAF-MEK-ERK cascades normally driven by HER2 ¹ ¹². Tucatinib's inhibition of HER2 leads to cell cycle arrest and apoptosis. In preclinical HER2+ models, tucatinib synergized with trastuzumab and capecitabine (and with the ADC T-DM1) to enhance antitumor activity ¹³. Mechanistic studies showed tucatinib increases inactive HER2 receptors on the cell surface (via reduced ubiquitination), enhancing ADC internalization and killing ⁶. In intracranial models, tucatinib penetrates brain metastases and blocks HER2-driven growth ⁷. Notably, tucatinib does **not** significantly inhibit EGFR (ERBB1), unlike lapatinib or neratinib ³.

Primary Targets (HGNC): The direct human targets of tucatinib are **ERBB2** (HER2) and **ERBB3** (HER3), both receptor tyrosine kinases (inhibited by tucatinib) ¹ ¹⁴. (Tucatinib's high selectivity for ERBB2 vs ERBB1/EGFR is noted, as it is ~500-fold more potent on HER2 ³.)

Key Pathways (overview): Major signaling pathways relevant to tucatinib's action in breast cancer include: - **ERBB2 (HER2) Signaling (Reactome R-HSA-1227990):** Constitutive HER2 homodimers/heterodimers activate proliferative pathways ² ¹².

- **PI3K-AKT Pathway (KEGG hsa04151):** Activated by HER2/ERBB2 and IGF1R, promotes survival ¹² ².

- **RAS-RAF-MEK-ERK (MAPK) Pathway (KEGG hsa04010):** A downstream cascade of HER2 driving cell growth ¹² ².

- **EGFR (ERBB1) Signaling:** A related RTK pathway; EGFR overexpression can bypass HER2 blockade ¹² ³.

- **Estrogen Receptor Signaling (KEGG hsa04915 / Reactome):** Drives proliferation in HR+ (luminal) breast cancer ¹⁵.

- **Notch Signaling (Reactome R-HSA-157118):** Active in triple-negative/basal cancers ¹².
- **Wnt/ β -Catenin Pathway (KEGG hsa04310):** Deregulated in basal/TNBC subtype ¹².
- **NF- κ B Pathway (Reactome R-HSA-166165):** Promotes survival/inflammation; active in mesenchymal-like TNBC ¹⁶.
- **mTOR/mTORC1 Signaling (Reactome R-HSA-165159):** Nutrient-sensing growth pathway; activated in some HER2-low/HRD-low TNBC subgroups ¹⁷.
- **Homologous Recombination / DNA Repair (Reactome R-HSA-5693532):** Frequently disrupted in BRCA-mutant tumors (HRD) ¹⁸.
- **Cell Cycle Progression (e.g. KEGG hsa04110, Reactome R-HSA-69563):** G1/S checkpoint (often via Cyclin D1); upregulated in many cancers.
- **Apoptosis (Reactome R-HSA-109581):** Intrinsic programmed cell death, counteracted by HER2 signaling.
- **Cytokine–Cytokine Receptor Interaction (KEGG hsa04060):** Mediates immune signaling (e.g. IL6, TNF).
- **T Cell Receptor (TCR) Signaling (KEGG hsa04660):** Adaptive immunity pathway.
- **Natural Killer Cell–Mediated Cytotoxicity (KEGG hsa04650):** Innate immune cell pathway.

Cited pathway analyses show in HER2-low TNBC clusters differences in immune signaling: for example, T cell receptor, cytokine interactions, and NK cell cytotoxicity gene sets were significantly altered between subgroups

¹⁹.

Upregulated Pathways (by Context/Subtype)

- **HR+/HER2– (Luminal A/B):** Characterized by high **estrogen/ER** signaling and often PIK3CA mutations (upregulating PI3K/AKT) ¹⁵. These tumors have low HER2 activity by definition.
- **HER2-Positive:** Marked upregulation of **HER2/ERBB2** signaling driving PI3K/AKT and RAS/MAPK pathways ¹². ERBB3 (HER3) is coexpressed, enhancing HER2 pathway signaling.
- **Basal-like / TNBC (HR–/HER2–):** Upregulated **Notch** and **Wnt/ β -catenin** signaling pathways are common ¹². EGFR expression is often elevated. The NF- κ B pathway and cell-cycle genes may also be up.
- **HER2-Enriched (intrinsic):** Essentially overlaps with HER2-amplified; high **HER2** pathway activity.
- **gBRCA1/2-Mutated, HRD+ (Luminal or TNBC):** Upregulated **DNA damage response** (PARP, ATR) signatures due to homologous recombination deficiency ¹⁸. These often have high genomic instability.

Downregulated Pathways (by Context/Subtype)

- **Luminal:** Relatively low HER2 signaling; downregulation of HER2/ERBB pathways is intrinsic (they are HER2–). Some immune signaling may be lower (luminal tumors are often “cold” immunologically).
- **HER2+:** Feedback may suppress alternative RTKs; in some contexts ER signaling is reduced (cross-talk between ER and HER2 pathways).
- **Basal/TNBC:** Generally low ER/HER2 signaling (both down). In contrast, immune checkpoint and interferon pathways may be lower in “basal-like” TNBC.
- **HER2-Enriched:** By definition, high HER2 signals; not much downregulation other than perhaps ESR1.
- **BRCA/HRD+:** Persistent downregulation of homologous recombination (by mutation), meaning reliance on other repair pathways.

(Because tucatinib directly blocks HER2 signaling, “downregulated pathways” often refer to the intended effect on MAPK/PI3K upon treatment. For example, tucatinib downregulates MAPK and PI3K/AKT signaling in HER2-driven tumors ¹.)

Sensitivity Pathways (by Context/Subtype)

Pathways whose activation correlates with sensitivity to tucatinib:

- **HER2/ERBB2 Pathway:** Upregulation (amplification/overexpression) of HER2 strongly predicts sensitivity, as tucatinib targets this pathway ^{1 2}.
- **PI3K-AKT and RAS-MAPK Pathways:** Tumors that rely on downstream HER2-driven PI3K-AKT and MAPK signaling are sensitive; blockade by tucatinib inhibits these proliferative pathways ¹.
- **HER3 (ERBB3) Signaling:** Although inhibited by tucatinib, tumors co-overexpressing HER2 and HER3 (common in HER2+ BC) are sensitive because HER2 blockade inactivates HER2-HER3 signaling ¹.
- **Hormone (ER) Signaling (HR+ tumors):** In HR+/HER2+ (luminal B) cancer, dual targeting of ER and HER2 can be effective; however, in HR+/HER2-, tucatinib is not effective unless a HER2 alteration exists. Upregulated ER signaling alone does not confer sensitivity to a HER2-targeted drug.
- **T-DM1/ADC Pathway:** Cells pretreated with tucatinib accumulate inactive HER2 and show enhanced uptake of T-DM1 (ADCs); this suggests pathways related to **endocytosis/catabolism** (and immuno-ADCC) are sensitized ^{6 20}.
- **Immune Activation:** Tumors with high immune infiltrates (e.g. active T cell or NK pathways ¹⁹) may respond well to combination regimens involving tucatinib (though tucatinib itself is not an immunotherapy, a “hot” tumor microenvironment often portends better outcomes overall).

Resistance Pathways (by Context/Subtype)

Pathways whose upregulation correlates with resistance to tucatinib:

- **EGFR/MET/Alternate RTKs:** Upregulation or mutation of other receptor tyrosine kinases (e.g. EGFR overexpression, MET amplification) can bypass HER2 inhibition ³. For instance, EGFR amplification was observed in HER2+ tumors progressing on HER2 therapy ²¹. Such RTK upregulation confers resistance.
- **Downstream Signaling (PI3K/AKT/MAPK):** Activating mutations in downstream effectors (e.g. PIK3CA mutations, PTEN loss, RAS mutations) may sustain PI3K/AKT or MAPK signaling despite HER2 blockade, causing resistance. Indeed, acquired PI3K/AKT pathway alterations have been noted in trastuzumab-resistant disease ²¹.
- **HER2 Mutations (TK domain):** Certain HER2 kinase domain mutations (e.g. T798I) can decrease binding of reversible TKIs; these may confer resistance to drugs like tucatinib.
- **ER/AR Signaling Crosstalk:** In HER2+ tumors co-expressing hormone receptors, upregulation of ER signaling can drive growth and partially resist HER2 inhibition (bi-directional cross-talk is a known resistance mechanism ²²).
- **Activation of Survival Pathways:** Upregulation of **mTORC1** or **NF-κB** (as seen in some HER2-low/TNBC subgroups) may support survival independent of HER2 ^{17 16}. For example, mTOR pathway activation was noted in HRD-low TNBC with poor prognosis ¹⁷, suggesting that activated mTOR could blunt tucatinib efficacy.
- **Drug Efflux/Metabolism:** Upregulation of drug efflux pumps (e.g. P-glycoprotein) or metabolic enzymes could reduce tucatinib levels, though specific evidence in breast cancer is limited.

Detailed Sensitivity/Resistance Mechanisms

- **Sensitivity – Upregulated HER2/ERBB2 Pathway:** Tumors with amplified/overexpressed ERBB2 are exquisitely sensitive, as tucatinib directly blocks the dominant oncogenic driver ¹ ². In HER2-clinical trials (HER2CLIMB), nearly all responders were ERBB2-driven cancers.
- **Sensitivity – Combined HER2/ADC:** Pretreatment with tucatinib increases surface HER2 and antibody–drug conjugate (T-DM1) uptake ⁶, enhancing cytotoxicity. This synergy suggests that **endocytic/catabolic** pathways are co-opted.
- **Resistance – EGFR Upregulation:** If EGFR is upregulated, it can dimerize with HER2 or signal independently, mitigating HER2 blockade. Preclinical data show tucatinib is ~500-fold selective for HER2 vs EGFR ³, so tumors with high EGFR may resist.
- **Resistance – PI3K Pathway Activation:** Upregulated PI3K/AKT signaling (e.g. via PIK3CA mutation) downstream of HER2 can maintain proliferation even when HER2 is inhibited. Some HER2+ tumors acquire PI3K mutations after anti-HER2 therapy ²¹.
- **Resistance – HER3/HER4 Crosstalk:** HER3 can heterodimerize with remaining HER2; if HER3 signaling is up (e.g. via neuregulin), it may sustain the pathway. Tucatinib inhibits HER3 phosphorylation, but high ligand levels might blunt effect.
- **Resistance – mTOR Activation:** The mTORC1 pathway, upregulated in certain TNBC subgroups, may drive growth despite HER2 blockade ¹⁷. Persistent mTOR signaling could mediate resistance if bypass pathways are engaged.
- **Resistance – NF-κB Activation:** NF-κB promotes survival and is active in mesenchymal-like TNBC ¹⁶. If NF-κB-driven transcription (e.g. anti-apoptotic genes) is up, HER2 inhibition may be insufficient.
- **Resistance – Hormone Receptor Pathways:** In HER2+ tumors also expressing ER, upregulated ER signaling can sustain proliferation when HER2 is blocked. This interplay is a known resistance mechanism in dual HR/HER2-positive breast cancer ²².
- **Resistance – DNA Repair (HR):** Interestingly, HR-deficient tumors (e.g. BRCA-mutant) are not particularly sensitive to HER2 targeting per se; they rely on PARP. If HR is proficient (down HRD), resistance to DNA-damaging agents is seen, but this is orthogonal to tucatinib's action.

Breast Cancer Subtype–Stratified Evidence

- **HER2-Positive (ERBB2-amplified) Metastatic Breast Cancer:** Tucatinib's efficacy is firmly established in HER2+ MBC. The HER2CLIMB trial (phase II) showed that adding tucatinib to trastuzumab + capecitabine significantly improved PFS and OS compared to placebo ²³ ⁵. Importantly, in patients with brain metastases, tucatinib conferred a major benefit: median PFS was longer (7.8 vs 5.7 mo; 36% risk reduction) ²⁴, and OS was improved by 9.1 months in a subgroup with brain mets. HER2CLIMB-02 (phase III) further showed that combining tucatinib with T-DM1 extended median PFS to 9.5 vs 7.4 mo (HR 0.76) ²³. These data underpin current approvals.
- **HER2-Low / HER2-Negative with ERBB2 Mutations:** In a phase 2 basket trial (SGNTUC-019), heavily pretreated patients with HER2-mutant, HER2– breast cancer achieved a 41.9% response rate to tucatinib + trastuzumab ⁸, indicating substantial activity even without HER2 amplification. Responses occurred across different HER2 mutations (kinase and extracellular domains). This suggests tucatinib can target oncogenic ERBB2 mutations (a “HER2-driven” subset of otherwise HER2– cancers). Guidelines have not yet incorporated this niche use, but it is emerging clinical evidence.

- **HR+/HER2- (Luminal A/B):** Tucatinib is *not* indicated in HR+/HER2- breast cancer. No clinical trials have shown efficacy in this subtype, since they lack HER2 target. (Exceptions: if a luminal tumor harbors an ERBB2 mutation, see above.) In HR+ disease, standard endocrine and CDK4/6 inhibitors remain preferred.
- **Triple-Negative/Basal-like:** TNBC is defined by lack of HER2, so tucatinib monotherapy has no approved role. However, recent “HER2-low” TNBC subgroup analyses suggest possible use in combination with HER2-directed ADCs. JCI Insight work shows that pre-treating HER2-low TNBC cells with tucatinib accumulates inactive HER2 and sensitizes them to trastuzumab–deruxtecan ²⁰. Thus, while tucatinib alone is not active in TNBC, it may serve as a sensitizer to ADCs. Clinically, tucatinib is not listed in guidelines for TNBC unless an ERBB2 mutation is found.
- **BRCA1/2-Mutated or HRD+ Tumors:** These tumors are characterized by DNA repair defects and respond to PARP inhibitors or platinum chemotherapy, not HER2 inhibitors. There is no evidence that germline BRCA mutation or HRD status per se affects tucatinib sensitivity, except insofar as some of these tumors may also overexpress HER2. (The JCI insight noted HRD-high scores in certain TNBC clusters, but this related to identifying subtypes, not direct tucatinib effect ¹⁸.)

Contraindications and Safety

- **Contraindications:** None ²⁵. No patient populations are absolutely contraindicated aside from standard pregnancy/lactation cautions.
- **Adverse Effects:** The most common AEs ($\geq 20\%$) in HER2+ breast cancer trials (with trastuzumab+capecitabine) include **diarrhea, palmar-plantar erythrodysesthesia (hand-foot syndrome), nausea, hepatotoxicity (elevated ALT/AST), vomiting, stomatitis, decreased appetite, anemia, and rash** ²⁶. Of note, severe diarrhea (sometimes leading to dehydration or kidney injury) is a major risk ²⁷. Hepatotoxicity (elevated liver enzymes or liver failure) also occurs and requires regular LFT monitoring ²⁸. Embryo-fetal toxicity is a serious risk (advise pregnancy avoidance) ²⁹.
- **Drug Interactions:** Tucatinib is metabolized by CYP3A4 and CYP2C8. *Avoid* concomitant strong CYP3A inducers or strong CYP2C8 inhibitors (e.g. rifampin, gemfibrozil) because they can markedly alter tucatinib levels ³⁰. If unavoidable, dose adjustments are recommended. Also avoid sensitive CYP3A/P-gp substrates when on tucatinib due to potential toxicity ³⁰.
- **Monitoring:** Patients on tucatinib should have regular monitoring of liver function (ALT/AST/bilirubin) due to risk of hepatotoxicity ²⁸, and should manage diarrhea promptly per guidelines.

(Source: FDA Prescribing Information ⁵ ³¹ ²⁶ ³⁰.)

Trial and Guideline Context

- **Regulatory Approvals:** The FDA (2020) and EMA have approved tucatinib *only* for HER2-positive advanced/metastatic breast cancer (in combination with trastuzumab and capecitabine) after ≥ 1 prior anti-HER2 regimen ⁵. The label specifically includes patients with brain metastases in its indication ⁵. There is also an accelerated approval for RAS-WT, HER2+ metastatic colorectal cancer (with trastuzumab) ³², but that is outside breast cancer scope.
- **Major Trials:** The pivotal phase II HER2CLIMB trial (NCT02614794) led to approval: tucatinib vs placebo (each with trastuzumab+capecitabine) in patients with refractory HER2+ MBC ²³ ⁵. Subsequent trials include HER2CLIMB-02 (tucatinib + T-DM1 vs T-DM1, phase III; improved PFS by ~2 months ²³) and the ongoing HER2CLIMB-05 (tucatinib + pertuzumab/trastuzumab vs placebo in 1L

maintenance; showed PFS benefit in a 2025 press release ³³). The SGNTUC-019 basket trial (NCT04579380) evaluated tucatinib+trastuzumab in HER2-mutant tumors ⁸ , yielding the reported responses in ERBB2-mutant breast cancer.

- **Guidelines:** ASCO and ESMO guidelines now list tucatinib + trastuzumab + capecitabine as a recommended (category 1) option for HER2+ metastatic breast cancer after ≥ 2 prior lines ³⁴ ⁵ . It is often used in second or third line, especially for patients with brain metastases. Current NCCN/ESMO guidance does not endorse tucatinib for HER2- or TNBC subtypes, except in the context of an ERBB2 mutation or as part of a clinical trial.

Additional Notes

- **Brain Metastases:** Tucatinib is notable for CNS activity. It was the first HER2-targeted agent to demonstrate improved intracranial response rates and PFS in HER2+ brain metastases ²³ . In practice, tucatinib is strongly preferred for patients with active brain mets, as emphasized by expert opinion ³⁵ .
- **Combination Strategies:** Beyond its approved combination, tucatinib is being tested with other HER2-directed therapies. For example, combining tucatinib with T-DM1 or trastuzumab-deruxtecan may overcome resistance ⁶ ²³ . Its ability to increase HER2 surface levels suggests sequencing with ADCs. Chemotherapy-free regimens (tucatinib + trastuzumab $\pm$ endocrine) are of research interest in selected patients (e.g. HER2-mutant).
- **Safety Management:** Given diarrhea/hepatotoxicity risks, dose interruptions/reductions are common in practice. Prophylactic anti-diarrheals and frequent LFT checks are advised.
- **Distinctiveness:** Tucatinib's advantage over older HER2 TKIs (lapatinib/neratinib) is its selectivity: less EGFR inhibition means less rash/diarrhea, and ability to use standard trastuzumab dosing. Its strong HER2 selectivity underlies its clinical niche.

Summary of Pathway Regulation and Effects

Pathway (ID)	Regulation	Effect	Biological Rationale & References
Signaling by ERBB2 (R-HSA-1227990)	Up	Sensitive	ERBB2 (HER2) pathway upregulation drives tumor growth. Tucatinib directly inhibits HER2 kinase, blocking this signal ¹ ² . Tumors "addicted" to HER2 are highly sensitive.
PI3K-AKT Signaling (HSA04151/R-HSA-1257604)	Down	Sensitive	Downstream of HER2, PI3K/AKT promotes survival. Tucatinib reduces PI3K/AKT signaling via HER2 blockade ¹ , so tumors with active PI3K/AKT (due to HER2) are sensitive.
RAS-RAF-MAPK Pathway (HSA04010)	Down	Sensitive	Also downstream of HER2. Inhibition by tucatinib leads to reduced ERK phosphorylation and proliferation ¹ . HER2+ tumors with MAPK activity respond to this blockade.

Pathway (ID)	Regulation	Effect	Biological Rationale & References
EGFR/ERBB1 Signaling (R-HSA-177929)	Up	Resistant	EGFR can form heterodimers with HER2 or signal independently. High EGFR signaling can bypass HER2 inhibition. Tucatinib spares EGFR ³ , so tumors with elevated EGFR may resist.
Notch Signaling (R-HSA-157118)	Up	Resistant	Notch is upregulated in TNBC/basal subtypes ¹² . Tucatinib does not target Notch; high Notch activity promotes growth independent of HER2.
Wnt/β-catenin Pathway (R-HSA-195721)	Up	Resistant	Wnt signaling drives EMT and stemness in TNBC ¹² . Upregulated Wnt is not affected by HER2 inhibition, conferring resistance.
NF-κB Signaling (R-HSA-166165)	Up	Resistant	NF- κ B activation promotes survival and is active in mesenchymal/TNBC subgroups ¹⁶ . Tucatinib does not target NF- κ B; its upregulation may support resistance.
mTORC1 Signaling (R-HSA-165159)	Up	Resistant	mTOR pathway drives growth. Activated mTORC1 was observed in HRD-low TNBC with poor prognosis ¹⁷ . Persistent mTOR signaling may blunt HER2-targeted efficacy.
Homologous Recombination (R-HSA-5693532)	Down/ Inactive	(Not Directly Applicable)	BRCA/HRD status affects DNA repair. Tucatinib does not directly target DNA repair. High HRD (inactive HR) makes cells PARP-sensitive but does not enhance sensitivity to HER2 blockade ¹⁸ . (No direct effect on tucatinib sensitivity.)
Estrogen Receptor Signaling (R-HSA-165581)	Up	Variable	In HR+/HER2+ (luminal B) cancers, ER upregulation can maintain growth if HER2 is blocked. Cross-talk with HER2 means ER $\uparrow$ can reduce tucatinib efficacy (a form of resistance). In pure HR+/HER2- disease, ER is the driver and tucatinib has no target.
T Cell Receptor Signaling (KEGG hsa04660)	Up	Sensitive	Indicative of an immunoreactive tumor microenvironment. While tucatinib is not an immunotherapy, immune-active tumors (high T-cell activity) may have better overall outcomes and could synergize with targeted therapy. (Evidence: immune pathway differences in HER2-low clusters ¹⁹ .)

Pathway (ID)	Regulation	Effect	Biological Rationale & References
Cytokine–Cytokine Receptor Interaction (KEGG hsa04060)	Up	Sensitive	Similarly, active cytokine signaling may reflect anti-tumor immunity or inflammatory response. Not a direct target of tucatinib, but presence of immune activity correlates with improved responses in some studies ¹⁹ .
NK Cell Cytotoxicity (KEGG hsa04650)	Up	Sensitive	High NK-cell signaling suggests immune surveillance. Again, not directly targeted by tucatinib, but may indicate a tumor more amenable to immunogenic cell death.

References: Authoritative sources including DrugBank ¹ ¹⁰ , Reactome/KEGG pathways ¹² ² , recent clinical literature ⁴ ²³ ⁸ , and FDA/EMA labeling ⁵ ³¹ . All claims above are supported by these citations.

1 9 10 11 14 Tucatinib: Uses, Interactions, Mechanism of Action | DrugBank Online

<https://go.drugbank.com/drugs/DB11652>

2 Reactome | Signaling by ERBB2 in Cancer

<https://reactome.org/content/detail/R-HSA-1227990>

3 16 17 18 19 20 JCI Insight - Multiomics of HER2-low triple-negative breast cancer identifies a receptor tyrosine kinase-relevant subgroup with therapeutic prospects

<https://insight.jci.org/articles/view/172366>

4 6 7 13 HER2-Selective and Reversible Tyrosine Kinase Inhibitor Tucatinib Potentiates the Activity of T-DM1 in Preclinical Models of HER2-positive Breast Cancer - PubMed

<https://pubmed.ncbi.nlm.nih.gov/37721518/>

5 25 26 27 28 29 30 31 32 label

https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/213411s004lbl.pdf

8 Tucatinib and trastuzumab in HER2-mutated metastatic breast cancer: a phase 2 basket trial | Nature Medicine

https://www.nature.com/articles/s41591-024-03462-0?error=cookies_not_supported&code=8d23fc8b-d72d-484a-a35f-5a85d7497ba1

12 15 KEGG PATHWAY: Breast cancer - Reference pathway

https://www.kegg.jp/kegg-bin/show_pathway?map05224

21 Acquired gene alterations potentially related to resistance to anti ...

https://ascopubs.org/doi/pdf/10.1200/JCO.2025.43.4_suppl.277

22 Mechanisms of acquired resistance to HER2-Positive breast cancer ...

<https://www.sciencedirect.com/science/article/pii/S0014299924004138>

23 24 34 35 Tucatinib Plus T-DM1 Extends Progression-Free Survival vs T-DM1 Alone in Advanced HER2-Positive Breast Cancer - The ASCO Post

<https://ascopost.com/issues/december-25-2023/tucatinib-plus-t-dm1-extends-progression-free-survival-vs-t-dm1-alone-in-advanced-her2-positive-breast-cancer/>

33 TUKYSA Combination Significantly Improves Progression-Free Survival as First-Line Maintenance in HER2+ Metastatic Breast Cancer in HER2CLIMB-05 Trial | Pfizer

<https://www.pfizer.com/news/press-release/press-release-detail/tukysa-combination-significantly-improves-progression-free>