

## Letrozole – Drug Summary

Letrozole is a third-generation non-steroidal aromatase inhibitor (AI) used in estrogen receptor-positive (ER<sup>+</sup>) breast cancer, especially in postmenopausal women <sup>1</sup> <sup>2</sup>. It is indicated for all stages of hormone-receptor-positive breast cancer in postmenopausal women <sup>1</sup> (often in luminal A/B subtypes) and is often combined with targeted agents (e.g. CDK4/6 inhibitors) in HR<sup>+</sup>/HER2<sup>-</sup> metastatic disease <sup>3</sup> <sup>4</sup>. By blocking estrogen synthesis, letrozole causes regression of ER<sup>+</sup> tumors <sup>2</sup>. It is *not* effective in ER-negative (e.g. triple-negative) tumors, and its use in HER2<sup>+</sup> disease is only when the tumor is also HR<sup>+</sup> (often Luminal B); HER2<sup>+</sup>/HR<sup>-</sup> tumors do not benefit from aromatase inhibition.

## Identifiers & Synonyms

- **ChEMBL ID:** ChEMBL1444 <sup>5</sup>
- **DrugBank ID:** DB01006 <sup>6</sup>
- **Synonyms:** Chemical synonyms include *Letrozol* <sup>7</sup> (the active moiety). Brand names include **Femara** <sup>8</sup> (Novartis) and generic letrozole. Other synonyms (e.g. CGS-20267) are documented <sup>9</sup>.

## Mechanism of Action

Letrozole competitively inhibits the aromatase enzyme (CYP19A1), binding its active site and blocking the conversion of androgens (testosterone/androstenedione) into estrogens <sup>2</sup>. In postmenopausal women – where peripheral fat and adrenal glands are the main estrogen sources – this causes profound estrogen depletion. The resulting drop in estradiol levels deprives ER $\alpha$ -expressing tumor cells of growth signals, leading to tumor regression (through cell-cycle arrest and apoptosis) in ER<sup>+</sup> breast cancers <sup>2</sup> <sup>10</sup>. As a type-II (triazole) inhibitor, letrozole has a long half-life (~2 days) and is highly selective for CYP19A1, with minimal effect on other steroidogenic enzymes (e.g. cortisol or aldosterone synthesis) <sup>2</sup> <sup>11</sup>. Inhibition of aromatase also causes feedback increases in gonadotropins (LH/FSH), but postmenopausally these have little effect. In summary, letrozole's high-level mechanism is estrogen deprivation: it **decreases** estrogen signaling in ER<sup>+</sup> tumors <sup>2</sup>.

## Primary Targets

- **CYP19A1 (Aromatase):** The principal pharmacologic target is the aromatase enzyme, encoded by gene CYP19A1 (HGNC symbol *CYP19A1*) <sup>2</sup>. Letrozole binds to CYP19A1's heme active site and halts the final step of estrogen biosynthesis <sup>2</sup>. No other direct human molecular targets (HGNC symbols) are known for letrozole.

# Pathways (Overview)

Letrozole affects multiple cancer-relevant pathways by altering hormone-driven signaling. Key pathways include:

- **Estrogen Response (MSigDB/HALLMARK\_ESTROGEN\_RESPONSE\_EARLY/LATE; Reactome “Estrogen-dependent gene expression”):** *Downregulated* by letrozole, since estrogen levels fall. Suppression of this pathway (including ERα target genes) is central to therapy <sup>2</sup> <sup>10</sup>. *Effect:* Downregulation correlates with tumor shrinkage (sensitivity).
- **Steroid Hormone Biosynthesis (KEGG hsa00140):** The enzymatic pathway inhibited by letrozole. Letrozole blocks CYP19A1 in this pathway, *downregulating* estrogen and estrone production <sup>2</sup>.
- **Cell Cycle (Reactome “Mitotic G1-G1/S phases” or KEGG Cell Cycle hsa04110):** ER signaling promotes cell-cycle progression. Letrozole-induced estrogen deprivation causes *downregulation* of proliferation genes (cyclins/CDKs). In endocrine-sensitive tumors, G1/S and mitotic pathways are suppressed, leading to growth arrest <sup>2</sup> <sup>12</sup>.
- **Apoptosis (Reactome “Intrinsic apoptotic signaling”; HALLMARK\_APOPTOSIS):** Loss of ER signaling triggers apoptotic pathways in HR<sup>+</sup> cancer cells. Letrozole tends to *upregulate* pro-apoptotic signals (e.g. increased caspases) <sup>13</sup> <sup>2</sup>, correlating with tumor cell death (sensitivity).
- **PI3K/AKT/mTOR (KEGG hsa04151; Reactome PI3K cascade):** ER<sup>+</sup> cells often activate PI3K/AKT as a growth/survival pathway. In many letrozole-resistant cancers, this pathway becomes *upregulated* (compensatory) <sup>14</sup>. Hyperactivation of PI3K/AKT/mTOR (e.g. via AKT1/2, FGFR3) is associated with resistance <sup>14</sup> <sup>12</sup>.
- **MAPK/ERK (KEGG hsa04010; Reactome “RAF/MAPK signaling”):** Similarly, Ras/MAPK signaling can be activated in resistance, leading to continued proliferation despite low estrogen. Letrozole often *downregulates* MAPK activity in sensitive tumors, but resistant clones may *upregulate* it to bypass ER dependency (not directly cited but consistent with PI3K crosstalk).
- **FOXM1 Transcriptional Network (Reactome “FOXM1 transcription factor network”):** FOXM1 drives G2/M progression. Pathway analysis of letrozole-resistant ER<sup>+</sup> tumors shows the FOXM1 network *activated* <sup>15</sup>, promoting cell-cycle entry in non-responders.
- **PLK1 Signaling Events (Reactome “PLK1 signaling events”):** Polo-like kinase 1 controls G2/M transition. PLK1 pathway is *upregulated* in letrozole-resistant cells <sup>16</sup> <sup>15</sup>, indicating cell-cycle drive contributes to resistance.
- **Syndecan-1 (SDC1) Pathway:** Syndecan-1 (SDC1) is involved in cell-matrix adhesion and growth-factor signaling. The SDC1 signaling pathway is *activated* in resistant ER<sup>+</sup> tumors <sup>17</sup>, often inversely correlated with ER expression, and may promote invasion.
- **Matrix Metalloproteinases (MMPs) / Extracellular Matrix (Reactome “Extracellular matrix organization”):** MMP2 and MMP9 are *upregulated* in resistant or aggressive ER<sup>+</sup> cancers <sup>18</sup>, facilitating invasion and metastasis. They are known markers of endocrine resistance.
- **Angiogenesis (Reactome “VEGF signaling”; HALLMARK\_ANGIOGENESIS):** Letrozole indirectly reduces angiogenic signaling by lowering tumor growth, but VEGF-A can still be produced. Pathway analysis identifies VEGFA as a top target in resistant settings <sup>19</sup>. Inhibition of estrogen may mildly upregulate VEGF in stroma, but overall the effect of letrozole is to slow angiogenesis (as tumors regress).
- **Hypoxia / HIF-1α (Reactome “HIF-1α transcription factor network”):** In resistant tumors, HIF-1α network appears *suppressed* <sup>20</sup>, possibly because oxygen demand is altered. HIF-1α can drive aggressive behavior; its inhibition was noted in non-responders.

- **Immune Response (ImmuneDB/GO “immune response”; cytokine signaling):** Letrozole modifies the tumor microenvironment. Neoadjuvant studies show that letrozole increases tumor-infiltrating lymphocytes (TILs) <sup>21</sup>, indicating an *upregulation* of immune activity. Paradoxically, patients with higher TILs during AI therapy had **poorer** response <sup>21</sup>, perhaps reflecting a pro-inflammatory milieu associated with resistance. Immune-related pathways (e.g. interferon response, cytokine signaling) may thus be modulated by endocrine therapy.

Overall, letrozole’s action **downregulates** estrogen-driven proliferation and survival pathways (ER signaling, cell cycle) while **upregulating** apoptosis and, in resistant disease, alternative growth (PI3K/AKT, PLK1) and invasion (MMP/EMT) pathways. These pathway changes drive *sensitivity* or *resistance* to therapy: for example, suppression of ER and cell-cycle pathways correlates with good response, whereas upregulation of PI3K/AKT or PLK1 pathways marks resistance <sup>2</sup> <sup>16</sup>.

## Upregulated Pathways (with Letrozole)

- **Apoptotic signaling:** Letrozole induces pro-apoptotic genes (e.g. caspases). Preclinical studies show increased apoptosis markers with AIs <sup>13</sup>. This *upregulation* of cell death is a mechanism of sensitivity.
- **Adaptive Immune Pathways:** Increased TILs and immune gene expression are seen during letrozole therapy <sup>21</sup>. (Hallmark sets: TNFα signaling, IFNγ response may become active). Enhanced immune signaling may help anti-tumor activity, though its net effect on response is complex <sup>21</sup>.
- **Stress Response:** Oxidative stress and autophagy markers (e.g. PRDX4) can rise in resistant cells <sup>22</sup>. These pathways help cells cope with estrogen starvation; their activation is often seen in resistance.
- **Matrix Remodeling:** Genes like MMP9/MMP2 are *upregulated* in resistant ER<sup>+</sup> cancers <sup>18</sup>, reflecting epithelial–mesenchymal transition (EMT) and invasion. This also correlates with poorer endocrine response.
- **Angiogenic factors:** VEGFA and related genes can be modestly elevated as tumors adapt; pathway analysis highlights VEGFA in the resistant signature <sup>23</sup>.

## Downregulated Pathways (with Letrozole)

- **Estrogen/ER Signaling:** Pathways driven by ESR1 (ERα) are directly suppressed <sup>2</sup>. Estrogen-responsive gene sets (MSigDB HALLMARK\_ESTROGEN\_RESPONSE) are strongly downregulated in treated tumors.
- **Cell Proliferation (E2F, G1/S):** Letrozole downregulates E2F targets and cyclins (e.g. CCND1, CDK1) <sup>2</sup> <sup>12</sup>. (Hallmark\_E2F\_Targets, KEGG\_Cell\_Cycle). This reduction in proliferation correlates with favorable response.
- **Growth Factor Signaling (EGF/IGF):** Some growth factor pathways (EGFR/MAPK, IGF-1R) may be indirectly downregulated as ER cross-talk is blocked. Studies of combined letrozole+trastuzumab suggest reduced HER signaling when ER is shut off.
- **Steroid Biosynthesis:** Enzymatic steps upstream of aromatase may feedback down (via reduced substrate flow), though main effect is the block at CYP19A1.
- **Cholesterol Metabolism:** Letrozole-treated women often show *upregulated* cholesterol (see Safety), implying downregulation of pathways that consume cholesterol in estrogen synthesis <sup>24</sup>.

## Sensitivity vs Resistance Pathways

- **Sensitivity-Associated (Downregulated):** Effective letrozole response is marked by sustained suppression of ER-regulated genes and cell-cycle drivers. For instance, tumors that remain Ki67-low (indicating cell-cycle arrest) during therapy show better outcomes. Gene profiles of responders show lower PLK1/FOXM1 and intact apoptotic activation <sup>2</sup> <sup>21</sup>.
- **Resistance-Associated (Upregulated):** Letrozole resistance arises when alternate pathways compensate. Key upregulated pathways in resistant (non-responder) cases include PLK1/G2-M signaling <sup>16</sup>, FOXM1, PI3K/AKT (via targets like AKT1/2) <sup>14</sup>, and growth-factor/EMT pathways (Syndecan-1, MMPs) <sup>17</sup> <sup>18</sup>. The identification of PLK1 and FOXM1 activation in non-responders <sup>15</sup> illustrates this shift. Resistance is also linked to reactivation of ER signaling via ESR1 mutations, which keep ER pathways on despite letrozole <sup>25</sup> (ESR1 gain-of-function mutations mediate endocrine escape).

## Subtype-Stratified Evidence

- **HR<sup>+</sup>/HER2<sup>-</sup> (Luminal A/B):** The primary indication group. Letrozole as monotherapy (or with CDK4/6 inhibitors) is standard first-line in postmenopausal HR<sup>+</sup>/HER2<sup>-</sup> advanced or adjuvant settings <sup>3</sup> <sup>26</sup>. Luminal A tumors (high ER/PR, low Ki67) typically respond well to AIs, while luminal B (often higher grade or HER2+) may have a higher proliferation setpoint and sometimes need combined targeted therapy (e.g. CDK4/6). Trials show adding palbociclib/letrozole improves outcomes in luminal B-like subtypes.
- **HR<sup>+</sup>/HER2<sup>+</sup>:** Endocrine therapy (including letrozole) is used only if ER<sup>+</sup>; these “double-positive” tumors benefit from combined HER2-targeted and endocrine therapy. Some studies suggest similar disease-free survival whether tamoxifen or an AI is used in HER2<sup>+</sup>/HR<sup>+</sup> adjuvant therapy <sup>27</sup>. Letrozole can be combined with trastuzumab or dual blockade; e.g. a phase II trial combined trastuzumab+letrozole in ER<sup>+</sup>/HER2<sup>+</sup> metastatic BC (showing efficacy). However, HER2-driven signaling may reduce AI sensitivity, and estrogen-independent growth can occur <sup>28</sup>.
- **Triple-Negative (HR<sup>-</sup>/HER2<sup>-</sup>):** Letrozole has no target (no ER), so it is *not used*. TNBC is inherently resistant to aromatase inhibition.
- **gBRCA1/2 Mutation Carriers:** Most BRCA1-related BC are triple-negative, but BRCA2 tumors are often HR<sup>+</sup>. In BRCA-mutant HR<sup>+</sup> cases, endocrine treatment follows standard guidelines for HR<sup>+</sup> disease. There is no specific sensitivity of BRCA-mutant tumors to letrozole beyond their hormone status, and no unique biomarker-driven indication. (PARP inhibitors are used for BRCA-mutants, not AIs.)

## Contraindications and Safety

- **Contraindications:** Letrozole is contraindicated in **pregnancy** (FDA category X) and in women who may become pregnant <sup>29</sup>. It causes fetal harm and must not be used in gestation <sup>29</sup>. Its benefit in premenopausal women alone is not established, so it is *not given* to women with functioning ovaries unless combined with ovarian suppression <sup>29</sup>.
- **Bone Loss:** Aromatase inhibition markedly lowers estrogen, often causing **osteopenia/osteoporosis**. In trials, letrozole caused greater BMD loss than tamoxifen and increased fracture risk <sup>30</sup>. Patients on letrozole should have bone density monitored, and may need calcium/vitamin D or bisphosphonates to mitigate loss <sup>30</sup>.

- **Cardiovascular/Thrombotic:** Letrozole can raise cholesterol ( $\geq 52\%$  had hypercholesterolemia vs 28% on tamoxifen <sup>24</sup>) and increase cardiovascular risk. In NSABP B-42, prolonged letrozole raised thrombotic events modestly ( $HR \approx 1.85$ ) <sup>31</sup>. Conversely, it has a lower risk of uterine cancer and thromboembolism than tamoxifen. Still, patients' lipid profiles and cardiovascular status should be monitored.
- **Other Adverse Effects:** Common side effects include hot flashes, arthralgias/myalgias, fatigue, and joint stiffness <sup>32</sup>. Letrozole may cause **QT prolongation** (monitor with caution) and rare pulmonary issues (ILD/pneumonitis) <sup>33</sup>. It can cause elevated liver enzymes and mild neutropenia <sup>33</sup>. Fatigue and dizziness are noted; patients should be cautioned about operating machinery <sup>34</sup>. Ovarian failure (menopause) can occur if underlying ovarian reserve is suppressed, so it is avoided or given with ovarian suppression in younger women.
- **Drug Interactions:** Letrozole is metabolized by CYP2A6 and CYP3A4 <sup>35</sup>, so potent inhibitors/inducers of these enzymes can alter its levels. No major food effects are noted.

## Clinical Trial & Guideline Context

Current guidelines (NCCN, ESMO, ASCO) recommend 5–10 years of adjuvant endocrine therapy for all HR<sup>+</sup> breast cancers, with an aromatase inhibitor as first-line in postmenopausal women <sup>26</sup> <sup>36</sup>. Recent trials reinforce letrozole's role: for example, the NRG/NSABP B-42 trial (postmenopausal HR<sup>+</sup> patients disease-free after 5 years of ET) found that 5 additional years of letrozole significantly improved 10-year disease-free survival ( $HR \sim 0.84$ ) compared to placebo <sup>37</sup>. Extended therapy modestly reduces recurrence risk but must balance risks (bone loss, thrombosis) <sup>37</sup> <sup>31</sup>. The MA.17 trial similarly demonstrated that letting women switch to letrozole after 5 years of tamoxifen improved DFS <sup>38</sup>. In metastatic disease, letrozole (with or without CDK4/6 inhibitors) is standard first-line therapy for ER<sup>+</sup> disease. PALOMA-2 and other trials showed adding palbociclib to letrozole markedly improves progression-free survival (e.g. from  $\sim 14.5$  to 24.8 months) <sup>39</sup>. Guidelines also address sequencing: letrozole alone or with targeted agents is recommended initially; upon progression, other endocrine agents (fulvestrant, SERDs, or estrogens pathway antagonists) or chemotherapy may follow. In the last 5 years, trials have focused on combinations (e.g. CDK4/6 or PI3K inhibitors with letrozole) and biomarkers (e.g. multigene assays to select candidates for extended letrozole) <sup>40</sup> <sup>41</sup>.

## Summary Table of Pathway Effects

| Pathway (ID)                                   | Letrozole Effect (↑ / ↓) | Sensitivity vs. Resistance                | Rationale / Source                                                       |
|------------------------------------------------|--------------------------|-------------------------------------------|--------------------------------------------------------------------------|
| <b>Estrogen Response</b><br>(MSigDB, GO)       | ↓ (downregulated)        | ↓ Down = ↑ sensitivity (tumor regression) | Letrozole blocks estrogen synthesis <sup>2</sup> .                       |
| <b>Steroid Biosynthesis</b><br>(KEGG hsa00140) | ↓                        | — (target pathway)                        | Aromatase (CYP19A1) inhibition blocks estrogen production <sup>2</sup> . |

| Pathway (ID)                             | Letrozole Effect ( ↑ / ↓ )         | Sensitivity vs. Resistance                | Rationale / Source                                                                                                       |
|------------------------------------------|------------------------------------|-------------------------------------------|--------------------------------------------------------------------------------------------------------------------------|
| <b>Cell Cycle (Reactome, KEGG)</b>       | ↓                                  | ↓ Down = ↑ sensitivity; ↑ Up = resistance | ER-driven proliferation suppressed by letrozole <sup>2</sup> ; PLK1/FOXM1 up in resistance <sup>16</sup> <sup>15</sup> . |
| <b>PI3K/AKT/mTOR (KEGG hsa04151)</b>     | ↑ (in resistance)                  | ↑ Up = ↑ resistance                       | Compensatory survival signaling; AKT1/2 activation in non-responders <sup>14</sup> <sup>12</sup> .                       |
| <b>MAPK/ERK (KEGG hsa04010)</b>          | ↓ (initially)                      | ↑ Up (resistance) – thought               | Downstream of ER; letrozole suppresses MAPK, but alternative activation contributes to resistance.                       |
| <b>Apoptosis (Reactome)</b>              | ↑                                  | ↑ Up = ↑ sensitivity                      | Letrozole induces apoptosis in ER <sup>+</sup> cells <sup>13</sup> <sup>2</sup> .                                        |
| <b>FOXM1 (Reactome)</b>                  | ↑ (in resistance)                  | ↑ Up = ↑ resistance                       | FOXM1-driven G2/M network activated in non-responders <sup>15</sup> .                                                    |
| <b>PLK1 (Reactome R-HSA-157018)</b>      | ↑ (in resistance)                  | ↑ Up = ↑ resistance                       | PLK1/G2–M signaling up in letrozole-resistant cells <sup>16</sup> .                                                      |
| <b>Syndecan-1 (Reactome)</b>             | ↑ (in resistance)                  | ↑ Up = ↑ resistance                       | SDC1 signaling activated in non-responders <sup>17</sup> (promotes proliferation/EMT).                                   |
| <b>MMP/ECM Remodelling (GO/Reactome)</b> | ↑ (in resistance)                  | ↑ Up = ↑ resistance                       | MMP2/9 upregulated in aggressive/resistant ER <sup>+</sup> tumors <sup>18</sup> .                                        |
| <b>Angiogenesis (Reactome, HALLMARK)</b> | ↓ (due to overall tumor shrinkage) | ↓ Down = ↑ sensitivity                    | Tumor regression reduces angiogenic drive; VEGFA target in resistant context <sup>23</sup> .                             |
| <b>Hypoxia (HIF-1α network)</b>          | ↓ (in resistance)                  | ↓ Down = resistance                       | HIF-1α TF network was inhibited among non-responders <sup>20</sup> .                                                     |
| <b>Immune (ImmuneDB/GO)</b>              | ↑ (TIL infiltration)               | ↑ Up = ↑ resistance (paradoxically)       | Letrozole increases TILs <sup>21</sup> ; higher TILs signaled poorer response.                                           |

**Sources:** Authoritative databases and studies (DrugBank <sup>2</sup> ; MyCancerGenome <sup>42</sup> ; clinical guidelines and literature <sup>26</sup> <sup>21</sup> <sup>16</sup> <sup>4</sup> ) provide evidence for these pathways and their roles in letrozole response and resistance. (See citations in text.)

1 10 **Letrozole (Femara): What to Expect, Side Effects, and More**

<https://www.breastcancer.org/treatment/hormonal-therapy/femara>

2 3 5 6 7 8 9 11 33 35 **Letrozole: Uses, Interactions, Mechanism of Action | DrugBank Online**

<https://go.drugbank.com/drugs/DB01006>

4 12 14 15 16 17 18 19 20 22 23 25 **Secondary Analysis of Human Bulk RNA-Seq Dataset Suggests Potential Mechanisms for Letrozole Resistance in Estrogen-Positive (ER+) Breast Cancer**

<https://www.mdpi.com/1467-3045/46/7/424>

13 **Differential biological effects of aromatase inhibitors: Apoptosis ...**

<https://www.sciencedirect.com/science/article/abs/pii/S0303720721002707>

21 **Tumour-infiltrating lymphocytes and response to neoadjuvant letrozole in patients with early oestrogen receptor-positive breast cancer: analysis from a nationwide phase II DBCG trial | Breast Cancer Research | Full Text**

<https://breast-cancer-research.biomedcentral.com/articles/10.1186/s13058-020-01285-8>

24 29 30 34 **label**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2014/020726s027lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2014/020726s027lbl.pdf)

26 32 **Hormone Therapy for Breast Cancer | Breast Cancer Treatment | American Cancer Society**

<https://www.cancer.org/cancer/types/breast-cancer/treatment/hormone-therapy-for-breast-cancer.html>

27 **Adjuvant endocrine therapy in HER2-positive breast cancer patients**

<https://www.sciencedirect.com/science/article/pii/S2059702921000454>

28 **Current Trends in the Treatment of HR+/HER2+ Breast Cancer**

<https://www.tandfonline.com/doi/full/10.2217/fon-2020-0504>

31 36 **Extended adjuvant endocrine therapy in early breast cancer**

[https://www.esmopen.com/article/S2059-7029\(25\)00926-3/fulltext](https://www.esmopen.com/article/S2059-7029(25)00926-3/fulltext)

37 **10-year results of NRG Oncology/NSABP B-42 trial | Science Codex**

<https://sciencecodex.com/10-year-results-nrg-oncologynsabp-b-42-trial-636794>

38 **Preventing relapse beyond 5 years: the MA.17 extended adjuvant trial**

<https://pubmed.ncbi.nlm.nih.gov/16730271/>

39 **Pfizer Announces Overall Survival Results from Phase 3 PALOMA-2 ...**

<https://www.pfizer.com/news/press-release/press-release-detail/pfizer-announces-overall-survival-results-phase-3-paloma-2>

40 **SABCS 2019: Extended letrozole therapy improves disease-free ...**

<https://ecancer.org/en/news/17068-sabcs-2019-extended-letrozole-therapy-improves-disease-free-survival-in-postmenopausal-women-with-hr-positive-breast-cancer>

41 **10-year results of NRG Oncology/NSABP B-42 trial**

<https://www.nrgoncology.org/News/News-Post-Template/extended-letrozole-therapy-improves-disease-free-survival-dfs-in-postmenopausal-women-with-hr-breast-cancer-10-year-results-of-nrg-oncologynsabp-b-42-trial>

42 **letrozole - My Cancer Genome**

<https://www.mycancergenome.org/content/drugs/letrozole/>