

Readout errors and estimator statistics (M=2 bell_tilted for a-b; M=1 tilted, eps2=0.005 for c-d)
(a) Signed visibility vs readout error
$t(q)/t(0)$
Readout error q
Measured accumulator, N=4
Coherent streaming VD, N=4
Measured accumulator, N=8
Coherent streaming VD, N=8
(b) Measured-tree / coherent variance ratio
$v_{\mathrm{tree}}/v_{\mathrm{coh}}$
simulated, N=4
$(1-2q)^{-2(N-2)}$, N=4
simulated, N=8
$(1-2q)^{-2(N-2)}$, N=8
Readout error q
(c) Monte Carlo sd / delta-method sd (95% CI), N=4
$\mathrm{sd}(\hat{\mu})/\sqrt{v/S}$
Coherent streaming VD, q=0
Coherent streaming VD, q=0.05
Coherent streaming VD, q=0.1
Measured accumulator, q=0
Measured accumulator, q=0.05
Measured accumulator, q=0.1
Shots S
(d) Ratio bias x S at q=0.05, N=4 (95% CI)
$S(\bar{\hat{\mu}}-\mu_A)$
Measured accumulator: c_A
Coherent streaming VD: c_A
Measured accumulator: MC
Coherent streaming VD: MC
Shots S