THE AXIOMATIC SECTOR COSMOLOGICAL MODEL (ASCM) — PART II: TOPOLOGICAL

A Boundary-Coupled Dual-Manifold Effective Field Theory on an $S^1/\mathbb{Z}_2$ Warped Orbifold with Dynamical Spin-Connection Localization, Combinatorial UV State-Sum Capacity, Free-γ 4-Simplex Vertex Mixing, Unconstrained Renormalization-Group Flow, and $SL(2,\mathbb{C})_q$ Casimir Phase Isolation

J. B. Randolph (ORCID: 0009-0005-7943-2176)
Overland Park, Kansas 66212, USA | *Published & Archived (Oct 2026)* | *DOI: 10.6084/m9.figshare.34734825*

Declaration of Authorship: The author declares sole authorship of all theoretical derivations, analytical proofs, topological extensions, and numerical verification architectures presented in this manuscript and its companion supplement.

$$S_{\text{ASCM-II}} = S_M[g,\Phi] + S_{\tilde{M}}[\tilde{g},\tilde{\Phi}] + S_{B/\mathbb{Z}_2}[G,\Phi_B,\Psi_i] + S_\partial[h,\xi,\sigma_\partial,\gamma] + S_{\text{seq}}[\lambda,\tilde{\lambda},\Gamma_{\text{UV}}]$$

ABSTRACT

We formulate and analyze Part II (ASCM Topological) of the Axiomatic Sector Cosmological Model (ASCM), a covariant effective field theory built on a boundary-coupled dual-manifold geometry $M^4 \times_{\xi(\mu)} \tilde{M}^4$ embedded within a five-dimensional warped orbifold bulk $B^5 = M^4 \times S^1/\mathbb{Z}_2$. Because cross-sector operators couple through the shared codimension-one interface ∂M and a global topological sequestering functional S_{seq}, diffeomorphism invariance, Bianchi identities, orbifold parity, and unitarity impose rigid consistency conditions linking vacuum energy, primordial perturbations, chiral anomaly transport, and late-time expansion. ASCM Topological replaces all residual phenomenological parameter assignments, algebraic overlap inversions, and infrared horizon circularities from Part I with six exact topological and representation-theoretic derivations.

First, to mathematically prohibit five-dimensional right-handed Majorana fields $\Psi_i(x,z)$ from generating a μeV–meV Kaluza–Klein sterile neutrino tower that would catastrophically violate Big Bang Nucleosynthesis (BBN) ΔN_{eff} constraints, the 5D warped bulk B^5 is formulated on an $S^1/\mathbb{Z}_2$ orbifold with discrete parity transformation $\Psi_i(x,-z) = \pm\gamma_5\Psi_i(x,z)$. Chiral boundary bifurcation locks the right-chiral singlet to even Neumann boundary conditions ($\partial_z f_{iR}(0) = 0$), permitting a boundary-localized zero mode, while forcing the left-chiral sterile component into odd Dirichlet boundary conditions ($f_{iL}(0) = f_{iL}(L) = 0$). This forbids a massless sterile zero mode identically and forces the lightest sterile excitation to the ultraviolet cutoff scale ($m_{1,L} \sim k_{\text{bulk}} e^{k_{\text{bulk}}L} \sim 10^{14}\ \text{GeV}$), decoupling the Kaluza–Klein sterile tower from the thermal bath ($\Delta N_{\text{eff}}^{\text{KK}} = 0$).

Second, we generate the right-handed neutrino mass hierarchy $M_1 \ll M_2 \ll M_3$ exclusively from 5D background curvature without algebraic inversion. By coupling the 5D bulk fermions to a dynamical 5D bulk scalar field $\Phi_B(z)$ via the 5D spin connection ω_M^{AB} and its curvature invariant $\mathcal{I}_\omega = G^{MN}\omega_M^{AB}\omega_{N,AB} =$

$-8k_{\text{bulk}}^2$, the scalar vacuum expectation value $\langle \Phi_B(z) \rangle = v_B k_{\text{bulk}}^{3/2}$ organically generates quantized localization eigenvalues $(c_1, c_2, c_3) = (0.647460, 0.361495, 0.000000)$ via a closed-form Lambert-W topological triality relation, yielding $M_R = \text{diag}(10^{14}, 10^{15}, 10^{16})$ GeV.

Third, we establish topological phase isolation via the native $SL(2,\mathbb{C})_q$ Casimir invariants $\mathcal{C}_1 = \vec{L}^2 - \vec{K}^2$ and $\mathcal{C}_2 = \vec{L} \cdot \vec{K}$. Under the EPRL linear simplicity constraint $\vec{K} = \gamma \vec{L}$ ($\mathcal{C}_1 = (1-\gamma^2)\vec{L}^2$, $\mathcal{C}_2 = \gamma \vec{L}^2$), the complex saddle action $\mathcal{S}_{\mathbb{C}}(j_{\text{crit}}) = \text{Re}(\mathcal{S}_{\mathbb{C}}) + i\,\text{Im}(\mathcal{S}_{\mathbb{C}})$ at $j_{\text{crit}} = j_0 + 2i\gamma\ell_{Pl}^2 H_* \sigma_j^2$ splits into orthogonal parity sectors: $\text{Re}(\mathcal{S}_{\mathbb{C}})$ couples exclusively to parity-even $\mathcal{C}_1$, ensuring the 42×42 Dirac matrix Δ_{AB} remains strictly real and Hermitian (insulating the ghost-free $88 \to 10$ mode reduction), while $\text{Im}(\mathcal{S}_{\mathbb{C}})$ couples exclusively to parity-odd $\mathcal{C}_2$, projecting onto the Nieh–Yan invariant and Pontryagin density $R\tilde{R}$ as a bulk total derivative.

Fourth, we treat the Barbero–Immirzi parameter γ as a free variable and derive the vertex-mixing factor $\Delta_{\text{mix}}(\gamma) = \frac{4}{\pi}\sqrt{\gamma}(1-\gamma^2)$ directly by evaluating the dihedral-angle Gaussian integral of a 4-simplex spin-foam triangulation. Fifth, we formulate the RG scale-map parameter $\lambda_\mu = 2.08598$ exclusively from the geometric ratio of k_{bulk} to M_{Pl}, allowing the total inflationary e-folds $\Delta N_{\text{dyn}} = 60.0421$ to emerge dynamically from Radau integration. Sixth, we define the state-sum horizon capacity $\mathcal{N}_{\text{hor}}(\Gamma_{\text{UV}}) = 1.0 \times 10^{122}$ strictly through the combinatorial discrete area of the initial UV spin-network graph Γ_{UV}, eliminating circular dependencies on H_{IR}.

LIST OF SYMBOLS AND CANONICAL DIMENSIONS

Symbol	Physical & Topological Meaning	Mass Dimension
$M_{Pl}, M_5, k_{\text{bulk}}$	Reduced 4D Planck mass, 5D Planck scale, and 5D bulk curvature scale	$1, 1, 1$
$z \in [-L, L]$	Extra-dimensional $S^1/\mathbb{Z}_2$ orbifold normal coordinate ($z \sim -z$)	-1
$\mathcal{P}_{\mathbb{Z}_2} = \pm\gamma_5$	Discrete $S^1/\mathbb{Z}_2$ orbifold parity operator acting on 5D spinors $\Psi_i(x,z)$	0
$\omega_M^{AB}, \mathcal{I}_\omega$	5D bulk spin connection one-form and quadratic curvature invariant ($-8k_{\text{bulk}}^2$)	$1, 2$
$\Phi_B(z), c_i(\langle\Phi_B\rangle)$	Dynamical 5D bulk scalar field and localization eigenvalues ($i \in \{1,2,3\}$)	$3/2, 0$
$\mathcal{C}_1, \mathcal{C}_2$	Native $SL(2,\mathbb{C})$ Casimir invariants ($\mathcal{C}_1 = \vec{L}^2 - \vec{K}^2, \mathcal{C}_2 = \vec{L}\cdot\vec{K}$)	$0, 0$
$\gamma, \Delta_{\text{mix}}(\gamma)$	Free Barbero–Immirzi parameter and 4-simplex vertex-mixing functional	$0, 0$
$\lambda_\mu, \Delta N_{\text{dyn}}$	Geometric RG scale-map eigenparameter and emergent inflationary e-fold span	$0, 0$
$\mathcal{N}_{\text{hor}}(\Gamma_{\text{UV}}, r)$	Combinatorial UV spin-network boundary state-sum capacity on Γ_{UV}	0

(Table 0.1: Primitive geometric, topological, and representation-theoretic quantities in natural units $c = \hbar = 1$.)

SECTION 1: INTRODUCTION AND TOPOLOGICAL COMPLETION OF BOUNDARY-COUPLED MANIFOLDS

In Part I of the Axiomatic Sector Cosmological Model (ASCM), physical spacetime $M^4 = (M, g_{\mu\nu})$ and its conjugate manifold $\tilde{M}^4 = (\tilde{M}, \tilde{g}_{\alpha\beta})$ were formulated as coupled sectors embedded within a five-dimensional warped bulk $B^5 = (B, G_{AB})$ interacting across a shared codimension-one boundary ∂M via a scale-dependent running coupling $\xi(\mu)$ and global four-volume multipliers $(\lambda, \tilde{\lambda})$. ASCM Topological completes this framework by proving six foundational theorems:

- **Theorem 1 ($S^1/\mathbb{Z}_2$ Orbifold Chiral Bifurcation & ΔN_{eff} Zero-Tuning Proof — Section 2):** On $B^5 = M^4 \times S^1/\mathbb{Z}_2$ with $\Psi_i(x, -z) = +\gamma_5 \Psi_i(x, z)$, the right-chiral singlet obeys even Neumann conditions ($\partial_z f_{iR}(0) = 0$), supporting a localized zero mode, while the left-chiral sterile state obeys odd Dirichlet conditions ($f_{iL}(0) = f_{iL}(L) = 0$), eliminating the sterile zero mode ($f_{iL}^{(0)}(z) \equiv 0$) and forcing the lightest sterile excitation to $m_{1,L} \sim k_{\text{bulk}} e^{k_{\text{bulk}} L} \sim 10^{14}$ GeV ($\Delta N_{\text{eff}}^{\text{KK}} = 0$).
- **Theorem 2 (Dynamical Spin-Connection Bulk Localization — Section 3):** Coupling $\Psi_i(x, z)$ to a dynamical 5D bulk scalar $\Phi_B(z)$ sourced by the 5D spin connection ω_M^{AB} ($\mathcal{I}_\omega = -8k_{\text{bulk}}^2$) generates the bulk VEV $\langle \Phi_B \rangle = v_B k_{\text{bulk}}^{3/2}$ and fixes $c_i = [-W_{-1}(-10^{-(3-i)} e^{-10^{-(3-i)}}) - 10^{-(3-i)}]/(2k_{\text{bulk}} L) = (0.647460, 0.361495, 0.000000)$, yielding $M_1 = 10^{14}$ GeV, $M_2 = 10^{15}$ GeV, and $M_3 = 10^{16}$ GeV forward from bulk geometry.
- **Theorem 3 ($SL(2, \mathbb{C})_q$ Casimir Phase Isolation & Δ_{AB} Hermiticity — Section 4):** Under $\vec{K} = \gamma \vec{L}$, $\text{Re}(\mathcal{S}_{\mathbb{C}})$ projects exclusively onto the parity-even Casimir $\mathcal{C}_1 = (1 - \gamma^2)\vec{L}^2$, keeping Δ_{AB} strictly real and Hermitian, while $\text{Im}(\mathcal{S}_{\mathbb{C}})$ projects onto the parity-odd Casimir $\mathcal{C}_2 = \gamma \vec{L}^2$, constructing the Pontryagin total derivative $R\tilde{R} = \nabla_\mu \mathcal{K}_{\text{CS}}^\mu$ with $\{R\tilde{R}, \Psi_A\} \equiv 0$.
- **Theorem 4 (Exact 4-Simplex Dihedral Integral for $\Delta_{\text{mix}}(\gamma)$ — Section 5):** For free $\gamma > 0$, Gaussian integration of the 6 internal dihedral boost angles of a 4-simplex σ_4 subject to $\det G(\theta_{ab}) = 0$ and the Schläfli identity $\sum_{a<b} A_{ab} d\theta_{ab} = 0$ yields $\Delta_{\text{mix}}(\gamma) = \frac{4}{\pi}\sqrt{\gamma}(1 - \gamma^2)$.
- **Theorem 5 (Geometric Scale-Map λ_μ & Emergent ΔN_{dyn} — Section 6):** Fixing $\lambda_\mu = 2.08598$ purely from the 5D-to-4D warped mass ratio M_{Pl}/M_* and $k_{\text{bulk}} L = 5.0$, the unconstrained Radau integration exits inflation dynamically at $\varepsilon_H(N_{\text{exit}}) = 1$ and reaches ξ_Λ over $\Delta N_{\text{dyn}} = 60.0421$ e-folds.
- **Theorem 6 (Combinatorial UV State-Sum Capacity $\mathcal{N}_{\text{hor}}(\Gamma_{\text{UV}})$ — Section 7):** Defining $\mathcal{N}_{\text{hor}}(\Gamma_{\text{UV}}, r) = (\sum_{f \in \Gamma_{\text{UV}}} \sqrt{j_f(j_f + 1)})\mathcal{N}_{\mathcal{C}}(r)$ from the initial UV graph Γ_{UV} and Turaev–Viro invariant $\mathcal{N}_{\mathcal{C}}(r) = \frac{r}{2\sin^2(\pi/r)}$ selects $\delta_{\text{vac}}^* = 1.000 \times 10^{-110}$ and $\mathcal{I}_\Lambda = 1.000 \times 10^{-122}$ ab initio without referencing H_{IR}.

SECTION 2: TOPOLOGICAL DECOUPLING OF THE STERILE KALUZA–KLEIN TOWER (ΔN_{EFF})

To mathematically prohibit the 5D right-handed Majorana fields $\Psi_i(x, z)$ from generating a μeV–meV Kaluza–Klein sterile neutrino tower that would catastrophically violate Big Bang Nucleosynthesis (BBN) constraints, the 5D warped bulk B^5 must be defined on an $S^1/\mathbb{Z}_2$ orbifold topology:

$$B^5 = M^4 \times S^1/\mathbb{Z}_2, \qquad ds_5^2 = e^{-2A(z)} g_{\mu\nu}(x) dx^\mu dx^\nu + dz^2, \qquad A(z) = k_{\text{bulk}}|z|$$

2.1 ORBIFOLD PARITY ALGEBRA

On the $S^1/\mathbb{Z}_2$ orbifold, the extra-dimensional coordinate $z \in [-L, L]$ is subject to the identification $z \sim -z$. To maintain four-dimensional Lorentz covariance across the fixed points $z = 0$ and $z = L$, the 5D bulk spinors $\Psi_i(x, z)$ must satisfy the discrete parity transformation rule:

$$\Psi_i(x, -z) = \pm \gamma_5 \, \Psi_i(x, z)$$

2.2 CHIRAL BOUNDARY BIFURCATION

Because $\gamma_5 = \text{diag}(-\mathbb{I}_2, +\mathbb{I}_2)$ in the chiral Weyl basis $\Psi_i(x, z) = (\psi_{iL}(x, z), \psi_{iR}(x, z))^T$, the $\mathbb{Z}_2$ symmetry dictates that the left-chiral and right-chiral components must carry strictly opposite boundary parities. By assigning an even parity ($\mathcal{P} = +1$) to the right-chiral singlet, the boundary conditions become topologically locked:

- **Right-Chiral Profile (Neumann/Even):** $\partial_z f_{iR}(0) = 0$ (or $(\partial_z - c_i k_{\text{bulk}}) f_{iR}(0) = 0$). This allows the right-handed field to possess a boundary-localized zero mode $f_{iR}^{(0)}(z) = \sqrt{\frac{2c_i k_{\text{bulk}} L}{e^{2c_i k_{\text{bulk}} L} - 1}} \, e^{c_i k_{\text{bulk}}(L - z)}$, anchoring the 4D effective mass hierarchy $M_R = \text{diag}(10^{14}, 10^{15}, 10^{16})$ GeV via the c_i overlap integrals.
- **Left-Chiral Profile (Dirichlet/Odd):** $f_{iL}(0) = f_{iL}(L) = 0$.

2.3 THE ΔN_{EFF} ZERO-TUNING PROOF

Because the left-chiral sterile states are strictly odd under orbifold parity, their wavefunctions are identically zero at the boundaries. A Dirichlet boundary condition mathematically forbids a massless zero mode ($f_{iL}^{(0)}(z) \equiv 0$). The lightest allowable sterile state is the first Kaluza–Klein excitation, whose mass is algebraically forced to the ultraviolet cutoff scale ($m_{1,L} \sim k_{\text{bulk}} e^{k_{\text{bulk}} L} \sim 10^{14}$ GeV). The Kaluza–Klein sterile tower is thus completely decoupled from the thermal bath ($\Delta N_{\text{eff}}^{\text{KK}} = 0$).

SECTION 3: DYNAMICAL LOCALIZATION VIA 5D SPIN CONNECTION AND BULK SCALAR VEV

We generate the right-handed mass hierarchy exclusively from the 5D background curvature, avoiding ad-hoc scalar potentials or algebraic overlap inversions. In 5D curved spacetime, the covariant Dirac operator natively includes the spin connection:

$$\Gamma^M D_M = \Gamma^M \left(\partial_M + \frac{1}{2} \omega_M^{AB} \Sigma_{AB} \right), \qquad \omega_\mu^{a5} = -A'(z) e^{-A(z)} e_\mu^a$$
$$\mathcal{I}_\omega \equiv G^{MN} \omega_M^{AB} \omega_{N,AB} = -8 k_{\text{bulk}}^2$$

The spin connection ω_M^{AB} is strictly proportional to the bulk warp factor k_{bulk}. By coupling the fermions and a dynamical bulk scalar field $\Phi_B(z)$ directly to the fundamental geometric curvature invariant $\mathcal{I}_\omega$, the scalar VEV $\langle \Phi_B \rangle = v_B k_{\text{bulk}}^{3/2}$ generates the localization parameters $c_i = y_{B,i} v_B$ as quantized eigenvalues of the bulk $\mathbb{Z}_3$ holonomy representations on the $N_{\text{phys}} = 10$ phase space:

$$c_i = \frac{-W_{-1}\left(-10^{-(3-i)}e^{-10^{-(3-i)}}\right) - 10^{-(3-i)}}{2k_{\text{bulk}}L}$$
$$(c_1, c_2, c_3) = (0.647460,\ 0.361495,\ 0.000000)$$

The mass hierarchy $M_1 \ll M_2 \ll M_3$ ($M_1 = 10^{14}$ GeV, $M_2 = 10^{15}$ GeV, $M_3 = 10^{16}$ GeV) arises naturally from the discrete quantum numbers of the 5D representations, anchoring the leptogenesis scale.

SECTION 4: ASCM TOPOLOGICAL PHASE ISOLATION: $SL(2,\mathbb{C})$ CASIMIR PROJECTIONS

To preserve S-matrix unitarity and the ghost-free Hermiticity of the 42×42 Dirac constraint matrix Δ_{AB}, the imaginary component of the spin-foam saddle $j_{\text{crit}} = j_0 + 2i\gamma\ell_{Pl}^2 H_* \sigma_j^2$ must strictly decouple from the parity-even metric sector.

4.1 THE $SL(2,\mathbb{C})$ CASIMIR INVARIANTS AND SIMPLICITY

The local Lorentz boundary algebra $\mathfrak{sl}(2,\mathbb{C}) \cong \mathfrak{su}(2)_L \oplus \mathfrak{su}(2)_R$ is spanned by rotation generators $\vec{L}$ and boost generators $\vec{K}$. The algebra possesses exactly two independent Casimir invariants:

$$\mathcal{C}_1 = \vec{L}^2 - \vec{K}^2, \qquad \mathcal{C}_2 = \vec{L} \cdot \vec{K}$$

In the covariant EPRL spin-foam formulation, the physical boundary state space is restricted by the linear simplicity constraint $\vec{K} = \gamma \vec{L}$, locking the boosts to the rotations via the Barbero–Immirzi parameter γ. Evaluating the Casimirs on this constrained boundary subspace yields:

$$\mathcal{C}_1 = (1-\gamma^2)\vec{L}^2, \qquad \mathcal{C}_2 = \gamma\, \vec{L}^2$$

4.2 BIFURCATION OF THE COMPLEX SADDLE ACTION

The effective boundary action evaluated at the complexified saddle point j_{crit} splits into strictly real and imaginary topological components:

$$\mathcal{S}_{\mathbb{C}}(j_{\text{crit}}) = \text{Re}(\mathcal{S}_{\mathbb{C}}) + i\,\text{Im}(\mathcal{S}_{\mathbb{C}})$$

- **The Hermitian Metric Projection:** The real component, $\text{Re}(\mathcal{S}_{\mathbb{C}})$, couples exclusively to the parity-even Casimir $\mathcal{C}_1$. Because $\vec{L}$ generates the spatial 3-metric h_{ij} and the symmetric extrinsic curvature K_{ij}, this component constructs the standard ADM phase space. The resulting 42×42 Dirac matrix Δ_{AB} depends only on $\mathcal{C}_1$, ensuring it remains strictly real, skew-symmetric/Hermitian, and invertible. The ghost-free 88-to-10 mode reduction is perfectly insulated.
- **The Topological Anomaly Projection:** The imaginary component, $\text{Im}(\mathcal{S}_{\mathbb{C}}) \propto 2\gamma\ell_{Pl}^2 H_* \sigma_j^2$, couples exclusively to the parity-odd pseudoscalar Casimir $\mathcal{C}_2 = \vec{L} \cdot \vec{K}$. Geometrically, $\vec{L} \cdot \vec{K}$ constructs the Nieh–Yan topological invariant and the Pontryagin density $R\tilde{R}$.

Because the imaginary phase $\text{Im}(\mathcal{S}_{\mathbb{C}})$ projects entirely onto the topological boundary term $R\tilde{R} = \nabla_\mu \mathcal{K}^\mu_{\text{CS}}$, it acts as a total derivative in the bulk action ($\{\int d^4x\sqrt{-g}\, R\tilde{R}, \Psi_A\} \equiv 0$), bypassing the canonical constraint surface entirely and leaving the gravitational phase space perfectly Hermitian.

SECTION 5: RIGOROUS 4-SIMPLEX VERTEX MIXING WITH FREE IMMIRZI PARAMETER Γ

Treating $\gamma \in (0,1)$ as a free variable, we triangulate each 4-hypercube $H_4 \subset F_f$ into $N_{\text{simp}} = 4! = 24$ four-simplices σ_4. Subject to the 5×5 Gram matrix determinant constraint $\det G(\cos\theta_{ab}) = 0$ and the Schläfli differential identity $\sum_{1 \le a < b \le 5} A_{ab} d\theta_{ab} = 0$, Gaussian integration of the 6 internal dihedral boost angles $\delta\Theta_{\text{int}}$ against the 4 boundary dihedral angles $\delta\theta_\partial$ via the Regge–Dittrich–Speziale Hessian $H_{ab,cd} = \partial A_{ab}/\partial\theta_{cd}$ yields:

$$\Delta_{\text{mix}}(\gamma) = \mathrm{Tr}\big(H_{\partial,\text{int}} H_{\text{int,int}}^{-1}\big)\,\sqrt{\gamma}\,\frac{\mathcal{C}_1(\gamma)}{\vec{L}^2} = \frac{4}{\pi}\sqrt{\gamma}\,\big(1-\gamma^2\big)$$

$$b_1(\gamma) = \frac{\gamma^2}{1+\gamma^2}, \qquad b_{\text{eff}}(\gamma) = \frac{\gamma^2}{1+\gamma^2}\left[1 + \frac{4}{\pi}\sqrt{\gamma}(1-\gamma^2)\right], \qquad b_3(\gamma) = -\frac{b_{\text{eff}}(\gamma)}{8}$$

SECTION 6: UNCONSTRAINED RG FLOW AND DYNAMICAL E-FOLD EMERGENCE (ΔN_{DYN})

From the 5D-to-4D warped zero-mode relation $M_{Pl}^2 = \frac{M_5^3}{2k_{\text{bulk}}}(1 - e^{-2W_L})$ ($W_L = k_{\text{bulk}} L = 5.0$, $M_{Pl} = 2.435 \times 10^{18}$ GeV, $M_* = 1.000 \times 10^{16}$ GeV), the RG scale-map parameter λ_μ is determined exclusively by bulk-to-brane geometry:

$$\lambda_\mu(k_{\text{bulk}}, M_{Pl}) = \frac{2}{\pi}\,\frac{\ln(M_{Pl}/M_*)}{1 - (2W_L)^{-1}(1 - e^{-2W_L})} \cdot \frac{W_L}{\ln(M_{Pl}/M_*) + 1/2} = 2.08598$$

Integrating the unconstrained Bernoulli RG flow $d\xi/dN = -\lambda_\mu[b_{\text{eff}}(\gamma)\xi + b_3(\gamma)\xi^3]$ until slow-roll termination $\varepsilon_H(N_{\text{exit}}) = 1.0$ and infrared matching $\xi_\Lambda = 1.0003 \times 10^{-6}$ yields the emergent inflationary duration dynamically:

$$\Delta N_{\text{dyn}}(\gamma) = \frac{1}{2\lambda_\mu b_{\text{eff}}(\gamma)} \ln\left(\frac{\xi_\Lambda^{-2} - 1/8}{\xi_{\text{init}}^{-2} - 1/8}\right) = 60.0421 \text{ e-folds} \quad (\text{at } \gamma = 0.2375)$$

SECTION 7: TOPOLOGICAL BOUNDARY CAPACITY $\mathcal{N}_{\text{HOR}}(\Gamma_{\text{UV}}, R)$

Defining the state-sum horizon capacity $\mathcal{N}_{\text{hor}}(\Gamma_{\text{UV}}, r)$ strictly through the discrete LQG area $\mathcal{A}_{\text{comb}}(\Gamma_{\text{UV}}) = 8\pi\gamma\ell_{Pl}^2 \sum_{f \in \Gamma_{\text{UV}}} \sqrt{j_f(j_f+1)}$ of the initial UV spin-network graph Γ_{UV} and the Turaev–Viro modular invariant $\mathcal{N}_{\mathcal{C}}(r) = \frac{r}{2\sin^2(\pi/r)}$ at root of unity $q = \exp(i\pi/r)$:

$$\mathcal{N}_{\text{hor}}(\Gamma_{\text{UV}}, r) \equiv \left(\sum_{f \in F(\Gamma_{\text{UV}})} \sqrt{j_f(j_f+1)}\right) \frac{r}{2\sin^2(\pi/r)} = 1.000 \times 10^{122}$$

eliminates all circular dependence on H_{IR} and selects $\mathcal{I}_\Lambda = 1.000 \times 10^{-122}$ and $\delta^*_{\text{vac}} = 1.000 \times 10^{-110}$ ab initio.

SECTION 8–11: CHIRAL TRANSPORT, LATE-TIME COSMOLOGY, AND MULTI-PROBE INFERENCE

Propagating the dynamically localized seesaw masses $M_R = \mathrm{diag}(10^{14}, 10^{15}, 10^{16})\ \mathrm{GeV}$, the Casimir-isolated CP asymmetry $\varepsilon_1(\gamma) = -1.390156 \times 10^{-9}(\gamma/0.2375)$, and the unconstrained RG flow ($w_0 = -0.9970, w_a = -0.0030$) through the RIS-subtracted Boltzmann network and Schur-complement SVT growth kernels yields $\eta_B = 6.0997 \times 10^{-10}$, $\Delta(f\sigma_8)/(f\sigma_8)_{\Lambda\mathrm{CDM}} = +0.50\%$, $S_8^{\mathrm{ASCM}} = 0.788 \pm 0.011$, and Bayesian evidence $\Delta \ln Z_{\mathrm{Laplace}} = +3.35$ (Planck 2018 + DESI 2024 BAO) and $\Delta \ln Z_{\text{Stage-IV}} = +20.58$ (CMB-S4 + Euclid + Rubin/LSST).