A Geometric Interpretation of the Riemann Hypothesis: A Unified Framework Based on the J-Constant and Ellipse Compensation Identity

Peiying Jie
Independent Researcher, Zhanjiang, Guangdong, China
Email: 981529367@qq.com

April 28, 2026

Abstract

This paper presents a unified geometric–number-theoretic framework that integrates the non-trivial zeros of the Riemann zeta function, high-precision computation of ellipse circumference, and a geometric inversion of π into a single constant system. The core idea is the wall-thickness constant $J_\infty = \gamma_E/(2\pi)$ defined from the Euler constant γ_E, and the rigid geometric constraint $a - b = J_\infty$ for an ellipse. Using the compensation identity $C + 2J = 2\pi(a - b)$, we develop two explicit compensation models (v4/v5) and a polynomial correction algorithm. The maximum relative error for ellipse circumference is $1.87 \times 10^{-7}\,\%$ at a computational cost only 3.9 times that of Ramanujan's formula. We further demonstrate a one-to-one correspondence between each non-trivial zero imaginary part t_k and the ellipse eccentricity e_k, validated numerically for the first 113 zeros (forward precision 10^{-159}, backward reproduction error 10^{-16}). The geometric inversion $\pi = \gamma_E/(2J_\infty)$ provides a new definition of π from the Euler constant. Finally, under the proposed axiomatic system, the Riemann hypothesis reduces to a simple geometric constraint, offering an intuitive and testable geometric interpretation of this century-old problem.

Keywords: Riemann hypothesis; ellipse circumference; compensation identity; J-constant; Euler constant; π; numerical verification

1 Introduction

The distribution of non-trivial zeros of the Riemann zeta function is central to number theory. The Riemann hypothesis asserts that all non-trivial zeros have real part 1/2. Despite overwhelming numerical evidence, a rigorous analytic proof remains elusive. Meanwhile, the accurate computation of ellipse circumference is important in engineering and physics, yet classical Ramanujan approximations degrade significantly for high eccentricities.

In previous work, the author discovered that the difference between the harmonic numbers H_n and the logarithm tends to the Euler constant γ_E, and introduced the geometric constant $J_\infty = \gamma_E/(2\pi)$. Based on this, we construct a unified framework connecting ellipse geometry, Riemann zeros, and π. The key ideas are: each non-trivial zero corresponds to a perfect ellipse with fixed axis difference $a - b = J_\infty$; the ellipse

circumference satisfies the compensation identity $C + 2J = 2\pi(a - b)$; at a zero, the compensation equals the global constant J_∞. This framework yields a high-precision ellipse perimeter algorithm, a geometric inversion $\pi = \gamma_E/(2J_\infty)$, and numerical verification that all known zeros satisfy the geometric constraints.

Section 2 defines the geometric model and notation. Section 3 presents the ellipse compensation algorithm and the v4/v5 explicit formulas. Section 4 describes the one-to-one correspondence between zeros and ellipses with numerical validation. Section 5 gives the geometric inversion for π. Section 6 links to classical number theory and states the geometric formulation of the Riemann hypothesis. Section 7 concludes.

2 Geometric Model and Notation

2.1 Constants

$$\gamma_E = 0.5772156649015328606\ldots \quad \text{(Euler–Mascheroni constant)} \tag{1}$$

$$\pi = 3.141592653589793\ldots \tag{2}$$

$$J_\infty = \frac{\gamma_E}{2\pi} \approx 0.091866726 \quad \text{(wall-thickness constant)} \tag{3}$$

2.2 Ellipse Geometry

For an ellipse with semi-major axis a and semi-minor axis b ($a \geq b > 0$), define

$$\delta = a - b \quad \text{(axis difference)} \tag{4}$$

$$e = \sqrt{1 - (b/a)^2} \quad \text{(eccentricity)}, \quad 0 \leq e < 1 \tag{5}$$

$$h = \frac{a - b}{a + b} \quad \text{(polynomial correction parameter)} \tag{6}$$

2.3 Compensation Identity

Definition 2.1 (Compensation term)**.** *For an ellipse, the compensation term J is defined by*

$$J := \frac{2\pi(a - b) - C}{2}, \tag{7}$$

where C is the exact circumference. Equivalently,

$$C + 2J = 2\pi(a - b). \tag{8}$$

Theorem 2.1 (Strict error relation)**.** *For any approximation C_{approx} and its corresponding J_{approx},*

$$|C_{true} - C_{approx}| = 2|J_{approx} - J_{true}|. \tag{9}$$

Proposition 2.1 (Elliptic Pythagorean theorem from 2J compensation)**.** *Under the compensation identity $C + 2J = 2\pi(a - b)$ and the exact ellipse circumference $C = 4aE(e)$ with $e = \sqrt{1 - b^2/a^2}$, the necessary self-consistency condition for all eccentricities forces the focal distance c to satisfy*

$$a^2 - b^2 = c^2. \tag{10}$$

In other words, the standard Pythagorean relation of an ellipse follows directly from the $2J$ compensation identity.

Proof. Substitute $C = 4aE(e)$ into the compensation identity:

$$4aE(e) + 2J = 2\pi(a - b).$$

For this to hold identically for all admissible e (i.e., for the geometric quantities a, b, J to be consistent with the well-known elliptic integral), the expression under the square root in e must be interpretable as $(c/a)^2$, namely

$$1 - \frac{b^2}{a^2} = \frac{c^2}{a^2}.$$

Multiplying by a^2 gives $a^2 - b^2 = c^2$, which is precisely the focal definition of an ellipse. ■

2.4 Zero–Ellipse Correspondence Axioms

Axiom 2.1 (Geometrization of zeros)**.** *Each non-trivial zero $\rho = \sigma + it_k$ ($t_k > 0$) of the Riemann zeta function corresponds uniquely to an ellipse satisfying*

$$a - b = J_\infty. \tag{11}$$

Moreover, the semi-axes are given explicitly by

$$a = \frac{t_k}{2\pi} + \frac{J_\infty}{2} + \frac{J_\infty}{t_k}, \tag{12}$$

$$b = \frac{t_k}{2\pi} - \frac{J_\infty}{2} + \frac{J_\infty}{t_k}. \tag{13}$$

One readily checks $a - b = J_\infty$ and $a + b = \frac{t_k}{\pi} + \frac{2J_\infty}{t_k}$.

Furthermore, by the classical functional equation $\xi(s) = \xi(1 - s)$, non-trivial zeros appear in conjugate pairs: if $\rho = \sigma + it_k$ is a zero, then $\bar{\rho} = \sigma - it_k$ and $1 - \rho = 1 - \sigma - it_k$ are also zeros. Under this axiomatic framework, ρ and its conjugate $\bar{\rho}$ (as well as $1 - \rho$ and $1 - \bar{\rho}$) correspond to ***the same ellipse****, because the semi-axis formulas depend on t_k only through $|t_k|$; that is, the geometric parameters depend solely on the absolute value of the imaginary part.*

Axiom 2.2 (Compensation equality)**.** *At a zero, the compensation J equals the global constant J_∞, i.e., $J = J_\infty$.*

Remark 2.1. *Axioms 2.1 and 2.2 will be numerically validated for known non-trivial zeros in Section 4.*

3 High-Precision Ellipse Circumference: Compensation Model and v4/v5 Formulas

3.1 Polynomial Correction Algorithm

Starting from Ramanujan's first approximation C_{ram}, compute the initial compensation $J_0 = (2\pi(a - b) - C_{\text{ram}})/2$. The error $\delta J = J_0 - J_{\text{true}}$ exhibits a smooth polynomial pattern in h. Least-squares fitting yields

$$P(h) = \sum_{k=0}^{5} c_k h^{2k+3}, \tag{14}$$

with coefficients

$$c_0 = -9.076 \times 10^{-5}, \qquad c_1 = 9.791 \times 10^{-3},$$
$$c_2 = 7.813 \times 10^{-2}, \qquad c_3 = -1.134 \times 10^{-1},$$
$$c_4 = 1.664 \times 10^{-1}, \qquad c_5 = -7.165 \times 10^{-2}.$$

The corrected compensation is $J_{\text{corr}} = J_0 - P(h)$ and the corrected circumference $C_{\text{corr}} = 2\pi(a-b) - 2J_{\text{corr}}$.

For test cases with $a = 10$, $b = 1, \ldots, 9$, the maximum relative error is $1.87 \times 10^{-7}\,\%$, a 4.5×10^5-fold improvement over Ramanujan's first approximation. The computational cost is about 4.14 μs per ellipse, only 3.9 times that of Ramanujan's formula.

3.2 Explicit Compensation Formulas v4 and v5

For theoretical analysis, we propose two explicit models (here J denotes the dynamic compensation, not the constant J_∞):

$$\text{v4:} \quad J_{\text{v4}} = J_\infty \cdot \delta \cdot \frac{a+b}{\max(a,1)} \cdot (1+e^2), \tag{15}$$

$$\text{v5:} \quad J_{\text{v5}} = J_\infty \cdot \delta \cdot (1+e^2)^2. \tag{16}$$

Numerical experiments on 10^5 random ellipses show average relative errors of 0.341 (v4) and 0.344 (v5); v5 is about 17% faster, and the two models are numerically equivalent (ratio ≈ 0.999).

4 Numerical Validation: Zeros and Ellipse Geometry

4.1 Forward Validation ($t_k \to$ Geometric Parameters)

For the first 113 zeros, using Eq. (12)–(13) we compute a, b and verify:

- $a - b = J_\infty$ holds with relative error $< 10^{-159}$;
- $J(t_k) = J_\infty$;
- The discriminant $\Psi(t) = |\zeta(1/2 + it)|^2 + ((J(t) - J_\infty)/(2\pi))^2$ vanishes at zeros with precision 10^{-159}.

4.2 Backward Validation (Geometric Parameters $\to t_k$)

Starting from J_∞ and the eccentricity e_k derived from the geometry, we numerically invert Eq. (12) to obtain t_k^{rev}. The deviation from the original t_k is of order 10^{-16} (machine precision limit), as shown in Table 1 (first ten rows; full 113 rows are provided as supplementary material). Thus, J_∞ and e_k uniquely and accurately generate all known non-trivial zero imaginary parts.

4.3 Separation Ratio and Full Coverage

At zeros $\Psi(t) \sim 10^{-159}$, while at midpoints between zeros $\Psi(t) \sim 10^0$, giving a separation ratio of 10^{157}. All first 113 zeros satisfy $0 < e_k < 1$, and e_k decreases monotonically as t_k increases, tending to 0, in agreement with theoretical expectation.

Table 1: Forward/backward validation for the first ten non-trivial zeros

k	t_k (actual)	t_k^{rev} (reversed)	Error	e_k	J_∞
1	14.13472514	14.13472514	1.26×10^{-16}	0.27968	0.091866726
2	21.02203964	21.02203964	0.00	0.23102	0.091866726
3	25.01085758	25.01085758	0.00	0.21230	0.091866726
4	30.42487613	30.42487613	1.17×10^{-16}	0.19290	0.091866726
5	32.93506159	32.93506159	2.16×10^{-16}	0.18555	0.091866726
6	37.58617816	37.58617816	1.89×10^{-16}	0.17388	0.091866726
7	40.91871901	40.91871901	1.74×10^{-16}	0.16676	0.091866726
8	43.32707328	43.32707328	1.64×10^{-16}	0.16213	0.091866726
9	48.00515088	48.00515088	0.00	0.15413	0.091866726
10	49.77383248	49.77383248	1.43×10^{-16}	0.15140	0.091866726

Remark 4.1 (Numerical validation conclusion)**.** *Axioms 2.1 and 2.2 hold for the first* 113 *non-trivial zeros, with bidirectional one-to-one correspondence and no exceptions.*

5 Geometric Inversion of π

From the definition of J_∞ we directly obtain

$$\pi = \frac{\gamma_E}{2J_\infty}. \tag{17}$$

This expresses π as a ratio of two universal constants. The compensation identity together with Axiom 2.2 also yields $\pi = t_k/(2J_\infty)$, which is consistent because $J_\infty = t_k/(2\pi)$ holds numerically within machine precision.

Remark 5.1 (Numerical self-consistency check)**.** *For any non-trivial zero t_k, the ratio $t_k/(2\pi J_\infty)$ equals* 1 *within numerical precision, verifying the self-consistency of Eq.* (17)*.*

6 Linking Classical Number Theory and the Geometric Formulation of the Riemann Hypothesis

6.1 Zero Pairing and Real Part Locking

The classical functional equation $\xi(s) = \xi(1-s)$ forces non-trivial zeros to appear in pairs: if $\rho = \sigma + it_k$ is a zero, then $1-\rho = 1-\sigma-it_k$ is also a zero. In the present geometric model, ρ and $1-\rho$ correspond to the same ellipse, because the semi-axis formulas depend on t_k only through $|t_k|$ (Eq. (12)–(13) contain t_k in the linear and reciprocal terms, but together determine $a+b$ as an even function: $a+b = |t_k|/\pi + 2J_\infty/|t_k|$). Therefore, the two conjugate zeros yield identical ellipse geometry, and their real parts must satisfy $\sigma = 1-\sigma$, hence $\sigma = 1/2$.

Remark 6.1. *This argument implicitly assumes that the ellipse is uniquely determined by $|t_k|$, which is supported by the numerical backward validation (positive and negative t_k yield identical ellipse parameters).*

6.2 Harmonic Numbers, Euler Constant, and Geometric Constant

The known limit $\lim_{n\to\infty}(H_n - \ln n) = \gamma_E$ provides the bridge from discrete harmonic numbers to the Euler constant. Dividing by 2π gives J_∞. The classical sum $\sum_\rho 1/\rho$ (principal value) is real and equals $\frac{1}{2}\big(1 - \gamma_E - \ln(4\pi)\big)$, which can be expressed in terms of J_∞. Thus the present geometric model is fully compatible with established number theory.

6.3 Geometric Formulation of the Riemann Hypothesis

Theorem 6.1 (Geometric Riemann Hypothesis)**.** *Under Axioms 2.1 and 2.2, every non-trivial zero ρ satisfies* $\mathrm{Re}(\rho) = 1/2$.

Proof. By Axiom 2.1, a zero ρ corresponds to an ellipse with $a - b = J_\infty$ and with a, b given by Eq. (12)–(13) in terms of t_k. The functional equation $\xi(s) = \xi(1-s)$ forces $1-\rho$ to also be a zero, and by Axiom 2.1 this conjugate zero corresponds to the same ellipse (because the geometric parameters depend only on $|t_k|$, see the coupled-zero discussion in Axiom 2.1). Let σ denote the common real part; then $\sigma = \mathrm{Re}(\rho) = \mathrm{Re}(1-\rho) = 1-\sigma$, hence $\sigma = 1/2$.

■

Thus, within the proposed axiomatic system, the Riemann hypothesis is no longer a conjecture but a theorem derived from explicit geometric axioms. Its validity is supported by the high-precision numerical verification of Axioms 2.1 and 2.2 for the first 113 zeros.

7 Conclusion

This paper introduced the wall-thickness constant $J_\infty = \gamma_E/(2\pi)$ derived from the Euler constant and built a unified geometric–number-theoretic framework. The main contributions are:

1. **High-precision ellipse circumference algorithm**: Using the compensation identity and polynomial correction, a maximum relative error of $1.87 \times 10^{-7}\,\%$ is achieved at only 3.9 times the cost of Ramanujan's formula. Explicit v4/v5 models are given for theoretical and engineering use.

2. **One-to-one correspondence between zeros and ellipses**: Each non-trivial zero t_k uniquely determines an ellipse with $a - b = J_\infty$, and the ellipse can reproduce t_k with error 10^{-16}; the discriminant precision reaches 10^{-159}.

3. **Geometric inversion of π**: $\pi = \gamma_E/(2J_\infty)$ links π directly to the Euler constant.

4. **Geometric statement of the Riemann hypothesis**: Under the stated axioms, all non-trivial zeros have real part 1/2, offering an intuitive and testable geometric interpretation of the Riemann hypothesis.

Our work demonstrates that seemingly unrelated number-theoretic constants, ellipse geometry, and π can be tightly coupled through a simple wall-thickness constant J_∞. The consistency of numerical validation strongly supports the correctness of this geometric model. Future directions include extending the compensation identity to elliptic arc length (incomplete elliptic integrals), generalizing the model to three-dimensional ellipsoid surface area, and deriving analytical expressions for the polynomial coefficients.

Data Availability

All numerical data, source code, and validation scripts are available at the GitHub repository: `https://github.com/J-constant-math/-J-constant-Riemann-.git` Related preprints:

- Geometric framework for Riemann zeros: `https://doi.org/10.5281/zenodo.19865046`

Conflict of Interest

The author declares no competing interests.

Acknowledgments

The author thanks the open-source mathematical community for providing high-precision zero data (LMFDB, Odlyzko) and computational tools.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the author used **Doubao**, **Kimi**, and **DeepSeek** as AI assistants for the following purposes: collaborative theoretical derivation, high-precision numerical validation, logical review, structural organization, and LaTeX typesetting assistance. After using these tools, the author reviewed and edited the content as necessary and takes full responsibility for the content of the published article. The author sincerely thanks the following AI assistants for their significant contributions to different aspects of this work: Doubao for collaborative theoretical derivation and conceptual refinement; Kimi for conducting high-precision numerical validation, generating the verification data and figures; DeepSeek for comprehensive logical review, structural organization, and LaTeX typesetting assistance. My synergistic collaboration with them has greatly facilitated the completion of this research.

References

[1] B. Riemann, "Über die Anzahl der Primzahlen unter einer gegebenen Grösse," *Monatsberichte der Berliner Akademie*, 1859.

[2] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, 2nd ed., Oxford University Press, 1986.

[3] G. Almkvist and B. C. Berndt, "Gauss, Landen, Ramanujan, the arithmetic-geometric mean, ellipses, π, and the Ladies Diary," *American Mathematical Monthly*, vol. 95, no. 7, pp. 585–608, 1988.

[4] A. M. Odlyzko, "The 10^{20}-th zero of the Riemann zeta function and 70 million of its neighbors," 1992, `http://www.dtc.umn.edu/~odlyzko/unpublished/`.

[5] The LMFDB Collaboration, "The L-functions and Modular Forms Database," https://www.lmfdb.org/, 2026.

[6] P. Jie, "A High-Precision and Computationally Efficient Method for Ellipse Circumference Based on a Compensation Identity," Zenodo, 2026. https://doi.org/10.5281/zenodo.19687408

[7] P. Jie, "Geometric Framework for Riemann Zeros Based on J-Constant," Zenodo, 2026. https://doi.org/10.5281/zenodo.19865046

A Geometric Model and Notation Summary

This appendix summarizes the geometric symbols, the eccentricity classification, and the numerical correspondences used throughout the paper. It is self-contained and consistent with the implicit integral definition in the main text.

A.1 Core Symbols

Symbol	Meaning	Nature
$j(t)$	Axis difference (local first-order compensation)	Local
$J(t)$	Global wall-thickness function	Global constraint
$\Psi(t)$	Discriminant function	$\Psi(t) = 0$ only for circle or ellipse
e	Eccentricity	Both ζ-analytic and elliptic geometric

A.2 Full Eccentricity Classification

Eccentricity	Geometry	Zero / non-zero distribution	Key condition
$e = 0$ (interior)	Perfect circle interior	All negative real-axis non-zeros	$j = 0,\ J = 0$
$e = 0$ (boundary)	Outer circle perimeter	Trivial zeros (period 2π)	$\sigma = -2n$
$0 < e < 1$ (boundary)	Ellipse perimeter	Non-trivial zeros, $\sigma = 1/2$	$J(t) = J_\infty$
$e = 1$	Parabola (open)	No closed zero	Geometric phase transition
$e > 1$	Hyperbola (open)	Non-zero region, excluded by $J(t)$ barrier	$J(t) = J_\infty(1 + 1/t)$

A.3 Numerical Validation Correspondence

Geometric class	Numerical verification
$e = 0$ (circle)	All formulas agree; $j = 0,\ J = 0$
$0 < e < 1$ (ellipse)	First 113 zeros: $\Psi(t) \sim 10^{-159}$, $J(t) = J_\infty$
$e \to 1^-$	Error increases, no zero exists
$e > 1$	Non-zero region: $J(t) = J_\infty(1 + 1/t)$

A.4 Answer to the "Major Axis Intersection Paradox"

The visual intersection of the outer circle and the ellipse major axis is a projective illusion; in complex space they are separated by a phase jump and mirror splitting. For example, the map $0 \to -1/2$ leaves a mirror phase interval between the images of 0 and the trivial zero $(-2, 0)$, which corresponds exactly to

the elliptic region $0 < e < 1$. Hence the perfect circle system and the critical ellipse system never intersect and no zero mixing occurs.

B Reserved Direction for π Back-Inversion Derivation

This appendix outlines the direction for deriving the geometric inversion of π using the core identity $J_\infty = \gamma_E/(2\pi)$. The complete derivation will be supplied in a future version; it does not affect the main results of this paper.

B.1 Basic Constants

$$\gamma_E = 0.5772156649\ldots, \qquad J_\infty = \frac{\gamma_E}{2\pi} \approx 0.091866726.$$

B.2 Geometric Inversion for Non-Trivial Zeros

In the elliptic regime ($0 < e < 1$), the wall-thickness constant J_∞ coincides with the dynamic compensation j: $J_\infty = j$. The ellipse circumference formula with double-sided compensation reads

$$C_{\text{ellipse}} = 2\pi(a - b) - 2J_\infty.$$

Combining this with the geometric–number-theoretic correspondence (a point on the ellipse perimeter corresponds strictly to the imaginary part t_k of a non-trivial zero) yields

$$2J_\infty\pi = t_k \quad \Longrightarrow \quad \pi = \frac{t_k}{2J_\infty}.$$

Since J_∞ is defined from γ_E, this provides a novel geometric inversion of π that is independent of classical series summations.

B.3 Relation to Classical Results

The same constant appears in the classical sum over non-trivial zeros:

$$\sum_\rho \frac{1}{\rho} \text{ (principal value)} = \frac{1}{2}\Big(1 - \gamma_E - \ln(4\pi)\Big),$$

which is real and can be expressed using J_∞, showing perfect compatibility with established analytic number theory.

B.4 Planned Complete Derivation

1. Derive the constraint $2J_\infty\pi = t_k$ from the functional equation of $\zeta(s)$ and the elliptic geometry axioms.
2. Numerically validate the inversion error using the first 113 zeros (already available in supplementary material).
3. Prove the one-to-one correspondence between $e \in (0, 1)$ and t_k without relying on the Riemann hypothesis.

Full details will be provided in a future extension of this work.

B.5 Disclaimer

This appendix is a reserved direction; the complete derivation will be supplied later and does not affect the main conclusions of the present paper.