

Environmental fluoxetine promotes skin cell proliferation and wound healing

by Roa Hanoun and Jero Eyra

Abstract Pharmaceutical pollutants, including the antidepressant fluoxetine (FLX), are increasingly detected in aquatic environments, raising concerns about their effects on non-target organisms. The original study demonstrated that environmentally-relevant FLX concentrations promote wound closure in human keratinocytes through enhanced cell proliferation and serotonin signalling. Building on these findings, the present study extends the transcriptomic analysis by examining gene expression in scratched HaCaT keratinocytes exposed to 540 ng/L FLX at 6 and 36 hours. We identified 147 DEGs at 6 hours and 242 DEGs at 36 hours, with upregulated genes associated with cell proliferation and cellular stress defence. The difference in DEG numbers compared to the original study reflects sample quality filtering applied during data processing. These results support the conclusion that environmental FLX exposure modulates gene expression in ways that may meaningfully impact skin wound healing.

Introduction

Pharmaceutical pollution has become a growing environmental concern, particularly the presence of antidepressants in aquatic ecosystems. Fluoxetine (FLX), commercially known as Prozac, is one of the most widely prescribed selective serotonin reuptake inhibitors (SSRIs). Due to incomplete metabolism in the human body and inadequate removal by wastewater treatment plants, FLX is consistently detected in surface waters at concentrations ranging from nanograms to micrograms per liter.

Serotonin plays a well-established role in peripheral tissues, including the skin, where it is involved in regulating cell proliferation and wound healing. Wound healing proceeds through three overlapping phases: inflammation, proliferation, and tissue remodeling. During the proliferation phase, keratinocytes migrate toward the wound edge and actively divide to restore the skin barrier. Given that FLX modulates serotonin signalling, it is biologically plausible that environmental FLX exposure could influence skin repair mechanisms.

The original study by (Rodriguez-Barucg et al., 2024) was the first to investigate this link experimentally, identifying 350 DEGs in FLX-exposed keratinocytes, with upregulated genes linked to cell proliferation and downregulated genes associated with mitochondrial function (Rodriguez-Barucg et al., 2024).

The present study focuses exclusively on the transcriptomic component of the original research, analysing gene expression in scratched HaCaT keratinocytes exposed to 540 ng/L FLX at 6 and 36 hours. Our central research question is: does FLX exposure significantly alter gene expression in human keratinocytes in ways that support enhanced wound healing, and how does this response evolve between 6 and 36 hours?

Materials and Methods

Cells and clinical samples

HaCaT cells are immortalised human keratinocytes derived from normal adult human skin, widely used as a model system in dermatological research, and were obtained from AddexBio. Cells were maintained in 5% CO₂ at 37 °C and subcultured every 3 days or at approximately 80% confluency. A scratch assay was performed on HaCaT monolayers to simulate wound healing, followed by FLX treatment at 540 ng/L. Conditions were defined as a combination of treatment (FLX vs. control) and time post-scratch (6 h and 36 h).

RNA sequencing

HaCaT cells were scratched three times per well in 6-well plates, and total RNA was isolated at 6 h and 36 h post-scratch using the PureLink™ RNA Mini Kit (Thermo Fisher Scientific), following the manufacturer's instructions. Library preparation and RNA sequencing were carried out by Novogene Co. Ltd. (Cambridge, UK) using the Illumina NovaSeq 6000 platform. The sequencing output consisted

of two files, one for each comparison (6 h and 36 h), each containing one FLX-exposed sample and one untreated control sample. Each file included gene annotation columns (gene name, gene ID, chromosome, start and end positions, strand, length, and biotype), raw read counts per sample, FPKM-normalised expression values, and statistical outputs including $\log_2$ fold change, p-value, and adjusted p-value (padj). Raw read counts were used as input for all downstream differential expression analysis.

Data Pre-processing

Prior to differential expression analysis, sample quality was assessed using multiple visualizations. Raw count data were normalized using Variance Stabilizing Transformation (VST) as implemented in DESeq2, and Poisson distance-based normalization was applied to measure dissimilarity between samples. Sample-to-sample distances were visualized using heatmaps and Multidimensional Scaling (MDS) plots, generated with `PoiClu`, `ggplot2` and `pheatmap` in R, alongside DESeq2 built-in plotting functions. Additionally, bar plots and density plots were produced to inspect the distribution of counts across samples.

Based on these quality control visualizations, three irregular samples were identified and removed from the analysis: C36-R1 and E36-R3 from the 36-hour dataset, and C6-R4 from the 6-hour dataset. Sample removal was performed while ensuring that each experimental group retained a minimum of three biological replicates, in order to maintain statistical reliability in the downstream analysis.

Differential Expression & Enrichment Analysis

Differential gene expression analysis was performed separately for each time point (6 h and 36 h) using the DESeq2 R package, with genes considered differentially expressed at an unadjusted p-value threshold of $p < 0.05$. Unadjusted p-values were used in place of adjusted p-values (padj), as this approach yielded results most comparable to those reported in the original study.

Three complementary enrichment analyses were performed using the `clusterProfiler` and `ReactomePA` R packages, with gene annotations provided by `org.Hs.eg.db` and `Entrez` gene IDs (ENTREZID) as the primary identifier, all at a significance threshold of $p < 0.05$ without p-value adjustment. Gene Ontology (GO) enrichment was performed across all three ontology domains — Biological Process (BP), Cellular Component (CC), and Molecular Function (MF) — as each captures a distinct aspect of gene function. KEGG pathway enrichment was applied to identify significantly represented biological pathways among the DEGs, with gene set sizes restricted between 10 and 500 genes. Reactome pathway enrichment was additionally used to identify precise molecular reactions and sub-pathways driven by the DEGs.

Results

No significant differentially expressed genes were found using a p-adjusted value $\text{padj} < 0.05$. Thus, we proceed with p-values, although with the warning that we are not using p-adjusted values. Some significance can be seen in the following p-value distribution plots for both samples.




Figure 1: P-value distribution for both sample groups. Namely, distribution shows enough values with significance to proceed with unadjusted p-values.

With a p-value cutoff of 0.05 we end up with a similar number of DEGs as the original article. Of course, due to the data cleaning we end up with a lower amount, while without cleaning it would

result in the exact number. The following volcano plots showcase the resulting DEGs.




Figure 2: Volcano plots for both sample groups. A total of 147 DEGs at +6 hours, 86 upregulated and 61 downregulated, and 242 DEGs at +36 hours, 110 upregulated and 132 downregulated.

Go enrichment analysis for the +6 hours group shows reduced metabolic functions in the cell as well as regulation of Wnt signaling, commonly implicated in regeneration, cancer and epithelial development. For the +36 hours group upregulated genes are related to cell signaling and membrane organization.




Figure 3: go terms biological process, (a) +6 hours sample group, (b) +36 hours sample group




Figure 4: go terms cellular component, (a) +6 hours sample group, (b) +36 hours sample group




Figure 5: go terms molecular function, (a) +6 hours sample group, (b) +36 hours sample group

Kegg and reactome analysis results suggest response to oxidative and chemical stress rather than disease-specific processes for the +6 hours group, typically a detoxification response. Whereas for the +36 hours group suggests regulation of cellular stress responses, proliferative signaling, metabolic adaptation, and inflammatory processes.




Figure 6: kegg enrichment, (a) +6 hours sample group, (b) +36 hours sample group




Figure 7: reactome enrichment, (a) +6 hours sample group, (b) +36 hours sample group

Conclusion

Given the results shown above cell signaling and proliferation seem to be the most notable upsides given the exposure to FLX. Notably, an initial exposure fires negative responses related to cell stress and detoxification processes in the cell. There are some weak links to formation of illnesses, specially cancer since we are talking about skin cells. The most important aspect is that we cannot entirely ensure these effects given that the p adjusted values proved no significance, possibly related to sample

noise. However, p values showed some interesting significance which could prompt further research into the topic.

Discussion

The aim of this analysis was to reproduce the differential gene expression results reported by (Rodriguez-Barucg et al., 2024) using the same RNA-seq dataset of FLX-exposed HaCaT cells at two time points (6 h and 36 h post-scratch). Prior to sample cleaning, our DESeq2 analysis yielded exactly the same number of DEGs as reported in the original article: 165 DEGs at 6 h and 350 DEGs at 36 h. Following the removal of three irregular samples identified during quality control, the number of DEGs decreased to 147 at 6 h (86 upregulated, 61 downregulated) and 242 at 36 h (110 upregulated, 132 downregulated). The genes identified after cleaning appear to be a subset of those found before cleaning, suggesting that sample removal reduced statistical power rather than introducing different signals. Notably, replicating the original DEG counts was only achievable using unadjusted p-values, despite the article stating padj was used, which remains an open question and a limitation of the current analysis. Finally, a Venn diagram comparison revealed three genes shared across both time points — NRARP, TMEM238, and IKBKGP1 — suggesting they may play an ongoing role in the cellular response to FLX during wound healing.

Bibliography

Q. Rodriguez-Barucg, A. A. Garcia, B. Garcia-Merino, T. Akinmola, T. Okotie-Eboh, T. Francis, E. Bringas, I. Ortiz, M. A. Wade, A. Dowle, D. A. Joyce, M. J. Hardman, H. N. Wilkinson, and P. Beltran-Alvarez. Environmental fluoxetine promotes skin cell proliferation and wound healing. *Environmental Pollution*, 2024. doi: 10.1016/2024/124952. URL <https://doi.org/10.1016/j.envpol.2024.124952>. [p1, 5]

Roa Hanoun
Hanze University of Applied Sciences

rjm.hanoun@st.hanze.nl

Jero Eyraas
Hanze University of Applied Sciences

j.eyraas.jimenez@st.hanze.nl