Prepared for submission to JHEP

MadStree: a package for quickly evaluating arbitrary tree-level cosmological correlator

Spongebob Squarepants, Patrick Star

Abstract: `MadStree` is a Mathematica package to compute arbitrary tree-level cosmological correlators with scalar propagators numerically. It can deal with both massless and massive propagators. In this framework, a given dS tree diagram will be reduced to corresponding master integrals, and these master integrals will be solved by IBP and differential equations numerically. The usage of this package is described in detail via some examples.

Contents

1 Introduction **1**

2 Statement on AI Use **3**

3 IBP reduction and differential equations for tree-level dS integrals **4**
- 3.1 Notation: propagators, the integral family and index conventions 4
- 3.2 IBP relations from the three building blocks 9
- 3.3 d log-form differential equations 13
- 3.4 The remaining term and the contact blocks of the DE 16
- 3.5 Boundary conditions by nested blow-up 19
- 3.6 Several propagators sharing one theta 22

4 Using the package **24**
- 4.1 Installation and loading 24
- 4.2 Input formats and core functions 25
- 4.3 Numerical evaluation 31
- 4.4 Laurent-series reconstruction 34
- 4.5 Options, validation and common pitfalls 36

5 Examples **38**
- 5.1 Package examples 38
- 5.2 Application examples 39

6 Summary and outlook **50**

A Common-theta bundles and the odd-subset expansion **51**

1 Introduction

Tree-level cosmological correlators in de Sitter (dS) space are the basic objects of inflationary perturbation theory: after the in-in Feynman rules are applied, a correlator is a sum of time integrals whose integrands contain Hankel functions for massive propagators, exponentials for the external energies, and theta functions for the time-ordered propagators[1, 2]. Many analytic methods have been developed, including the cosmological bootstrap [3–18], Mellin techniques [12, 19–26], spectral decompositions [17, 27–32] and dispersive methods [33, 34]. See also [35–39] for some recent numerical studies.

A rather different route, and the one taken in this paper, is the route that precision calculations in flat-spacetime QFT rely on: Integration-By-Parts (IBP) relations [40] and the differential equations (DE) that they induce [41–44]. An IBP relation is a linear relation

inside one integral family, so it reduces every integral of the family to a small set of master integrals; differentiating the masters and reducing again gives a closed system of first-order linear DE, and when that system is written in d log form [45] the masters follow from boundary values by plain transport. In flat space this is by now a systematic and efficient standard that has been widely adopted, and the surrounding software is a large part of the reason. Generation and solving of the IBP system are automated by FIRE [46, 47] and by Kira [48–50], while the DE are solved numerically by DiffExp [51] and by the auxiliary mass flow method with its implementation AMFlow [52–55]. The auxiliary mass flow series also shows how far the underlying logic can be pushed: [56] argues that the master integrals of a family are completely fixed once the linear relations among them are given, and works this out on examples, so computing an integral becomes a problem of linear algebra plus numerical transport, with no integration input left to supply.

That combination of generality and automation is the reason to build the same machinery for dS space, and it was set up in [39, 57, 58]. Reference [57] extends IBP from the polynomial integrands of flat space to the Hankel integrands of dS, and gives universal formulas for the iterative reduction and for the d log-form DE of an arbitrary vertex integral family; these formulas govern all tree-level correlators and are the kernel of the loop-level construction of [58]. Reference [39] solves that system by power series expansion, obtains multivariate hypergeometric expressions for the homogeneous part of an arbitrary tree, and fixes the boundary data by blowing up the degenerate point produced by the multi-energy limit. Read together, these papers indicate that the IBP/DE framework has in dS the potential it has realized in flat space: reduction and DE are algorithmic, a topology is reduced and solved once, and the cost of a computation follows the topology rather than the analytic complexity of the answer.

This paper concerns the tree level, and it starts from [39, 57] rather than rederiving them. Two ingredients needed by a numerical implementation are not yet available there. The first is the massless internal propagator. Reference [57] covers the massless case only briefly and never uses the extra reduction relations that such a propagator provides, namely the compression of the four states at its two vertices into a single two-state factor shared by the whole propagator. Because a tree of physical interest usually contains several of these propagators, we build the IBP relations from the outset out of local building blocks that already include that relation, and we re-derive from them the general iterative reduction formula and the general d log-form DE formula of an arbitrary tree (Sec. 3.2, Sec. 3.3). The second ingredient is the boundary data. Reference [57] leaves the determination of the boundary conditions outside its scope, and [39] works it out for a two-vertex correlator, remarking that the same route should extend to any number of vertices. We give that extension as one construction: a strict ordering of the vertex energies freezes every theta function to 0 or 1, the factorized time integrals then return the boundary values of all masters as products of Gamma functions, and a vertex without an external leg is supplied with a private auxiliary energy inside the same system (Sec. 3.5). Because the reduced system and its boundary data are both algorithmic and the masters are obtained by numerical transport, one run delivers values at many kinematic points at a cost that does not grow with the analytic complexity of the answer, and analytic continuation becomes a transport

of the DE instead of the search for a new representation of the result.

`MadStree` is a Mathematica package that implements the construction above and numerically evaluates arbitrary tree-level cosmological correlators of scalar fields in de Sitter space. The reduction, the d log-form differential equation, the boundary data and the numerical transport are all derived by the package from the diagram the user supplies, so a tree is evaluated without writing down any intermediate formula by hand. The user input is an arbitrary tree diagram or loop diagram[1]. `MadStree` returns the master-integral values at every requested point, with their status and error estimates, exportable to CSV/JSON; in the regulator mode it returns the certified lowest power and the Laurent coefficients of the pole and finite parts. This mode is needed because whenever the vertex time powers take special values, the time integrals are defined only after regularization: the integral can sit on a pole and the time IBP becomes singular, while the boundary expansion at $\tau \to 0$ degenerates as soon as its branch exponents resonate. Introducing a common analytic regulator ε on the time powers makes the system generic, and the physical $\varepsilon \to 0$ limit is then taken from the reconstructed Laurent series; an end-to-end application of this mode is shown in Sec. 5.2.

The paper is organized as follows. In Sec. 3 we derive the formulas underlying the package. We review the definition of the vertex integral family and the derivation of their IBP relations, differential equations, and boundary conditions. We also discuss the simplification of the massless case in this section, and the reduction of several propagators that share one theta function, whose derivation is collected in App. A. In Sec. 4 we describe how to use the package: the loading procedure, the input formats and core functions, the numerical workflow and the reconstruction of the regulator series. We also summarize the principal options and common pitfalls. In Sec. 5 we collect the examples shipped with the release and present three application examples, the last of which evaluates a resonant configuration in the regulator mode. We close with a summary and outlook in Sec. 6.

2 Statement on AI Use

The notation, the functionality and the public interfaces of this package were designed by the authors, who also indicated the overall direction and the strategy of the derivation recorded in Sec. 3; the concrete derivation was then worked out by agent programs driven by large language models. The package was implemented by those same programs on the basis of the authors' guidance and of the derivation results above; to verify the reliability of the package, the authors designed the following verification tasks, listed together with the source of the reference data in each case.

- **Single-vertex families and the massive two-vertex correlator.** The exact differential equations, the boundary coefficients and the numerically transported results are compared with the known results of [39], which are computed by an independent multivariate hypergeometric construction.

[1] For a loop diagram the package evaluates the vertex time integrals of the Schwinger-Keldysh graph; the loop-momentum integral itself is not performed.

- **The massive three-vertex tree graph.** The package results are checked against two independent sources at once: the output of an agent program that received only the previously hand-written two-vertex code and generalized it to three vertices along its own route, and the published eight-branch result of [25]. The agent output was itself first checked against that published result before being used as a reference, so the package is validated both by independently generated code and by the published numbers.

- **IBP relations and differential equations.** These are compared with the `dSIBP` package [59], which is released in the same repository as this one, `https://github.com/J7C/dSIBP-Packages`. `dSIBP` samples naive IBP seeds and solves the resulting linear reduction system, whereas the present package applies the general formulas derived in Sec. 3 directly and never re-solves an IBP system; the agreement therefore tests the closed formulas themselves.

3 IBP reduction and differential equations for tree-level dS integrals

This section derives the objects the package computes: the integral family of a tree-level dS correlator, the IBP relations that reduce the family, the d log-form differential equation (DE) of its master integrals, and the boundary data that fixes the solution. Two references underlie the construction: [57] derives the generalized IBP relations, the one-step lowering/raising recurrences and the uniform d log-form DE of an arbitrary vertex integral family, and [39] solves that system by power series expansion and fixes its boundary data by blowing up the degenerate point produced by the multi-energy limit.

Two points distinguish the present section from those references and fix the organization of the material. First, we work in the prefactor convention that the package implements, in which the Hankel block carries $z^{+\nu}$ rather than the $z^{-\nu}$ of the printed formulas of [39, 57]; relative to those papers this is the single simultaneous replacement $\nu \to -\nu$ in every occurrence, and after performing it no further translation is needed anywhere below (Sec. 3.1). Second, the one and only structural difference from [57] is the massless internal propagator: that reference never uses the extra reduction relations of a massless bulk propagator to compress the four states at the two vertices of a massless *full* propagator into a single two-state factor shared by the whole propagator, while the package does. We therefore build the IBP relations from the start out of the three local building blocks of Sec. 3.2 rather than re-deriving the single-vertex family, and we mark every quantity quoted from [57] before that extra relation is imposed by a superscript unreduced (defined in eq. (3.20)). Section 3.6 and App. A treat a further case that [57] does not discuss: several propagators that share one and the same theta function.

3.1 Notation: propagators, the integral family and index conventions

In-in Feynman rules and asymptotic behavior. Cosmological correlators are evaluated in the Schwinger–Keldysh (SK) formalism: the operator product is ordered along a closed time path, and each vertex is labelled by a contour branch, written `"+"` or `"-"`, with

one time integration $\int_{-\infty}^{0} \mathrm{d}\tau_v$ per vertex. With the positive-frequency mode function

$$u(\tau;k) = -\mathrm{i}\sqrt{\frac{\pi}{2}}\, e^{\mathrm{i}\pi(\nu/2+1/4)} H^{(d-1)/2}_{\text{Hubble}}(-\tau)^{d/2} H^{(1)}_{\nu}(-k\tau), \tag{3.1}$$

where $k = |\boldsymbol{k}|$ and $\nu^2 = d^2/4 - m^2/H^2_{\text{Hubble}}$ for a minimally coupled scalar in d spatial dimensions (ν is real or imaginary for $d = 3$), the bulk-to-bulk propagators are

$$G_{>}(k;\tau_1,\tau_2) \equiv u(\tau_1,k)u^*(\tau_2,k), \qquad G_{<}(k;\tau_1,\tau_2) \equiv u^*(\tau_1,k)u(\tau_2,k), \tag{3.2}$$

and the time-ordered combinations used by the in-in rules are

$$\begin{aligned} G_{++} &= G_{>}\theta(\tau_1-\tau_2) + G_{<}\theta(\tau_2-\tau_1), & G_{--} &= G_{<}\theta(\tau_1-\tau_2) + G_{>}\theta(\tau_2-\tau_1), \\ G_{+-} &= G_{<}, & G_{-+} &= G_{>}. \end{aligned} \tag{3.3}$$

Throughout, $\theta_{ij} \equiv \theta(\tau_i - \tau_j)$ denotes the theta function of the time difference of vertices i and j. Bulk-to-boundary propagators of massless fields, $G_{\pm} \propto (1 \mp \mathrm{i}k\tau)e^{\pm \mathrm{i}k\tau}$, enter only as pure-exponential blocks. The Hankel functions obey

$$\partial_\tau^2 H^{(1,2)}_{\nu}(-k\tau) + \frac{1}{\tau}\,\partial_\tau H^{(1,2)}_{\nu}(-k\tau) + \left(k^2 - \frac{\nu^2}{\tau^2}\right) H^{(1,2)}_{\nu}(-k\tau) = 0, \tag{3.4}$$

with the two asymptotic regimes

$$\begin{aligned} &H^{(1)}_{\nu}(-k\tau) = c_1(-k\tau)^{\nu}(1+O(\tau^2)) + c_2(-k\tau)^{-\nu}(1+O(\tau^2)), \quad \tau \to 0^-, \\ &c_1 = e^{-\mathrm{i}\pi\nu}c[\nu]\,, \quad c_2 = c[-\nu]\,, \quad c[\nu] \equiv \frac{2^{-\nu}\Gamma(-\nu)}{\mathrm{i}\pi}\,. \end{aligned} \tag{3.5}$$

and

$$\begin{aligned} H^{(1)}_{\nu}(-k\tau) &\sim \sqrt{\frac{2}{\pi}}(-k\tau)^{-1/2} e^{-\mathrm{i}k\tau - \mathrm{i}\pi(\nu/2+1/4)}, \\ H^{(2)}_{\nu}(-k\tau) &\sim \sqrt{\frac{2}{\pi}}(-k\tau)^{-1/2} e^{\mathrm{i}k\tau + \mathrm{i}\pi(\nu/2+1/4)}, \quad -k\tau \to \infty. \end{aligned} \tag{3.6}$$

The first regime fixes the boundary conditions of the time integrals; the second justifies the Wick rotation used in the numerical evaluation of Sec. 4.3.

The two-state h block. A tree-level correlator with c_V vertices contains one time integration per vertex, and each integrand is a product of the elementary factors

$$\left\{ e^{\mathrm{i}k\tau},\ H^{(1,2)}_{\nu}(-k\tau),\ \partial_\tau H^{(1,2)}_{\nu}(-k\tau),\ \theta(\tau_j - \tau_k) \right\}. \tag{3.7}$$

Following [39], the Hankel prefactor and the time derivative are absorbed into two states of one function system. We use the convention of the package, in which the prefactor is

the *positive* power of z,

$$\begin{aligned} h^{(s)}(\nu,0;z) &\equiv z^{+\nu}H_\nu^{(s)}(z), \\ h^{(s)}(\nu,1;z) &\equiv -\frac{1}{k}\,\partial_\tau h^{(s)}(\nu,0;z) = \partial_z h^{(s)}(\nu,0;z), \qquad z=-k\tau, \end{aligned} \tag{3.8}$$

with $s\in\{1,2\}$ the Hankel branch. Superscripts on h will always denote the branch, never a power or a label. Wherever we write $h(\nu,n;z)$ with the branch superscript omitted, $n\in\{0,1\}$ being the state bit, the statement is meant for each branch separately and no branch choice is implied. The alternative normalization $h(\nu,0;z)=z^{-\nu}H_\nu(z)$ used in [39, 57] is related to (3.8) by $\nu\to-\nu$, because the reflection identities $H_{-\nu}^{(1)}=e^{\mathrm{i}\pi\nu}H_\nu^{(1)}$ and $H_{-\nu}^{(2)}=e^{-\mathrm{i}\pi\nu}H_\nu^{(2)}$ carry each branch into itself, multiplied only by a z-independent phase; the two phases are inverse to one another, so a normalization built from the pair of branches, such as eq. (3.47) below, is unaffected by the trade. Equivalently, with ν_{ref} the symbol printed in those two papers, $\nu_{\mathrm{ref}}=-\nu$ *everywhere at once* — in the equation of motion below, in the Wronskian, in M_1, in the contact powers, in the recurrences and in the d log potentials. For a heavy field $\nu=\mathrm{i}\mu$ with real μ this reduces to $e^{-\pi\mu}H_{-\mathrm{i}\mu}^{(2)}(z)=H_{\mathrm{i}\mu}^{(2)}(z)$, and we shall use exactly this identity in Sec. 3.4 to trade the conjugate-order basis at the two vertices of a contracted propagator in [39] for the common-order basis of the package.

Since the Hankel equation involves ν^2 only, the two normalizations satisfy different first-order systems but the same second-order one up to $\nu\to-\nu$. In the convention (3.8), with $\boldsymbol{h}=(h(\nu,0;z),h(\nu,1;z))^T$ and $P_1\equiv\frac{1}{2}(\mathbb{I}-\sigma_3)=\mathrm{diag}(0,1)$,

$$\partial_\tau\boldsymbol{h}=\left[-\mathrm{i}k\sigma_2-\frac{1-2\nu}{\tau}\,P_1\right]\boldsymbol{h}, \tag{3.9}$$

where $\sigma_{1,2,3}$ are the Pauli matrices and $\mathbb{I}$ the 2×2 identity; the equivalent dimensionless equation of motion is

$$\partial_z^2 h(\nu,0;z)+\frac{1-2\nu}{z}\,\partial_z h(\nu,0;z)+h(\nu,0;z)=0, \tag{3.10}$$

whose constant term is 1; the physical k^2 enters only through $\partial_\tau=-k\partial_z$.

The vertex integral family and its index conventions. Every integral in this paper is written with one symbol, I, whose arguments are the sector, the integer time-power shifts and the state bits. For a vertex carrying n massive legs the one-vertex family (an n-fold Hankel vertex integral family) reads

$$\begin{gathered} \mathrm{I}^{(a_0)}_{\tilde{\boldsymbol{a}};s_1\cdots s_n}=\int_{-\infty}^{0}\mathrm{d}\tau\,(-\tau)^{\nu_0+a_0}e^{\mathrm{i}p_0\tau}\prod_{i=1}^{n}h^{(s_i)}(\nu_i,a_i;-k_i\tau), \\ a_0\in\mathbb{Z},\quad a_i\in\{0,1\},\quad s_i\in\{1,2\}, \end{gathered} \tag{3.11}$$

where the indices have the following meanings, which we reuse for the whole graph:

- ν_0 is the power of the vertex time, $s_i\in\{1,2\}$ is the Hankel branch chosen at leg i

by the SK type of the propagator that ends there, $a_i \in \{0,1\}$ is the *state bit* — the index of the two components of (3.8) — and a_0 is the integer *shift* of the vertex time power;

- $p_0 = \varsigma k_0$ is the signed energy of the vertex (Sec. 3.2), and $k_i = |\boldsymbol{k}_i|$ the momentum modulus of leg i;
- the state bits are packed into one integer by the binary convention of [57],

$$\tilde{\boldsymbol{a}} = 1 + \sum_{i=1}^{n} a_i \, 2^{\,n-i}, \tag{3.12}$$

so that the first bit is the most significant and the 2^n entries $\mathrm{I}_{\tilde{\boldsymbol{a}}} = \mathrm{I}_{\{\boldsymbol{a}\}}$ of the master vector are ordered exactly as the Kronecker products of the leg spaces; in the package this is `IntegerDigits[...,2,n]` and the same order is used for the whole graph.

For a graph the same three kinds of index are kept, only made global. Fix a sector s on a graph with c_V vertices; it has c_s time components, counted below, and n_s^{slot} two-state slots — one for each massive propagator at each of the two vertices it joins, and one shared by the two vertices of a massless full propagator — and write a master as

$$\mathrm{I}_{\tilde{\boldsymbol{r}}} \equiv \mathrm{I}_s(\boldsymbol{n};\boldsymbol{a}), \qquad \boldsymbol{n} \in \mathbb{Z}^{c_s}, \quad \boldsymbol{a} \in \{0,1\}^{n_s^{\mathrm{slot}}}, \tag{3.13}$$

where $\boldsymbol{n}$ assigns one integer time-power shift to each component (a component is a maximal set of vertices merged by the contacts; a single vertex is one component), $\boldsymbol{a}$ is the ordered string of state bits of all slots, packed into $\tilde{\boldsymbol{r}}$ by (3.12), and the sector s itself is recorded by a *sector key*: with the root propagators listed in input order $E = (e_1, \ldots, e_{N_E})$ and L_s^{ctr} the set of contracted propagators,

$$\mathrm{key}(s) = \xi_1 \xi_2 \cdots \xi_{N_E}, \qquad \xi_j = \begin{cases} 0, & e_j \in L_s^{\mathrm{ctr}}, \\ 1, & e_j \notin L_s^{\mathrm{ctr}}, \end{cases} \tag{3.14}$$

a fixed-length bit *string*, not a binary integer, so leading zeros are kept and the top sector is the all-1 string. Each contracted propagator merges two components and removes one time integration, and the graph is a tree, so

$$c_s = c_V - \left|L_s^{\mathrm{ctr}}\right|, \tag{3.15}$$

and only the top sector has as many shifts as the graph has vertices. Sectors are partially ordered by $t \prec s$ when t is reachable from s by at least one contact, i.e. when $L_t^{\mathrm{ctr}} \supsetneq L_s^{\mathrm{ctr}}$. This ordering is the tree-level counterpart of the *subsector* relation of flat-space IBP, where S is a subsector of S' if every propagator power of S is less than or equal to the corresponding power of S' [49]: the integer propagator powers that order flat-space sectors are here replaced by the contact indicator bits ξ_j, since a propagator of a tree is either kept ($\xi_j = 1$) or contracted away ($\xi_j = 0$).

These are precisely the three arguments of the integral object of the package: the master (3.13) is `MSIntegral[sectorKey, timeShifts, stateBits]`, where `sectorKey` is the string (3.14), `timeShifts` the vector $\boldsymbol{n}$ and `stateBits` the string $\boldsymbol{a}$; the same slots are used by the returned records `"sectorKey", "stateBits", "timeShifts", "integral", "normalization"`. The object maps slot-by-slot onto the corresponding `dSIBP` object `J[sectorKey, timeShifts, stateBits]` of the parent framework [59], with the same key alphabet, the same shift dimension c_s and the same bit order, so that a MadStree master can be read as a dSIBP J and back without any re-indexing.

What the notation hides. The compactness of (3.13) is paid for by hiding six pieces of data, all of which are fixed once and for all at initialization and never re-enter the algebra:

1. the Hankel prefactor $z^{\pm\nu}$ and the normalization of the mode function (3.1): the $(-\tau)^{d/2}$ and the $z^{\pm\nu}$ powers are split between the explicit vertex power ν_0 and the definition (3.8), so no propagator carries a visible prefactor;

2. the choice of the small-τ branch and of the vanishing powers at $\tau \to 0$: both are contained in the superscript $(1, 2)$ and the order ν of $h^{(s)}$, which select between the two terms of (3.5);

3. the sector normalization $\mathcal{N}_s$: the object the package propagates is $J_s(\boldsymbol{n}; \boldsymbol{a}) = \mathcal{N}_s \, \mathrm{I}_s(\boldsymbol{n}; \boldsymbol{a})$, with
$$\mathcal{N}_s = \mathcal{N}_s^{(0)} \prod_{\substack{e \in L_s^{\mathrm{ctr}} \\ \nu_e \neq 1/2}} \left(-\sigma_e \, \frac{4\mathrm{i}}{\pi} \, q_e^{2\nu_e - 1} \right), \tag{3.16}$$
one factor per contracted massive full propagator, the constant that Sec. 3.4 attaches to the contact term (3.48); ν_e is the order and q_e the momentum modulus of that propagator, and σ_e its contour sign, $+1$ for G_{++} and -1 for G_{--}; a contracted massless full propagator contributes the factor 1, and $\mathcal{N}_s^{(0)}$ is the normalization of the sector seed, equal to 1 for every example of Sec. 5;

4. the SK contour signs: the vertex phase sign ς_v of (3.17), the incidence ϵ_{ve} of (3.28) and the propagator sign σ_e of (3.27) are fixed by the vertex types and by the vertex order of each propagator, and enter only through the signed energies $p_{0,v}$ and the letters κ;

5. the theta chamber: at a boundary chart each theta is frozen to 0 or 1 by the strict energy order (Sec. 3.5), and inside the bulk the ordering of a bundle is carried by the single δ of Sec. 3.6;

6. the regulator: when a time power sits on a pole, a common analytic shift $\nu_0 \to \nu_0 + \varepsilon$ is understood (Sec. 4.4).

All six may be hidden for the same reason: none of them changes the number of slots, the slot order or the count of master integrals, which are the only things the IBP recurrences and the DE block structure read. Every formula below is therefore an identity of

the reduced vector space, independent of the kinematics-independent constants that were absorbed.

3.2 IBP relations from the three building blocks

The relative normalization of the recurrences. Compared with [57] there is exactly one difference, already stated in Sec. 3: that paper gives each vertex of a propagator its own independent two-state factor, including the two vertices of a massless full propagator, and never imposes the extra massless relations that identify them. Everything else is its construction, translated to the $+\nu$ convention and to the $(-\tau)^{\nu_0}$ time power of (3.11). Note this last point: with $(-\tau)^A$ instead of τ^{ν_0}, the derivative of the explicit power picks up a sign, $\partial_\tau(-\tau)^A = -A(-\tau)^{A-1}$, so the one-step relation carries $-\mathrm{M}_1$ where the reference carries $+\mathrm{M}_1$; the diagonal part of the DE is unaffected but the contact block of the DE changes sign, which is why our (3.53) differs from the printed [39] form by that overall sign.

Block (i): the theta-free exponential of a vertex. Each vertex v carries exactly *one* theta-free exponential factor,

$$E_v(\tau_v) = \exp(\mathrm{i}\varsigma_v k_{0,v}\tau_v)\,, \qquad \varsigma_v = \begin{cases} -1, & v \text{ is a "+" vertex,} \\ +1, & v \text{ is a "-" vertex,} \end{cases} \qquad p_{0,v} = \varsigma_v k_{0,v}, \tag{3.17}$$

so that a `"+"` vertex gives $e^{-\mathrm{i}k_{0,v}\tau_v}$ and a `"-"` vertex gives $e^{+\mathrm{i}k_{0,v}\tau_v}$. Here $k_{0,v}$ is the *single summed* external-energy parameter of the vertex, the modulus sum of the theta-free external legs attached to v; in the package it is the input field `externalLegEnergy`. eq. (3.1) of [57] likewise writes one k_0 per vertex, and this document never introduces per-leg or per-block exponential parameters, nor any sum over them. A theta-free propagator branch is not folded into this symbol: it keeps its own modulus, and that modulus is added to the energy of whichever component holds the vertex it ends on. After a contact has merged a set C of vertices into one component, the same scalar becomes the component energy

$$p_{0,C} = \sum_{v\in C} \varsigma_v k_{0,v} + \sum_{\substack{\text{theta-free propagators } e \\ \text{with exactly one vertex in } C}} (-\varsigma_{v_e})\, q_e, \tag{3.18}$$

where $q_e = |\boldsymbol{q}_e|$ is the momentum modulus of the propagator and v_e is the vertex of e that lies in C; a branch whose two vertices both lie in C contributes $+q_e - q_e = 0$ and is therefore already absent from this sum. The second sum collects the massless cross and external branches, which create no slot and only add their moduli, each with the contour sign of the vertex that holds them, into the same scalar. The first sum alone is $p_{0,v}$, so for a one-vertex component (3.18) reduces to $p_{0,v}$ exactly when that vertex carries no theta-free propagator branch.

Because $\dim \mathcal{E}_v = 1$, this block creates no state bit and no slot: it contributes nothing to M_1 and contributes the scalar $\mathrm{i}p_{0,v}\mathbb{I}$ to M_0.

Block (ii): the h system of a massive propagator. A massive propagator of order ν contributes, at each of the two vertices it joins, one two-state system (3.8) in the $\mathcal{H} \simeq \mathbb{C}^2$

of generators (3.9). This is the block of [57] unchanged, with $1-2\nu$ replacing the printed $2\nu+1$. Only the branch $s\in\{1,2\}$, fixed by the SK type of the propagator, differs between the two vertices it joins, and it does not change the generator $-\mathrm{i}k\sigma_2$.

Block (iii): the massless full propagator. Before the extra massless relations are imposed, a massless full propagator $e=(u_e,v_e)$ is exactly like a massive one: the vertex u_e carries two states $a\in\{0,1\}$, the vertex v_e carries two states $b\in\{0,1\}$, and the local space of this propagator is $\mathcal{H}_{e,u}\otimes\mathcal{H}_{e,v}\simeq\mathbb{C}^4$ with, in the natural order,

$$\mathrm{I}_e^{(4)}=(\mathrm{I}_{00},\mathrm{I}_{01},\mathrm{I}_{10},\mathrm{I}_{11})^T, \tag{3.19}$$

where the I_{ab} are the integrals (3.21) below, in the convention (3.13). Writing the unreduced full-graph space as the tensor product of all such two-state factors and of the one-dimensional vertex exponentials, and the corresponding vertex matrices as

$$\begin{aligned}
\mathcal{V}_s^{\text{unreduced}} &= \bigotimes_{v\in V_s}\mathcal{E}_v\otimes\bigotimes_{\text{active two-state factors } i}\mathcal{H}_i, \qquad \dim\mathcal{V}_s^{\text{unreduced}}=2^{\#\{\text{two-state factors}\}},\\
\mathrm{M}_{1;v}^{\text{unreduced}} &= \nu_{0,v}\mathbb{I}-\sum_{i\in H_v^{\text{unreduced}}}(1-2\nu_i)\,\mathrm{Emb}_i(P_1),\\
\mathrm{M}_{0;v}^{\text{unreduced}} &= \mathrm{i}p_{0,v}\mathbb{I}-\mathrm{i}\sum_{i\in H_v^{\text{unreduced}}}k_i\,\mathrm{Emb}_i(\sigma_2),
\end{aligned}$$

reproduces [57] verbatim in our convention: the superscript unreduced means "the two states at each vertex of every propagator, massive or massless, form an independent two-state factor, with the $+\nu$ prefactor (3.8) and the $(-\tau)^{\nu_0}$ time power", $H_v^{\text{unreduced}}$ is the set of two-state factors contributed by the propagators that end on v — the unreduced analogue of the set $H_v(s)$ used on the reduced space below — and V_s is the set of active vertices of the sector s. Here $\mathrm{Emb}_i(A)$ is the embedding of a local operator, $\mathrm{Emb}_i(A)=\mathbb{I}\otimes\cdots\otimes A\otimes\cdots\otimes\mathbb{I}$ with A in the i-th factor.

Imposing the extra massless relations is what reduces these four states at the two vertices to one shared two-state system. At the conformal value $\nu=1/2$ the positive-prefactor block degenerates: the two branches are pure exponentials, $h^{(1)}(\frac12,0;z)=-\mathrm{i}\sqrt{2/\pi}\,e^{\mathrm{i}z}$ and $h^{(2)}(\frac12,0;z)=+\mathrm{i}\sqrt{2/\pi}\,e^{-\mathrm{i}z}$, (3.10) becomes $h''+h=0$, and the derivative state is a branch eigenstate of the zeroth-order state. Carrying the derivative out explicitly,

$$\begin{aligned}
h^{(1)}(\tfrac12,1;z)&=\partial_z\left(-\mathrm{i}\sqrt{\tfrac{2}{\pi}}\,e^{\mathrm{i}z}\right)=\sqrt{\tfrac{2}{\pi}}\,e^{\mathrm{i}z}=+\mathrm{i}\,h^{(1)}(\tfrac12,0;z),\\
h^{(2)}(\tfrac12,1;z)&=\partial_z\left(+\mathrm{i}\sqrt{\tfrac{2}{\pi}}\,e^{-\mathrm{i}z}\right)=\sqrt{\tfrac{2}{\pi}}\,e^{-\mathrm{i}z}=-\mathrm{i}\,h^{(2)}(\tfrac12,0;z).
\end{aligned} \tag{3.20}$$

More generally every half-integer order is a polynomial times an exponential and reduces to $\nu=1/2$. What decides the reduction is that the two branches carry *opposite* signs in (3.20), while a single *full* propagator puts the two branches on its two vertices in both possible orders, one per theta term. Write $\sigma_e=+1$ for G_{++} and $\sigma_e=-1$ for G_{--}. For a

massless full propagator $e = (u_e, v_e)$ of momentum modulus q_e, with $\tau_u \equiv \tau_{u_e}$ and $\tau_v \equiv \tau_{v_e}$, the four entries of (3.19) are, for G_{++},

$$\begin{aligned} \mathrm{I}_{ab} = & \int_{-\infty}^{0} \mathrm{d}\tau_u \int_{-\infty}^{0} \mathrm{d}\tau_v \; \Phi_e \, \theta_{uv} \, h^{(1)}\big(\tfrac{1}{2}, a; -q_e\tau_u\big) \, h^{(2)}\big(\tfrac{1}{2}, b; -q_e\tau_v\big) \\ & + \int_{-\infty}^{0} \mathrm{d}\tau_u \int_{-\infty}^{0} \mathrm{d}\tau_v \; \Phi_e \, \theta_{vu} \, h^{(2)}\big(\tfrac{1}{2}, a; -q_e\tau_u\big) \, h^{(1)}\big(\tfrac{1}{2}, b; -q_e\tau_v\big) \,, \qquad a, b \in \{0, 1\}, \end{aligned} \tag{3.21}$$

which is (3.3) written out in the h-state basis; for G_{--} the two Hankel branches are exchanged inside each of the two terms. Here Φ_e collects every other factor of the integrand: the vertex exponentials of u and v, their time powers, and the factors of all the remaining propagators. Name the two theta chambers of the propagator, taken undifferentiated,

$$\begin{aligned} A_e = & \int_{-\infty}^{0} \mathrm{d}\tau_u \int_{-\infty}^{0} \mathrm{d}\tau_v \; \Phi_e \, \theta_{uv} \, h^{(1)}\big(\tfrac{1}{2}, 0; -q_e\tau_u\big) \, h^{(2)}\big(\tfrac{1}{2}, 0; -q_e\tau_v\big) \,, \\ B_e = & \int_{-\infty}^{0} \mathrm{d}\tau_u \int_{-\infty}^{0} \mathrm{d}\tau_v \; \Phi_e \, \theta_{vu} \, h^{(2)}\big(\tfrac{1}{2}, 0; -q_e\tau_u\big) \, h^{(1)}\big(\tfrac{1}{2}, 0; -q_e\tau_v\big) \,, \end{aligned} \tag{3.22}$$

where the branch pair of each chamber is the one carried by G_{σ_e}, so that (3.21) displays A_e and B_e at $\sigma_e = +1$. A τ_u derivative then multiplies A_e by $+\mathrm{i}\sigma_e$ and B_e by $-\mathrm{i}\sigma_e$, and a τ_v derivative multiplies them by $-\mathrm{i}\sigma_e$ and $+\mathrm{i}\sigma_e$: the sign of the eigenvalue (3.20) is fixed by the branch, and the two chambers hold the two branches in opposite order. Hence

$$\mathrm{I}_{00} = \mathrm{I}_{11} = A_e + B_e, \qquad \mathrm{I}_{10} = -\mathrm{I}_{01} = \mathrm{i}\sigma_e \, (A_e - B_e) \,, \tag{3.23}$$

for both G_{++} and G_{--}. Only two of the four states are therefore independent, but exactly two: the four states are the *sum* and the *difference* of the two chambers, and the two chambers are different objects, with complementary regions of time ordering and the two Hankel branches interchanged between u and v. No constant λ can relate $\mathrm{I}_{10} = \lambda \, \mathrm{I}_{00}$, since A_e and B_e coincide at the exchange-symmetric point $\Phi_e(\tau_u, \tau_v) = \Phi_e(\tau_v, \tau_u)$, where I_{10} vanishes while I_{00} does not, and are unequal at a generic point, where I_{10} does not vanish. Differentiating therefore does not return the same integral up to a phase, as it would for a block made of a *single* theta term: the theta-free massless branches of (3.18) are of that kind, which is why they collapse to one dimension and create no slot, while a full propagator keeps a genuine two-state factor. Choosing the shared two-state $\boldsymbol{g}_e = (\mathrm{I}_{00}, \mathrm{I}_{10})^T \in \mathcal{M}_e \simeq \mathbb{C}^2$, the constant embedding and projection are

$$\mathrm{I}_e^{(4)} = S_e \boldsymbol{g}_e, \qquad S_e = \begin{pmatrix} 1 & 0 \\ 0 & -1 \\ 0 & 1 \\ 1 & 0 \end{pmatrix}, \qquad P_e = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \qquad P_e S_e = \mathbb{I}_2, \tag{3.24}$$

and every four-dimensional operator that preserves the relation subspace (3.23) reduces to

$P_e O S_e$. In particular

$$P_e(\sigma_2 \otimes \mathbb{I}_2)S_e = \sigma_2, \qquad P_e(\mathbb{I}_2 \otimes \sigma_2)S_e = -\sigma_2, \tag{3.25}$$

so the two vertices joined by a massless full propagator act with *the same* σ_2 on *the same* two-dimensional slot, with opposite signs. This is the point of the reduction: the shared slot carries one bit r_e that enters the state string of both vertices joined by e and cannot be duplicated into two independent two-state factors. Consequently the full-graph tensor structure survives, but the vertex-by-vertex factorization $\mathcal{V}_s \overset{?}{=} \bigotimes_{v\in V_s} \mathcal{V}_v$ fails, because $\mathcal{M}_e$ cannot belong to both u_e and v_e at once. The reduced space of a sector is

$$\mathcal{V}_s = \bigotimes_{v\in V_s} \mathcal{E}_v \otimes \bigotimes_{i\in H(s)} \mathcal{H}_i \otimes \bigotimes_{e\in E_m(s)} \mathcal{M}_e, \qquad \dim \mathcal{V}_s = 2^{|H(s)|+|E_m(s)|}, \tag{3.26}$$

where $H(s)$ is the set of two-state factors that the massive propagators of s still contribute and $E_m(s)$ the set of its uncontracted massless full propagators.

The contact vector of a theta derivative is $\boldsymbol{\rho}_e^{(4)} = (0,-1,+1,0)^T$, and (3.24) identifies it exactly as $\boldsymbol{\rho}_e^{(4)} = S_e|1\rangle_e$ with the fixed column $|1\rangle_e = (0,1)^T$; the contact selector of the shared slot therefore needs no further computation. What remains to fix is the numerical factor which this derivative carries at each of the two vertices. With the propagator sign σ_e of (3.23), the two vertices u_e and v_e, taken in that order, carry the *contact weights*

$$w_e^{(u_e)} = +2\mathrm{i}\sigma_e, \qquad w_e^{(v_e)} = -2\mathrm{i}\sigma_e, \tag{3.27}$$

in the h-state basis of (3.8). They are the numbers that multiply the local selector of the contracted propagator in the injection matrix $C_{st}^{(v,1)}$ of a contact event, which is the object built in (3.49) and carried into the contact block $\Omega_{st}^{\mathrm{contact}}$ of (3.53); the phase χ_v of (3.54) records how the pair (3.27) changes once the package basis is used. The package quotient instead factors the derivative phase of each vertex out of its state, which is the constant basis change $D_{\sigma_e} = \mathrm{diag}(1, \mathrm{i}\sigma_e)$; it turns $-\mathrm{i}q_e\sigma_2$ into $\pm\mathrm{i}\sigma_e q_e \sigma_1$ and (3.27) into the pair $(-2,+2)$, and it is diagonalized by the Hadamard matrix $\frac{1}{\sqrt{2}}\left(\begin{smallmatrix}1 & 1\\ 1 & -1\end{smallmatrix}\right)$ in place of T. Both bases give the same set of energy letters: on the states (3.23) the change of basis is $D_{\sigma_e}\boldsymbol{g}_e = (A_e + B_e,\ B_e - A_e)$, which carries no σ_e at all, so D_{σ_e} only relabels the shared bit.

Local generators and the full-graph $\mathbf{M}_1$, $\mathbf{M}_0$. For each massless full propagator, whose two vertices we order as (u_e, v_e), define the incidence

$$\epsilon_{ve} = \begin{cases} +1, & v = u_e, \\ -1, & v = v_e, \\ 0, & v \notin e, \end{cases} \tag{3.28}$$

and let $H_v(s)$ be the set of two-state factors that the massive propagators of s contribute at v. Each building block then contributes to the vertex v the pair of atoms

$$\Delta \mathrm{M}_1^{(E_v)} = 0, \qquad \Delta \mathrm{M}_0^{(E_v)} = \mathrm{i}p_{0,v}\mathbb{I}, \qquad \text{the exponential of the vertex } v, \tag{3.29}$$

$$\Delta \mathrm{M}_1^{(h_i)} = -(1-2\nu_i)P_1, \quad \Delta \mathrm{M}_0^{(h_i)} = -\mathrm{i}k_i\sigma_2, \qquad \text{a massive propagator of order } \nu_i \text{ at } v, \tag{3.30}$$

$$\Delta \mathrm{M}_1^{(\mathcal{M}_e)} = 0, \qquad \Delta \mathrm{M}_0^{(\mathcal{M}_e)} = -\mathrm{i}\epsilon_{ve}q_e\sigma_2, \quad \text{a massless propagator } (\nu = \tfrac{1}{2}) \text{ at } v. \tag{3.31}$$

The second row is one of the two states that a massive propagator contributes at the vertex v it ends on; the third is the internal propagator whose ν is a half-integer, i.e. a massless one. The massless row is written once for each of the two vertices: in (3.31) the factor ϵ_{ve} carries the sign with which the same operator $\mathrm{Emb}_e(\sigma_2)$ appears in the rows of u_e and of v_e. A massless internal propagator does not enter M_1 at all: its $\nu = 1/2$ makes $1-2\nu$ vanish identically in (3.30). The two complete matrices of a vertex v in a sector s are then the sum of these atoms, each embedded in the slot it acts on, together with the $\nu_{0,v}\mathbb{I}_s$ of the vertex time power,

$$\begin{aligned}
\mathrm{M}_{1;v,s}(\nu_{0,v}) &= \nu_{0,v}\mathbb{I}_s + \mathrm{Emb}_{\mathcal{E}_v}\Big(\Delta \mathrm{M}_1^{(E_v)}\Big) + \sum_{i\in H_v(s)} \mathrm{Emb}_i\Big(\Delta \mathrm{M}_1^{(h_i)}\Big) \\
&\quad + \sum_{e\in E_m(s)} \mathrm{Emb}_e\Big(\Delta \mathrm{M}_1^{(\mathcal{M}_e)}\Big) \\
&= \nu_{0,v}\mathbb{I}_s - \sum_{i\in H_v(s)} (1-2\nu_i)\,\mathrm{Emb}_i(P_1),
\end{aligned} \tag{3.32}$$

$$\begin{aligned}
\mathrm{M}_{0;v,s} &= \mathrm{Emb}_{\mathcal{E}_v}\Big(\Delta \mathrm{M}_0^{(E_v)}\Big) + \sum_{i\in H_v(s)} \mathrm{Emb}_i\Big(\Delta \mathrm{M}_0^{(h_i)}\Big) + \sum_{e\in E_m(s)} \mathrm{Emb}_e\Big(\Delta \mathrm{M}_0^{(\mathcal{M}_e)}\Big) \\
&= \mathrm{i}p_{0,v}\mathbb{I}_s + \sum_{i\in H_v(s)} \mathrm{Emb}_i(-\mathrm{i}k_i\sigma_2) + \sum_{e\in E_m(s)} \mathrm{Emb}_e(-\mathrm{i}\epsilon_{ve}q_e\sigma_2).
\end{aligned} \tag{3.33}$$

The first line of (3.32) and the first line of (3.33) are just the atoms (3.29)–(3.31) collected at v; the second line of each is what they give once those atoms are inserted. The per-vertex form is what (3.20) already gives, and the reduction only replaces the two two-state factors at the two vertices of a massless propagator by one shared factor: ∂_{τ_v} acts on the blocks belonging to v and on nothing else, so the vertex generators remain Kronecker *sums* on the common space (3.26) and different vertices meet only on a shared massless slot, where they carry the same σ_2 and opposite ϵ_{ve}.

3.3 d log-form differential equations

Master choice and the regular block. The masters of a sector s are the integrals (3.13) at vanishing time-power shift, $\mathrm{I}_s(\mathbf{0};\boldsymbol{a})$, collected in the bit order (3.12) into the

ordered vector

$$\mathrm{I}_s(\boldsymbol{n}) = \big(\mathrm{I}_s(\boldsymbol{n};\boldsymbol{a})\big)_{\boldsymbol{a}\in\{0,1\}^{n_s^{\mathrm{slot}}}}, \qquad \mathrm{I} = \big(\mathrm{I}_s(\boldsymbol{0})\big)_{\text{all sectors } s}, \tag{3.34}$$

the second entry being the stack over all sectors reachable by contacts. They obey a d log-form DE,

$$\begin{aligned} \mathrm{dI} &= (\mathrm{d}\Omega)\cdot \mathrm{I} = \sum_x \Omega_x \cdot \mathrm{Id}x\,, \\ \Omega_{ss}^{\mathrm{reg}} &= \Omega_{\mathrm{ex},s} - \mathrm{i}\sum_{v\in V_s} U_s^{-1}\cdot \widetilde{\Omega}_{0;v,s}\cdot U_s \cdot \mathrm{M}_{1;v,s}(\nu_{0,v}+1)\,, \end{aligned} \tag{3.35}$$

where x runs over the independent kinematic variables and $\Omega_x = \partial_x\Omega$. The derivation is that of [57]: differentiate the master vector, reduce the differentiated integrals back onto the masters by applying the one-step recurrence (3.41) iteratively, and read off the connection. The massless reduction changes nothing in that route except the two matrices the answer is expressed through: $\mathrm{M}_{1;v,s}$ and $\mathrm{M}_{0;v,s}$ are now built on the reduced space (3.26), and the diagonalizer of [57] is replaced by the U_s of (3.39), which carries an extra factor on every shared slot. Because the reduced generators still commute slot by slot, that one U_s diagonalizes every vertex matrix at once, so (3.35) follows from the printed formula of [57] by this single replacement. The two ingredients are diagonal matrices indexed by the state bits,

$$\begin{aligned} \big(\widetilde{\Omega}_{0;v,s}\big)_{\boldsymbol{ba}} &\equiv \begin{cases} -\mathrm{i}\log\kappa_{v,s;\boldsymbol{a}}, & \boldsymbol{b}=\boldsymbol{a} \\ 0, & \boldsymbol{b}\neq\boldsymbol{a} \end{cases}, \\ (\Omega_{\mathrm{ex,s}})_{\boldsymbol{ba}} &\equiv \begin{cases} -\sum_{i\in H(s)} a_i(1-2\nu_i)\log k_i, & \boldsymbol{b}=\boldsymbol{a} \\ 0, & \boldsymbol{b}\neq\boldsymbol{a} \end{cases}, \end{aligned} \tag{3.36}$$

with the energy letters

$$\kappa_{v,s;\boldsymbol{a}} = p_{0,v} + \sum_{i\in H_v(s)} (2a_i-1)k_i + \sum_{e\in E_m(s)} \epsilon_{ve}(2r_e-1)q_e, \tag{3.37}$$

in which $\boldsymbol{a}$ abbreviates the full bit string, massive bits a_i and shared bits r_e together. The same r_e enters the letters of u_e and v_e with opposite signs: this is the whole regular content of the correlation carried by one shared propagator, and it is why the letter set of a graph with shared slots is still a set of linear forms in the momenta. Only the states of massive propagators contribute to Ω_{ex}, because a massless slot has $1-2\nu=0$.

$\mathrm{M}_{1;v,s}$ is diagonal in the original basis and all $\mathrm{M}_{0;v,s}$ are made of σ_2 (or, in the package quotient, of σ_1) on individual slots, so they commute and are simultaneously diagonalized by one constant matrix, the Kronecker product of the massive diagonalizer T with the

diagonalizer of the shared slots,

$$(\mathrm{M}_{1;v,s}[\nu_{0,v},\boldsymbol{\nu}])_{\boldsymbol{ba}} = \begin{cases} \nu_{0,v} - \sum_i a_i(1-2\nu_i), & \boldsymbol{b}=\boldsymbol{a} \\ 0, & \boldsymbol{b}\neq\boldsymbol{a} \end{cases}, \tag{3.38}$$

$$(\mathrm{T}_n)_{\boldsymbol{ba}} = \prod_{i=1}^{n} \mathrm{T}_{b_i a_i}, \qquad \mathrm{T} = \frac{1}{\sqrt{2}}\begin{pmatrix} 1 & -\mathrm{i} \\ -\mathrm{i} & 1 \end{pmatrix},$$

$$U_s = \bigotimes_{i\in H(s)} \mathrm{T}_i \otimes \bigotimes_{e\in E_m(s)} \mathrm{T}_e, \qquad \widetilde{\mathrm{M}}_{0;v,s} = U_s \mathrm{M}_{0;v,s} U_s^{-1} = \mathrm{diag}(\mathrm{i}\kappa_{v,s;\boldsymbol{a}}). \tag{3.39}$$

Here T diagonalizes σ_2 ($\mathrm{T}\sigma_2\mathrm{T}^{-1} = \sigma_3$); on the shared massless slots of the package quotient the same factor is instead the Hadamard matrix, which diagonalizes σ_1, and the two choices give the same set of letters (3.37) because D_{σ_e} only relabels the shared bit. The condition for one constant U_s to diagonalize every vertex matrix slot by slot is that all the Pauli directions arriving on a given slot mutually commute, which holds here because every massive slot receives only σ_2 and every shared slot only σ_2 (or, in the quotient, only σ_1). For a single vertex all of whose legs are massive the sums reduce to one term with $U = \mathrm{T}_n$ and (3.35) is the uniform formula of [39], in our convention.

No dense $2^N \times 2^N$ inversion ever appears: inverting $\mathrm{M}_{0;v,s}$ means $U_s^{-1}\,\mathrm{diag}(1/(\mathrm{i}\kappa_{v,s;\boldsymbol{a}}))U_s$, i.e. the letter-by-letter reciprocal of the scalar letters (3.39), and M_1^{-1} is the reciprocal of its diagonal entries. Changing to user or Wick coordinates does not rederive the DE, it only pulls back the one-form, and changing basis to the package quotient is the constant similarity transformation generated by the D_{σ_e}.

What the extra relations change. Three statements isolate the effect of the massless reduction on the DE. (1) A massless shared slot contributes nothing to Ω_{ex} and nothing to M_1, so the diagonal logarithmic structure in the orders ν_i is exactly the massive one. (2) It contributes one state bit r_e to the letters of both vertices with opposite incidence (3.37), so the two vertices share a bit while their recurrences stay separate closed-form matrices. (3) The theta derivative of a full propagator, massive or massless, is not a regular operator: it injects the parent relation into strictly lower sectors, which produces the off-diagonal blocks (3.53); and the time power of the merged component counts vertex merges, not contracted propagators, so the base power of a component C is

$$A_{C,s} = \sum_{v\in C} \nu_{0,v} + \sum_{e\in E_s(C)} \varrho_e + |C| - 1, \tag{3.40}$$

where $E_s(C)$ is the set of propagators contracted into C, and ϱ_e is the raw contact power of the propagator — $\varrho_e = 2\nu_e - 1$ for a massive full propagator and $\varrho_e = 0$ for a massless one, as will follow from (3.48) and (3.23). This ϱ_e is a fixed number of the propagator, not to be confused with the shared state bit r_e of (3.37). The last term is the number of vertex merges that actually occur, which is why a simultaneous contact of k shared-theta propagators contributes once and not k times (App. A).

Iterative reduction. The recurrences are per vertex and per time component: every component C of a sector has its own integer shift n_C, and one step changes exactly one of them. Since a merged component is integrated over a single time, the power and the energy that a step reads are those of the component, not of the individual vertex: the relations below are written with $\nu_{0,v}$ and $p_{0,v}$ standing for $A_{C,s}$ of (3.40) and $p_{0,C}$ of (3.18) whenever v has been merged into a larger C, and they reduce to the vertex quantities exactly when $C = \{v\}$. The same reading applies to (3.35)–(3.39). Writing $\boldsymbol{e}_C$ for the unit vector of the component that contains v, the complete one-step IBP at v is

$$-\mathrm{M}_{1;v,s}(A_{C,s} + n_C)\,\mathrm{I}_s(\boldsymbol{n} - \boldsymbol{e}_C) + \mathrm{M}_{0;v,s}\,\mathrm{I}_s(\boldsymbol{n}) + \mathcal{R}_{v,s}(\boldsymbol{n}) = 0, \tag{3.41}$$

where the minus in front of M_1 is the $(-\tau)^A$ derivative and $\mathcal{R}_{v,s}$ collects the theta contact terms of the propagators incident to v (next subsection). Solving (3.41) for the vector at $\boldsymbol{n} - \boldsymbol{e}_C$, and the same relation taken one step higher for the one at $\boldsymbol{n} + \boldsymbol{e}_C$, gives the two directions

$$A_{-,v,s}(\boldsymbol{n}) = \mathrm{M}_{1;v,s}(A_{C,s} + n_C)^{-1}\mathrm{M}_{0;v,s}, \tag{3.42}$$

$$A_{+,v,s}(\boldsymbol{n}) = U_s^{-1}\widetilde{\mathrm{M}}_{0;v,s}^{-1}U_s\,\mathrm{M}_{1;v,s}(A_{C,s} + n_C + 1), \tag{3.43}$$

so $\mathrm{I}_s(\boldsymbol{n}\pm\boldsymbol{e}_C)$ is expressed through $\mathrm{I}_s(\boldsymbol{n})$ by two diagonal inverses only, up to $\mathcal{R}$: the lowering step carries $+\mathrm{M}_{1;v,s}^{-1}\mathcal{R}_{v,s}(\boldsymbol{n})$ and the raising step $-\mathrm{M}_{0;v,s}^{-1}\mathcal{R}_{v,s}(\boldsymbol{n}+\boldsymbol{e}_C)$. Note that both signs are the opposite of the printed [57] ones: there the explicit power is τ^A and the one-step relation carries $+\mathrm{M}_1$, whereas (3.41) carries $-\mathrm{M}_1$ for the $(-\tau)^A$ power of (3.11). The lowering step is singular when an eigenvalue of M_1 vanishes, $A_{C,s}+n_C-\sum_i a_i(1-2\nu_i) = 0$; this is a property of the state and the time powers, not of the kinematics, and it is the layer the regulator of Sec. 4.4 moves off. The raising step is singular on the kinematic hypersurface $\kappa_{v,s;\boldsymbol{a}} = 0$, i.e. when the momenta become collinear and split into two opposite groups; there the DE has its poles, and the letter count is exactly the number of such hypersurfaces.

3.4 The remaining term and the contact blocks of the DE

Two different remaining terms. The contact term $\mathcal{R}_{v,s}$ in (3.41) depends on the shift at which it is evaluated, and the two directions must not be confused. Reducing the lower time powers of the child first gives the *lowering* version

$$\mathcal{R}_{v,s}^{(0)} \equiv \mathcal{R}_{v,s}(\boldsymbol{n}), \qquad \text{child reached at shift } -1, \tag{3.44}$$

while the contact that appears in the raising row (3.43) is

$$\mathcal{R}_{v,s}^{(1)} \equiv \mathcal{R}_{v,s}(\boldsymbol{n} + \boldsymbol{e}_C), \qquad \text{child reached at shift } 0. \tag{3.45}$$

The reason is the base-power rule (3.40): the raw contact integrand carries one factor $(-\tau)^{\varrho_e}$ too few to be a master of the child at vanishing shift, so $\mathcal{R}^{(0)}$ lands on child shift -1 and only $\mathcal{R}^{(1)}$ — obtained by raising the parent power by one, which raises the merged

component by one — hits the child master itself. Since the DE differentiates the masters $\mathrm{I}_s(\mathbf{0})$ and therefore uses the raising row, *the off-diagonal blocks of the DE must be built from* $\mathcal{R}^{(1)}$; substituting $\mathcal{R}^{(0)}$ gives the wrong sign and a non-closing expression.

The two-vertex massive contraction. Take two vertices u, v joined by one massive full propagator with order ν at each of its two vertices, momentum modulus k, the legs i at u and j at v; the two other legs are collected in $\boldsymbol{a}_{\hat{\imath}}$ and $\boldsymbol{b}_{\hat{\jmath}}$. The SK combination of the two h systems at u and v is, with the branch convention (3.3),

$$\begin{aligned} &h(\nu_i, a_i, -k\tau_u)\,\theta_{uv}^{(i,j)}\,h(\nu_j, b_j, -k\tau_v) \\ &\qquad \equiv h^{(1)}(\nu_i, a_i, -k\tau_u)\,\theta_{uv}\,h^{(2)}(\nu_j, b_j, -k\tau_v) + h^{(2)}(\nu_i, a_i, -k\tau_u)\,\theta_{vu}\,h^{(1)}(\nu_j, b_j, -k\tau_v), \\ &\nu_i = \nu_j \equiv \nu, \qquad k_i = k_j \equiv k, \end{aligned}$$

where the relative sign between the two terms is that of G_{++}, and G_{--} exchanges the branches. Differentiating with respect to τ_u produces, besides the regular generators, $\partial_{\tau_u}\theta_{uv} = \delta(\tau_u - \tau_v)$ and $\partial_{\tau_u}\theta_{vu} = -\delta(\tau_u - \tau_v)$, so the delta sets the two times equal, $\tau_u = \tau_v \equiv \tau$, and leaves the antisymmetric difference

$$\delta(\tau_u - \tau_v)\Big[h^{(1)}(\nu, a_i; z)\,h^{(2)}(\nu, b_j; z) - h^{(2)}(\nu, a_i; z)\,h^{(1)}(\nu, b_j; z)\Big], \qquad z = -k\tau. \tag{3.46}$$

If $a_i = b_j$ the bracket is zero identically, since the two summands then coincide; only complementary states survive, which is the origin of the selector $\delta_{a_i,1-b_j}$. For $a_i = 0, b_j = 1$ the bracket is the Wronskian of the two branches of the h system, and with (3.8) and $H_\nu^{(1)}\partial_z H_\nu^{(2)} - \partial_z H_\nu^{(1)} H_\nu^{(2)} = -4\mathrm{i}/(\pi z)$,

$$\begin{aligned} h^{(1)}(\nu, 0; z)\,h^{(2)}(\nu, 1; z) - h^{(1)}(\nu, 1; z)\,h^{(2)}(\nu, 0; z) &= z^{2\nu}\left(H_\nu^{(1)}\partial_z H_\nu^{(2)} - \partial_z H_\nu^{(1)} H_\nu^{(2)}\right) \\ &= -\frac{4\mathrm{i}}{\pi} z^{2\nu-1}. \end{aligned} \tag{3.47}$$

For $a_i = 1, b_j = 0$ the bracket is the negative of (3.47), so both cases are collected by the factor $(-1)^{a_i+1}$. Finally $z^{2\nu-1} = k^{2\nu-1}(-\tau)^{2\nu-1}$ on the physical branch, and the surviving $(-\tau)^{2\nu-1}$ multiplies the two explicit vertex powers, which gives the raw contact power $\varrho_e = 2\nu - 1$ quoted after (3.40). The remaining term of the lowering row is therefore

$$\mathcal{R}^{(0)}_{\tilde{\boldsymbol{r}};\sigma_e} = -\sigma_e\,\delta_{a_i,1-b_j}\,(-1)^{a_i+1}\,\frac{4\mathrm{i}}{\pi}\,k^{\,2\nu-1}\,\mathrm{I}_t\big(-\boldsymbol{e}_C; \boldsymbol{r}\big), \qquad \tilde{\boldsymbol{r}} = \boldsymbol{a}_i\boldsymbol{b}_{\hat{\jmath}}, \tag{3.48}$$

where σ_e is the contour sign of the contracted propagator, of the same rule as in (3.27), k and ν are its momentum modulus and order, t is the child sector reached by the contact, and $\mathrm{I}_t(-\boldsymbol{e}_C; \boldsymbol{r})$ is the child integral (3.13) of the merged component C, with state-bit string $\boldsymbol{r}$ packed into $\tilde{\boldsymbol{r}}$ and with the shift -1 on that single component; here $c_t = 1$ by (3.15), so $-\boldsymbol{e}_C = (-1)$. The child base power is $\nu_{0,u} + \nu_{0,v} + 2\nu - 1$, one below $A_{C,t}$ of (3.40), which is what the shift -1 records. The version needed by the DE is (3.48) with $\mathrm{I}_t(-\boldsymbol{e}_C; \boldsymbol{r}) \to \mathrm{I}_t(\mathbf{0}; \boldsymbol{r})$, whose base power is $\nu_{0,u} + \nu_{0,v} + 2\nu$ and which is an actual master of the child sector; the two-vertex contact of the shipped examples is written out in this

notation in (5.7).

Two comments on the relation to [39] are in order. First, the printed form of this equation in that reference carries an extra Boltzmann factor $e^{\pi \operatorname{Im}\nu}$ and the power $k^{-2\nu-1}$: it is written in the negative-prefactor convention, which replaces $k^{2\nu-1} \to k^{-2\nu-1}$, and its basis of states at the two vertices of the contracted propagator uses conjugate Hankel orders. Our basis (3.8) uses the same order ν at both vertices, and the two bases are related by the constant identity $e^{-\pi\mu} H^{(2)}_{-\mathrm{i}\mu} = H^{(2)}_{\mathrm{i}\mu}$ for $\nu = \mathrm{i}\mu$, so that in the common-order basis the explicit Boltzmann factor is absorbed into this change of basis and does not appear in (3.48). Second, the same constant $-\sigma_e \frac{4\mathrm{i}}{\pi} k^{2\nu-1}$ is what the package assigns to the sector normalization $\mathcal{N}_s$ of the child as its pinch factor; because the contact blocks of the normalized masters $J_s(\boldsymbol{n};\boldsymbol{a}) = \mathcal{N}_s\, \mathrm{I}_s(\boldsymbol{n};\boldsymbol{a})$ carry the ratio $\mathcal{N}_s/\mathcal{N}_t$, that constant cancels from the normalized DE, and the block (3.53) below has no residual pinch prefactor.

Injection structure and the contact block. For a general graph the contact part of the row at v is a sum over lower sectors,

$$\mathcal{R}^{(1)}_{v,s}(\boldsymbol{e}_C) = \sum_{t \prec s} C^{(v,1)}_{st}\, \mathrm{I}_t(\boldsymbol{0}), \tag{3.49}$$

where $C^{(v,1)}_{st}$ is the constant injection matrix of the event that takes s to t when acted on by ∂_{τ_v}: it is the tensor product of the local contact selectors of the selected propagators — $|1\rangle_e$ on a massless shared slot, the complementary-bit selector of (3.48) on a pair of massive-propagator states — times the weights (3.27), embedded in the global bit order with all spectator bits preserved by Kronecker deltas, and times the odd-subset coefficient of Sec. 3.6. By the choice of base power, the child appears at vanishing shift and the right-hand side of (3.49) is already a master vector; only if a generalized event lands on a non-zero child shift does the same recurrence have to be applied inside the child.

Now differentiate the masters with respect to the component energy. Differentiating $e^{\mathrm{i}p_{0,C}\tau_C}$ brings down $\mathrm{i}\tau_C$, and τ_C is minus the $(-\tau_C)$ that the explicit power counts, so one factor of τ_C becomes one unit of extra time power together with a factor $-\mathrm{i}$: $\partial_{p_{0,C}} \mathrm{I}_s(\boldsymbol{n};\boldsymbol{a}) = -\mathrm{i}\, \mathrm{I}_s(\boldsymbol{n}+\boldsymbol{e}_C;\boldsymbol{a})$, which is where the $(-\tau)^A$ power shows its sign. The raising row of (3.41) at $\boldsymbol{n} = \boldsymbol{0}$,

$$\mathrm{M}_{0;v,s}\, \mathrm{I}_s(\boldsymbol{e}_C) = \mathrm{M}_{1;v,s}(\nu_{0,v}+1)\, \mathrm{I}_s(\boldsymbol{0}) - \mathcal{R}^{(1)}_{v,s}(\boldsymbol{e}_C), \tag{3.50}$$

gives

$$\partial_{p_{0,C}} \mathrm{I}_s(\boldsymbol{0}) = -\mathrm{i}\, \mathrm{M}^{-1}_{0;v,s} \mathrm{M}_{1;v,s}(\nu_{0,v}+1)\, \mathrm{I}_s(\boldsymbol{0}) + \mathrm{i}\, \mathrm{M}^{-1}_{0;v,s} \mathcal{R}^{(1)}_{v,s}(\boldsymbol{e}_C). \tag{3.51}$$

Since $\widetilde{\mathrm{M}}_{0;v,s} = \operatorname{diag}(\mathrm{i}\kappa_{v,s;\boldsymbol{a}})$ and $\partial_{p_{0,C}} \kappa_{v,s;\boldsymbol{a}} = \delta_{vC}$, the diagonalization of (3.39) is exactly

$$\mathrm{M}^{-1}_{0;v,s} = U_s^{-1} \operatorname{diag}\!\left(\frac{1}{\mathrm{i}\kappa_{v,s;\boldsymbol{a}}}\right) U_s = \partial_{p_{0,C}} \left(U_s^{-1} \widetilde{\Omega}_{0;v,s} U_s\right) \equiv \partial_{p_{0,C}} \mathcal{K}_{v,s}, \tag{3.52}$$

with $\mathcal{K}_{v,s} \equiv U_s^{-1} \widetilde{\Omega}_{0;v,s} U_s$. The first term of (3.51) is therefore the $p_{0,C}$ derivative of the regular potential in (3.35), and the second one, with (3.49), is the derivative of a purely

off-diagonal potential. Reading off the coefficient of $\mathrm{I}_t(\mathbf{0})$,

$$\Omega^{\mathrm{contact}}_{st} = +\mathrm{i}\sum_{v\in V_s} \chi_v(E_{st})\,\mathcal{K}_{v,s}\,C^{(v,1)}_{st}, \qquad t \prec s, \tag{3.53}$$

where E_{st} is the set of propagators actually selected by that simultaneous event and the phase χ_v separates the two ways a shared slot enters:

$$\chi_v(E_{st}) = \begin{cases} \varsigma_v, & \text{if } E_{st} \text{ contains no massless full propagator,} \\ (-1)^{N_0(E_{st})}, & \text{otherwise,} \end{cases} \tag{3.54}$$

where $N_0(E_{st}) = \#\{e \in E_{st} : e \text{ massless full}\}$ is the number of massless full propagators contracted by the event. For a purely massive event the only sign left in the derivative is that of the exponential quotient of the differentiated vertex, which is ς_v; a pure-massive component after a contact must have a single ς, and a mixture of positive and negative phases in one component is not a case the formula describes. For an event that contracts massless full propagators, the $-2, +2$ contact weights of the package quotient relative to the $\pm 2\mathrm{i}\sigma_e$ of (3.27) contribute one sign per shared slot, hence $(-1)^{N_0}$. All remaining sign correlations are already inside $C^{(v,1)}_{st}$. Differentiating the defining integral of a single pure-massless full propagator with respect to both vertex energies and substituting (3.53) makes the DE residual vanish component by component, which is the independent check the package runs on this block.

Ordering the sectors by the number of remaining full propagators, the complete connection is block triangular,

$$\mathrm{d}\begin{pmatrix} \mathrm{I}_{s_0} \\ \mathrm{I}_{s_1} \\ \vdots \end{pmatrix} = \mathrm{d}\begin{pmatrix} \Omega_{s_0 s_0} & \Omega_{s_0 s_1} & \cdots \\ 0 & \Omega_{s_1 s_1} & \cdots \\ \vdots & \vdots & \ddots \end{pmatrix}\begin{pmatrix} \mathrm{I}_{s_0} \\ \mathrm{I}_{s_1} \\ \vdots \end{pmatrix}, \tag{3.55}$$

so the DE is solved sector by sector, top sector first: contact only ever points from a parent to a strictly lower sector, never the reverse.

3.5 Boundary conditions by nested blow-up

The last ingredient is the boundary condition at the large-energy limit. The factor $e^{\mathrm{i}p_{0,v}\tau_v}$ is suppressed when $k_{0,v} \to \infty$ after a Wick rotation, so in this limit only the region $\tau \to 0$ contributes. Taken at face value this is one multivariate limit in which all energies diverge together, and the point it approaches is in general a multivariate degenerate singularity of the differential equation: several singular divisors meet there, so a direct series expansion about that point is not legitimate. The expansion is made legitimate by blowing the point up first; the nested coordinates introduced below are exactly the blow-up charts that replace the degenerate point by a chain of one-energy limits. This continues the route taken in [39], where the boundary condition of the multivariate hypergeometric system is fixed by the same resolution of its degenerate point. In a c_V-vertex correlator there are c_V

exponential factors with energies $k_{0,v}$; assuming the ordering

$$k_{\sigma(1)} \gg k_{\sigma(2)} \gg \cdots \gg k_{\sigma(c_V)} \gg 1, \tag{3.56}$$

with a permutation σ, we define the nested coordinates

$$x_j = \frac{k_{\sigma(j+1)}}{k_{\sigma(j)}} \quad (1 \le j < c_V), \qquad x_{c_V} = \frac{1}{k_{\sigma(c_V)}}, \tag{3.57}$$

so that

$$\frac{1}{k_{\sigma(j)}} = \prod_{r=j}^{c_V} x_r, \qquad x_1, \ldots, x_{c_V} \longrightarrow 0. \tag{3.58}$$

The exponential localization scales satisfy

$$|\tau_{\sigma(1)}| \ll |\tau_{\sigma(2)}| \ll \cdots \ll |\tau_{\sigma(c_V)}| \ll 1. \tag{3.59}$$

Since all τ approach zero from the negative side, large k corresponds to small $|\tau|$; hence $k_u \gg k_v$ implies $\tau_u > \tau_v$, and therefore

$$\theta(\tau_u - \tau_v) \longrightarrow 1, \qquad \theta(\tau_v - \tau_u) \longrightarrow 0. \tag{3.60}$$

For G_{++} or G_{--}, (3.60) must be applied to every explicit theta in the propagator expansion according to the order of its argument, and for a bundle of propagators sharing one theta it is applied to that single theta, which is the boundary counterpart of Sec. 3.6: the bundle contributes one frozen 0 or 1, not one per propagator. All thetas in an arbitrary time graph therefore become 0 or 1. If a term requires a cyclic inequality, for example

$$\theta(\tau_1 - \tau_2)\theta(\tau_2 - \tau_3)\theta(\tau_3 - \tau_1), \tag{3.61}$$

then it is automatically zero in every strict-total-order chart, instead of producing a new boundary region.

Applying the analysis above fixes every theta to 0 or 1 according to the strict energy order, so all master integrals — in the top sector as well as in every subsector — factorize into a product of one-dimensional time integrals, one per *time component* of the sector: once the thetas are frozen, a component is integrated over its single time, and the legs it carries enter that one integral through their small-τ endpoint coefficients instead of producing further integrals. At the boundary $\tau \to 0$ the massive legs behave like (3.5) and a massless leg contributes its leading power or a pure exponential, so each one-dimensional integral has the form $\int_{-\infty}^{0} \mathrm{d}\tau\, (-\tau)^A e^{\mathrm{i}K\tau} = (-\mathrm{i})^{A+1}\Gamma(A+1)K^{-A-1}$, which gives a Gamma function; worked examples of the resulting boundary data and of the blow-up charts are given in [39, 57]. The boundary conditions of all master integrals are therefore products of Gamma functions with some overall coefficients, as used in the numerical evaluation of Sec. 4.3.

Only one power of (3.5) enters each of these products, and it is worth stating which

one, because that also says what the boundary data do not contain. In the convention (3.8) the two-state block behaves as

$$h^{(1)}(\nu,0;z) = c_2 + c_1 z^{2\nu}\big(1+O(z^2)\big), \qquad h^{(1)}(\nu,1;z) = 2\nu\, c_1\, z^{2\nu-1}\big(1+O(z^2)\big), \tag{3.62}$$

with $c_{1,2}$ as in (3.5), and the second Hankel branch has the same two powers with its own constants. The constant term is the one that survives in the undifferentiated state, while differentiation removes it and leaves the second power, whose exponent differs from it by $2\nu-1$. The two powers of (3.5) are therefore not two alternatives between which one chooses: they are consumed one per state bit a, and the exponent of the one-dimensional integral moves by $2\nu-1$ exactly when a bit is switched on. Each master is thereby the anchor of one Frobenius solution, and it is the pair of powers at a single leg that separates the exponents of the two solutions: in the two-master single-vertex example of [39] the leading exponents are ν_0+1 and $\nu_0-2\nu_1$, and in each of the two solutions the component of the master that it does not anchor vanishes at leading order. Differentiation also turns the constant tower into $O(z)$, so the two powers carried by a leg are 0 and 2ν in the undifferentiated state and $2\nu-1$ and 1 in the differentiated one. The power of (3.62) that a state does not use is therefore not a subleading term of that state's own solution, since the gap to the power it does use is not an integer: it is exactly one integer step above the power used by the other state of the same leg, and it therefore arrives at an integer descendant of the solution anchored by some other master, where the differential equation induces a component on this master. [39] says so explicitly after its eq. (3.16), where that coefficient contributes only to the next-to-leading term of the solution anchored by the other master and is then dropped, and the package follows the same route by storing one leading record per master, as recalled in Sec. 4.3. Higher powers of the local expansion, logarithms and Jordan blocks are generated there by the power-log recurrence of the backend from the exact residue matrix, so they are not boundary input. What the boundary supplies beyond the leading coefficients is a linear statement: fixing the component of the anchor master of a solution to one and the components of the masters that this solution does not reach to zero turns the remaining components into the solution of one linear system, which is how the leading components of a subsector enter the masters of the sector above it. A master whose leading coefficient vanishes anchors no solution and produces no record, and a linear system that cannot be solved under those conditions returns a failure instead of an assumed boundary.

One more choice enters those endpoint coefficients, and it is a different object from the two powers just discussed: which of the two Hankel branches of (3.8) each endpoint of a propagator carries. For a propagator whose two vertices still lie in two different components that choice has already been made, because the frozen theta of (3.60) selects the ordered pair of branches of one chamber. For a propagator whose two endpoints have been merged into the component being integrated, no theta is left on that propagator at all: its integrand is just the product of the two h blocks, and the ordered pair supplied by the strict energy order is one of the two arrangements of a single equal-time configuration, the other being the arrangement with the two branches exchanged. Both arrangements

carry the same exponent A of the one-dimensional integral above and the same momentum factors, so they belong to one and the same integral, and a boundary coefficient that kept only one of them would not be constant along that configuration. The package therefore replaces the product of the two endpoint coefficients of such a propagator by the mean of its two arrangements. The mean is taken over the branch index alone: the state bits are not averaged, so the exponent, the energies and the normalizations do not move, and a propagator whose two bits are equal is unchanged, because the two arrangements then coincide. Each such propagator contributes one mean of its own, so arbitrarily many propagators joining the same two vertices are covered by the same rule, and this is the boundary counterpart of the split carried out in Sec. 3.6, where the antisymmetric difference survives only on the theta that is differentiated while every unselected propagator keeps its symmetric part. Two cases are deliberately not averaged: an unordered propagator G_{+-} or G_{-+} of (3.3), whose two branches are fixed at initialization and which carries no theta, and a coincident massless shared slot, which offers no branch choice and whose odd shared state is zero.

The construction above presumes that every vertex supplies an energy with which to be ranked in (3.59) and localized in τ_v. A vertex that carries no external leg at all has no such energy: its exponential factor is absent, and the strict-order chart that freezes the thetas is not defined for it. Rather than special-casing this situation, the package adds a private *auxiliary* external-leg energy $k_{0,v}^{\mathrm{aux}}$ to the vertex inside the very same dlog system. The auxiliary symbol is an ordinary coordinate: it enters the energy letters and the component signed energies exactly as a physical $k_{0,v}$ does, so the boundary data are first constructed from the damped limit $K_v^{\mathrm{aux}} \to \infty$ of the augmented system, and the transport then runs to the physical point, where every auxiliary coordinate takes its target value $k_{0,v}^{\mathrm{aux}} = 0$. Here $K_v \equiv \mathrm{i}p_{0,v} = \mathrm{i}\varsigma_v k_{0,v}$ is the *damping energy* of the vertex: on the damping ray $k_{0,v} \to -\mathrm{i}\varsigma_v K_v$ with real positive K_v, and the exponential of (3.17) becomes the decaying $e^{K_v \tau_v}$, which is the quantity ranked in (3.56) and, for a merged component, summed into $K_C = \mathrm{i}p_{0,C}$; the package records this substitution as `dampingDefinition` in the vertex coordinates. All auxiliary coordinates are added to the system and to every requested point at once, so this is a single transport in the enlarged coordinate space rather than a succession of limits. No re-derivation takes place, and the user never supplies or sees this coordinate. Transporting to zero is a limit inside the solution, and it can land on a singular point: if the only nonzero signed energy of some component was auxiliary, its letters vanish at $k_{0,v}^{\mathrm{aux}} = 0$. Such a finite singular point returns either a finite value or an explicit divergence instead of silently producing a number.

3.6 Several propagators sharing one theta

Up to now every contact has been attached to one full propagator. A tree graph can nevertheless contain two or more full propagators whose two vertices lie on the same pair of current components, so that all of them carry the identical theta argument $\Delta = \tau_u - \tau_v$; the package groups exactly such propagators into a *common-theta bundle* $\mathcal{B}$, with $|\mathcal{B}| \geq 2$, and a time-order cycle is the typical source. This case is absent from [57], where each contact is a single propagator. Figure 1 draws the minimal such graph: two vertices, no

external legs, and four full propagators that all join the same pair of vertices, the vertical dots between the inner two propagators marking that any number of further propagators may sit inside the same bundle. The handling is best summarized in four steps, with the details in App. A.




Figure 1. A common-theta bundle of four full propagators. The vertices u and v carry no external legs, and every propagator $e_1, \dots, e_4$ joins this same pair of vertices, hence the same pair of current components at every stage of the reduction. All four propagators therefore carry the identical theta argument $\Delta = \tau_u - \tau_v$ and form one bundle $\mathcal{B}$ with $|\mathcal{B}| = 4$. The two vertices sit at the left and the right end, the outer two propagators make up the halves of one circle, and the vertical dots between the inner two propagators stand for any number of further propagators joining the same pair, so the bundle is not restricted to the four propagators drawn.

Write the bundle as, per propagator e, an ordered pair of products α_e (coefficient of $\theta(\Delta)$) and β_e (coefficient of $\theta(-\Delta)$).

1. *Multiply the thetas out first.* Since inside a bundle all thetas are the same function, $\theta(\Delta)^2 = \theta(\Delta)$ and $\theta(\Delta)\theta(-\Delta) = 0$, the product of $|\mathcal{B}|$ propagators collapses to *two* terms rather than $2^{|\mathcal{B}|}$,

$$\prod_{e\in\mathcal{B}} \Big(\theta(\Delta)\alpha_e + \theta(-\Delta)\beta_e\Big) = \theta(\Delta)\prod_{e\in\mathcal{B}} \alpha_e + \theta(-\Delta)\prod_{e\in\mathcal{B}} \beta_e. \tag{3.63}$$

2. *One theta, one delta.* Differentiating the shared theta yields a single $\delta(\Delta)$ whose coefficient is the antisymmetric difference under $\alpha \leftrightarrow \beta$, $\partial_\Delta\big[\theta(\Delta)\prod \alpha_e + \theta(-\Delta)\prod \beta_e\big] = \delta(\Delta)\big(\prod_e \alpha_e - \prod_e \beta_e\big)$, so all contact information of the bundle is concentrated in one distribution and the two vertices are forced to coincide only once.

3. *Split and change basis.* Telescoping the antisymmetric difference and then collecting in $\mathsf{d}_e = \alpha_e - \beta_e$ and $\mathsf{m}_e = (\alpha_e + \beta_e)/2$ leaves only terms with an odd number of d factors, each with a numerical coefficient,

$$\prod_{e\in\mathcal{B}} \alpha_e - \prod_{e\in\mathcal{B}} \beta_e = \sum_{\substack{\varnothing\neq S\subseteq\mathcal{B}\\ |S| \text{ odd}}} 2^{1-|S|} \prod_{e\in S} \mathsf{d}_e \prod_{e\notin S} \mathsf{m}_e. \tag{3.64}$$

4. *Read off the events.* Every non-empty odd subset S is one simultaneous contact event: the propagators of S are contracted at once, their two vertices merge once,

the component count drops by one, and the base power of the merged component follows (3.40) with $|C|-1$ merge terms. The injection matrix $C_{st}^{(v,1)}$ of (3.49) for such an event is the tensor product of the local selectors of the propagators in S, multiplied by $2^{1-|S|}$ and by the spectator factors m_e.

The collected form (3.64) rather than the term-by-term telescoping is what the program uses, because a term-by-term enumeration would count the same sector transition several times; the argument, the parity proof and the treatment of unselected propagators whose two vertices become coincident are given in App. A. The same symmetric part governs the boundary data of a coincident propagator: Sec. 3.5 explains why the two branch arrangements of one merged propagator are replaced by their mean there, one mean per merged propagator.

4 Using the package

4.1 Installation and loading

Before using package we collect the system requirements and the loading commands.

System requirements.

- A Wolfram Language kernel: Mathematica or the free Wolfram Engine (the package is a `.wl` package; no external Wolfram libraries are needed).
- Python 3.10 or newer with `python-flint>=0.6` and `sympy>=1.12` installed, for the numerical backend. MadStree follows a zero-probe policy: the interpreter is selected only by the explicit `PythonExecutable` option or, when that is `Automatic` (the default), by the `MADSTREE_PYTHON` environment variable, falling back to `"python"`.

Install the dependencies in the Python interpreter used for numerical evaluation:

```
python -m pip install "python-flint>=0.6" "sympy>=1.12"
```

Here `python` should be replaced by the selected interpreter if necessary. MadStree includes a copy of FlintNDE under `MadStree/Vendor/FlintNDE`, so a separate installation of the FlintNDE package is not required.

Loading the package. Download or clone the official release repository https://github.com/J7C/dSIBP-Packages, which contains `dSIBP`, `MadStree` and `FlintNDE`. Set `madStreeDirectory` to the `MadStree` directory in the local repository, add it to `$Path`, and load with `Needs`:

```
madStreeDirectory = ".../dSIBP-Packages/MadStree";
AppendTo[$Path, madStreeDirectory];
Needs["MadStree`"];
```

Replace the example path with the absolute path to the local `MadStree` directory. This loads `MadStree.m`, which loads `Kernel/MadStree.wl` with explicit UTF-8. Equivalently, load the kernel file directly:

```
Get[FileNameJoin[{madStreeDirectory, "Kernel", "MadStree.wl"}],
  CharacterEncoding -> "UTF-8"];
```

Loading through `Needs` requires no user encoding option. MadStree uses its bundled FlintNDE backend without requiring `Needs["FlintNDE`"]`. Use a fresh kernel when switching versions, since `Needs` does not reload an already loaded package.

After loading, `MSFlintNDEConfiguration[]` returns an association. The following expression should return `"1.0"`:

```
MSFlintNDEConfiguration[]["version"]
```

The following expression should return `True`:

```
MSFlintNDEConfiguration[]["availableQ"]
```

It confirms that `flintnde/__init__.py` exists under the configured backend directory, which is `Vendor/FlintNDE` relative to `madStreeDirectory` by default.

4.2 Input formats and core functions

Tree diagrams: `MSInitTree`. A tree is given by an association of vertices and propagators. The propagator list sits under the input key `"lines"`, but we call these objects propagators throughout. A tree with two vertices and one massless propagator between them:

```
spec = <|
  "name" -> "masslessFullEdge",
  "vertices" -> {
    <|"id" -> v1, "externalLegEnergy" -> k1,
      "timePower" -> a1, "vertexType" -> "+"|>,
    <|"id" -> v2, "externalLegEnergy" -> k2,
      "timePower" -> a2, "vertexType" -> "+"|>
  },
  "lines" -> {
    <|"type" -> "massless", "endpoints" -> {v1, v2},
      "momentum" -> q, "nu" -> 1/2|>
  }
|>;
context = MSInitTree[spec];
```

Vertex fields:

field	meaning	note
`id`	vertex symbol	required
`externalLegEnergy`	external energy parameter (e.g. `k1`)	optional; default 0
`timePower`	time-power variable a_v (e.g. `a1`)	required
`vertexType`	contour assignment `"+"` or `"-"`	required

Propagator fields:

field	meaning	note
`type`	`"massive"` or `"massless"`	required
`endpoints`	list of one external or two internal vertices	required
`momentum`	momentum variable k_i (e.g. `q`)	required
`nu`	supplied Bessel/Hankel order ν	required for massive propagators; massless default 1/2

A massless propagator joining two internal vertices is classified internally as `masslessFull` and always uses the shared two-state quotient basis of Sec. 3.3; there is no field that selects another massless representation. The initialization option `NuConvention -> "Positive"` (default) or `"Negative"` chooses the prefactor sign of $h(\nu, 0; z) = z^{\pm\nu} H_\nu(z)$ introduced in Sec. 3.1; it is fixed for the whole context and cannot be changed per call.

Propagator numbers are assigned from the input order. The Full, Cross or External class and the contour signs are derived from the `vertexType` of the two vertices the propagator joins; they are not supplied as a type value or as separate sign fields. A `"+"` vertex has phase $e^{-\mathrm{i}E\tau}$ and a `"-"` vertex has phase $e^{+\mathrm{i}E\tau}$, where E is `externalLegEnergy`. An omitted or zero external energy is represented internally by an auxiliary variable whose target value is zero. Full propagators that join the same pair of vertices are grouped into one common-theta bundle automatically; the optional top-level field `"thetaBundles"` pins such a group explicitly as a list of propagator ids, and any group whose members do not join the same pair of vertices is rejected.

Loop diagrams without loop integral: `MSInitTimeGraph`. For a pure time-order graph the vertex and propagator fields are the same, but the graph may contain cycles. A triangle cycle reads

```
cycleContext = MSInitTimeGraph[<|
  "vertices" -> {
    <|"id" -> t1, "externalLegEnergy" -> k1,
      "timePower" -> a1, "vertexType" -> "+"|>,
    <|"id" -> t2, "externalLegEnergy" -> k2,
      "timePower" -> a2, "vertexType" -> "+"|>,
    <|"id" -> t3, "externalLegEnergy" -> k3,
      "timePower" -> a3, "vertexType" -> "+"|>
  },
  "lines" -> {
    <|"type" -> "massless", "endpoints" -> {t1, t2},
      "momentum" -> q12, "nu" -> 1/2|>,
    <|"type" -> "massless", "endpoints" -> {t2, t3},
      "momentum" -> q23, "nu" -> 1/2|>,
```

```
    <|"type" -> "massless", "endpoints" -> {t3, t1},
      "momentum" -> q31, "nu" -> 1/2|>
  }
|>];
```

Here every propagator joins two internal vertices, so all three are classified internally as `masslessFull`. Once one of them is contracted, the remaining two connect the same pair of components and therefore form the common-theta bundle of Sec. 3.6.

Single-vertex integral families: `MSInitVertexFamily`. No fake topology is needed for a single vertex. The compact input follows the reference `ki/nui` convention,

```
vertexContext = MSInitVertexFamily[<|
  "ki" -> {k0, k1, k2},
  "nui" -> {a0, nu1, nu2},
  "hankelBranches" -> {1, 2}
|>, NuConvention -> "Positive"];
```

where `First[ki]` is the pure vertex phase energy and `First[nui]` the base power of $(-\tau)$; the remaining entries give, position by position, the h-block momenta and their Hankel orders ν, which are used verbatim as in Sec. 3.1. The explicit alternative `externalLegEnergy/timePower/hBlocks/exponentialBlocks` is also accepted, where each h-block is an association with the keys `momentum`, `nu` and `hankelBranch`. Unlike tree and time-graph input, `MSInitVertexFamily` rejects any field outside its own key set.

Package conventions. The package implements the h-states of (3.8) with $z = -k\tau$ and the single function name h, using

$$h(\nu, 0; z) = z^{\pm\nu} H_\nu(z), \tag{4.1}$$

where the plus sign is the default (`NuConvention->"Positive"`) and the minus sign gives the convention of [57]; the two bases are related by $\nu \to -\nu$ (Sec. 3.1). The master integrals are the $\mathrm{I}_s(\mathbf{0}; \boldsymbol{a})$ of (3.34), which on a single vertex is the one-vertex family $\mathrm{I}^{(0)}_{\tilde{\boldsymbol{a}};s_1\cdots s_n}$ of (3.11); for multi-vertex correlators they are the products and sub-sector integrals of Sec. 3.3. Each master integral is represented by `MSIntegral[sectorKey, timeShifts, stateBits]`: `sectorKey` is a binary string over the root propagators in the input order, with `1` for a propagator that is still active and `0` for a propagator contracted by a contact; `timeShifts` assigns one integer shift of the base time power to every vertex component, where a component is a maximal set of vertices merged by the contractions; `stateBits` gives the state of every two-state slot of the sector in the slot order. The master integrals are the entries with vanishing shifts, and their connection is the d log-form (3.35) of Sec. 3.3.

Let us see some examples for `MSIntegral[sectorKey, timeShifts, stateBits]`. For the three-vertex chain v_1–v_2–v_3 with internal propagators l_{12}, l_{23}, vertex energies k_1, k_2, k_3 and time powers a_1, a_2, a_3, the contact-reachable sectors are the four strings `"11"`, `"01"`, `"10"`, `"00"`,

(i) both propagators massive: l_{12} with ν_{12}, l_{23} with ν_{23}.

`sectorKey`	contracted	components	master integrals
`"11"`	—	$\{v_1\},\{v_2\},\{v_3\}$	`MSIntegral["11",{0,0,0},{a,b,c,d}]` (16)
`"01"`	l_{12}	$\{v_1,v_2\},\{v_3\}$	`MSIntegral["01",{0,0},{c,d}]` (4)
`"10"`	l_{23}	$\{v_1\},\{v_2,v_3\}$	`MSIntegral["10",{0,0},{a,b}]` (4)
`"00"`	l_{12},l_{23}	$\{v_1,v_2,v_3\}$	`MSIntegral["00",{0},{}]` (1)

(ii) l_{12} massive with ν_{12}, l_{23} massless.

`sectorKey`	contracted	components	master integrals
`"11"`	—	$\{v_1\},\{v_2\},\{v_3\}$	`MSIntegral["11",{0,0,0},{a,b,r}]` (8)
`"01"`	l_{12}	$\{v_1,v_2\},\{v_3\}$	`MSIntegral["01",{0,0},{r}]` (2)
`"10"`	l_{23}	$\{v_1\},\{v_2,v_3\}$	`MSIntegral["10",{0,0},{a,b}]` (4)
`"00"`	l_{12},l_{23}	$\{v_1,v_2,v_3\}$	`MSIntegral["00",{0},{}]` (1)

(iii) both propagators massless.

`sectorKey`	contracted	components	master integrals
`"11"`	—	$\{v_1\},\{v_2\},\{v_3\}$	`MSIntegral["11",{0,0,0},{r,s}]` (4)
`"01"`	l_{12}	$\{v_1,v_2\},\{v_3\}$	`MSIntegral["01",{0,0},{r}]` (2)
`"10"`	l_{23}	$\{v_1\},\{v_2,v_3\}$	`MSIntegral["10",{0,0},{r}]` (2)
`"00"`	l_{12},l_{23}	$\{v_1,v_2,v_3\}$	`MSIntegral["00",{0},{}]` (1)

The vertex `vertexType` field (`"+"` or `"-"`) determines the sign of the pure vertex exponential $e^{\mathrm{i}k_0\tau}$ of (3.11), and the pair of vertex types determines the SK class of each propagator, from which the default Hankel branches of (3.8) follow; a propagator may override them with its own `hankelBranches` field. The package h-states are those of (3.8), i.e. $h(\nu,1;z)=\partial_z h(\nu,0;z)$, with the alternative prefactor $z^{+\nu}$ obtained by $\nu\to-\nu$; `MSHTohMatrix`, `MShToHMatrix` and `MSConvertBasis` implement the corresponding basis change and its Kronecker products, with the massless shared two-state of Sec. 3.2 acting trivially on the reduced states.

Once the context is initialized, the following functions expose the formula products of Sec. 3.

Core functions. Most functions take the initialized `context`; `MSExportEvaluationData` takes a completed evaluation result instead.

function	purpose
`MSSectors[context]`	contact-reachable sectors in deterministic order (keys, pinched propagators, master counts)
`MSMasterIntegrals[context]`	ordered master records; each `"integral"` field is an `MSIntegral[sectorKey, timeShifts, stateBits]`
`MSFormulaMatrices[context, key]`	per-component M_1, M_0 and energy letters $\kappa_{\boldsymbol{a}}$; sector diagonalizer `U` and `UInverse`
`MSDLogDE[context]`	block-triangular d log-form potential Ω, letters, certification status
`MSReduce[expr, context]`	iterative IBP reduction to the master basis; scalar, list or nested table input
`MSContactMaps[context, key]`	parent-to-sub-sector contact maps
`MSWriteFormulaArtifacts[context]`	write formula products beside the caller
`MSFormulaData[context]`	aggregate all-sector masters, recurrence metadata and the d log-form DE
`MSEvaluatePath[context, pointSequence, opts]`	single-stage evaluation: boundary initialization and segment transport in one backend process; optional FlintNDE path planning
`MSReconstructEpSeries[context, ep, pointSequence, opts]`	Laurent series in the common analytic regulator: certified lowest power, automatic sample points, incremental fitting
`MSExportEvaluationData[evaluation, opts]`	export the saved ordinary-point records of an evaluation to CSV/JSON

`MSSectors[context]` returns the sector list. Each sector key is the fixed-length bit string over the root propagators in input order (`"0"` pinched, `"1"` unpinched); the sub-sector structure is discussed in Sec. 3.3. Keys are strings, not integers; leading zeros are part of the identity.

`MSMasterIntegrals[context]` returns the ordered master records. Their `"integral"` fields contain the `MSIntegral[sectorKey, timeShifts, stateBits]` objects of Sec. 4.2, i.e. the master integrals $I_s(\mathbf{0};\boldsymbol{a})$ of (3.34). Use `Lookup[MSMasterIntegrals[context], "integral"]` to obtain this ordered basis. Each record also contains `"bareIntegral"`, `"normalization"` and `"definition"`. Their order is the unique authority for all matrices, boundary vectors and numerical results.

`MSFormulaMatrices[context, key]` returns the component records under `"components"`, with fields `"M1"`, `"M0"` and `"energyLetters"` for the matrices and letters of Sec. 3.2. The sector diagonalizer of M_0 and its inverse are returned as `"U"` and `"UInverse"`; M_1 is already diagonal in the state basis. Here `key` is a

sector-key string; omitting it, or passing `All`, returns an association for all sectors. `MSContactMaps` accepts the same sector selection.

`MSDLogDE[context]` returns `omegaPotential`, the matrix Ω of Sec. 3.3 whose derivative along a path is the DE matrix, together with the letters (the arguments of the logarithms of (3.36)), `letterMatrices`, `dlogResidual`, `constantResidueQ` and the certification `dlogStatus`. Only `"certifiedByFormulaChecks"` is a closed d log-form and may be consumed numerically.

`MSReduce[expr, context, MasterBasis -> Automatic]` reduces finite linear combinations of legal `MSIntegrals`, or lists and nested tables of such expressions, by the lowering/raising recurrences $A_{\pm}$ of Sec. 3.2 along the sub-sector structure of Sec. 3.3. Results include `result`, `masterBasis`, `coefficientVector`, `masterRules`, `nonMasterResidual`, `remainingShiftedIntegrals` and `singularLayers`; a full reduction requires a zero residual and an empty shifted list for every scalar expression. For list input, these fields preserve the input shape; `elementResults`, `failurePositions` and `incompletePositions` provide the element-level reports.

`MSFormulaData[context]` returns all-sector masters, the per-sector $\mathrm{M}_1/\mathrm{M}_0/\mathrm{T}_n$ recurrence metadata of Sec. 3.2 and the complete block-triangular d log-form DE in the master order of `MSDLogDE`; `TimePowerRules` may substitute the vertex time powers a_i in the recurrence metadata, but the returned `"dlogDE"` remains analytic and unsubstituted.

`MSEvaluatePath[context, pointSequence, opts]` is the single-stage numerical entry for evaluating the master integrals at the user points. `pointSequence` is a coordinate table: its first row gives the ordered coordinate symbols and subsequent rows give equal-width coordinate values. Fixed parameters are supplied once through `ParameterRules`. Boundary generation, transport and result records are described in Sec. 4.3; the principal options and point tags are summarized in Sec. 4.5.

`MSReconstructEpSeries[context, ep, pointSequence, opts]` reconstructs the Laurent series of the master integrals in a common analytic regulator `ep`, with `MaximumEpPower` the highest returned power (default 0, i.e. the pole terms and the finite part). It uses the same `pointSequence` and `ParameterRules` format as `MSEvaluatePath`. Only `ParameterRules`, not the coordinates, may depend on `ep`. By default, no `ep` values need to be supplied; `EpSamplePoints` and `EpValidationPoints` may instead specify production candidates and disjoint validation points. The certificate conditions, sampling and fitting procedure are described in Sec. 4.4; the corresponding options are summarized in Sec. 4.5, and an end-to-end application is presented in Sec. 5.2.

`MSExportEvaluationData[evaluation, opts]` exports the saved ordinary-point records of one `MSEvaluatePath` result, including the coordinates, the real/imaginary parts of every master, `status`, `relativeDifferenceInf` and `targetRelativeErrorMet`, to CSV and JSON for downstream plotting tools; transient records are excluded.

The two precision fields are the corresponding segment reports, not separate error estimates computed at every point. `ExportFormats` selects CSV, JSON or both, and `SignificantDigits` defaults to 16; see Sec. 4.2.

Output organization.

- Transient request JSON and backend logs: `results_temp/nde/` beside the caller by default. Request and log files are deleted after a successful run and retained for backend failures; structured singular-path or leading-order refusals also remove these files. Successful JSON results remain in `results_temp/cache/` for reuse. `MSRuntimeDirectory` may specify a different runtime directory.
- Saved point records: `MSEvaluatePath` returns the records under `"pointResults"`. An ordinary untagged coordinate row has `"status" -> "saved"`; `"saved"` is a status value, not a top-level key. A row `{{values...}, "tmp"}` gives a transient waypoint, represented by `"status" -> "transient"` and excluded from export. Singular-point classifications are returned under `"singularityClassifications"` by the same backend evaluation run.
- Formula artifacts: `MSWriteFormulaArtifacts[context]` writes `masters.wl`, `recurrence_metadata.wl`, `dlog_de.wl` and `manifest.wl` to `results/madstree_formula/run-UUID/` by default. `MSEvaluatePath` and `MSReconstructEpSeries` automatically write or reuse these analytic files before numerical transport and return their paths under `"formulaArtifacts"`.
- Evaluation export: `MSExportEvaluationData[result, MSOutputDirectory -> ...]` writes `evaluation_data.csv` and `evaluation_data.json` under the caller-chosen output directory (default `results/madstree_evaluation/run-UUID/`), for downstream plotting tools.
- Cache identity: a successful-result cache key includes the request and the SHA-256 identities of the backend adapter, the configured backend's `pyproject.toml` and sorted `flintnde/*.py`, so in-place source fixes invalidate stale cache entries automatically.

The package source directory never receives run artifacts.

4.3 Numerical evaluation

With an initialized context, numerical evaluation proceeds in two steps: boundary generation and transport. The workflow is single-stage: `MSEvaluatePath` partitions the user point sequence into maximal consecutive complex-affine one-variable segments, pulls the d log-form connection back once per segment, and sends all segments, the exact boundary conditions and the user options to FlintNDE in one UTF-8 JSON request. With the default `FlintNDEPathPlanning -> True`, FlintNDE plans the transport nodes inside each segment and evaluates user points by fast multipoint evaluation; with `False` it uses every supplied point as a node in input order, without inserting, removing, or replanning

points. The backend inside the version directory is a synchronized copy of the standalone `FlintNDE` package released in the same repository, whose mechanisms are described below. `MSBoundaryData` and `MSEvaluatePath` default to 200 decimal working digits; an explicit `WorkingPrecision` overrides the default.

Boundary generation.

```
targetRules = {k1 -> 9 I, k2 -> 3 I, q -> 1, a1 -> 1, a2 -> 1};

certificate = MSBoundaryChartCertificate[context, targetRules,
  RankOrder -> All];

boundary = MSBoundaryData[context, targetRules,
  BoundaryScale -> 4, WorkingPrecision -> 200];
```

`MSBoundaryChartCertificate` verifies the nested blow-up chart of Sec. 3.5 (normal crossing, theta fixing, quotient checks) for the chosen strict energy rank; `RankOrder -> All` checks every strict chart. `MSBoundaryData` consumes the certificate and returns the leading Frobenius data at the $K_v \to \infty$ boundary: for every master whose leading weight does not vanish, one branch record carrying the Frobenius exponent, the overall coefficient and the normalized leading vector, together with the anchor rules, the rank order and the certificate itself. If the chart is not certified, it returns `BoundaryChartNotCertified` and does not proceed to numerics. The energies in the example are positive imaginary so that the vertex factors $e^{-\mathrm{i}E\tau}$ damp at $\tau \to -\infty$; the rank order of the boundary is derived from the supplied target point, largest damping energy first, unless `RankOrder` fixes it. The example passes `WorkingPrecision -> 200` explicitly; 200 is also the default, and any explicit positive value overrides it.

Transport.

```
pointSequence = {{k1, k2}, {9 I, 3 I}};
parameterRules = {q -> 1, a1 -> 1, a2 -> 1};

result = MSEvaluatePath[context, pointSequence,
  ParameterRules -> parameterRules,
  BoundaryScale -> 4, RankOrder -> {v1, v2},
  WorkingPrecision -> 200, TransportOrder -> 80,
  ReferenceTransportOrder -> 104,
  TargetRelativeError -> "1e-20",
  MessageLanguage -> "EN"];

result["pointResults"]
```

The second argument is the *point sequence*: its first row lists the movable coordinate symbols, and every further row supplies one numerical value for each of them, in the same

order. The remaining symbols of the target point, here `q`, `a1` and `a2`, are fixed once for the whole run through `ParameterRules`. For several targets, append one row per point,

```
pointSequence = {{k1, k2}, {8 I, 2 I}, {7 I, 4 I}, {6 I, 5 I}};
```

and MadStree groups the points into maximal consecutive complex-affine segments and pulls each group back once, while FlintNDE evaluates every point in one process, including dense points inside a node's convergence disk, without changing the path. For a single-vertex integral family, run the same single-stage entry on the vertex context. The principal transport options and the point tags are summarized in Sec. 4.5. If every d log-form letter is constant on a segment, every pulled-back derivative vanishes; the segment is then a zero connection with no finite poles and is transported normally.

Reading the result. `result` is an association whose `"pointResults"` entry is the list of point records. Every saved ordinary point carries `coordinate`, `value` (the master vector in master order), `status` (equal to `"saved"` for ordinary points) and `userIndex`; transient `"tmp"` points participate only in the path. The current version does not provide leading-order point labels: every tag other than `"tmp"` is rejected. With the default `SingularityMode -> "Automatic"` a segment that meets a supported singularity is crossed by an explicit jump; `"Avoid"` refuses such a segment instead, and with `FlintNDEPathPlanning -> False` the backend is forced into avoid mode, so a user chain that crosses a pole or leaves a local convergence disk fails explicitly. The exported per-point `relativeDifferenceInf` and `targetRelativeErrorMet` are obtained from the embedded truncation certification of Sec. 4.3, not from a second independent transport chain. A route-robust access to a saved value is

```
savedPoints = Select[result["pointResults"], #["status"] === "saved" &];
firstPoint = First[savedPoints];
firstPoint["value"]
```

The numerical backend. FlintNDE is a general solver for first-order matrix differential equations with exact $Q(\mathrm{i})(s)$-coefficient systems; MadStree embeds a synchronized copy and only ever feeds it single-variable pullbacks of the d log-form connection. Three mechanisms are responsible for the current performance:

- *Pole–residue transport.* An ordinary segment is serialized as its pole–residue representation $A(x) = P(x) + \sum_j R_j/(x - p_j)$, with P of arbitrary degree and the residues R_j of coincident letters merged. The backend generates the solution Taylor coefficients by a pole-state recurrence, avoiding the Cauchy sampling and discrete-Fourier coefficient reconstruction of a generic analytic-matrix backend.

- *Embedded truncation certification.* Only one transport chain is run (`certificationMode -> "embedded"`); the primary result is re-evaluated from the coefficient prefix by Horner, and the per-segment truncation difference is returned as a local error estimate, so the reference-chain overhead is removed while the `relativeDifferenceInf` and `targetRelativeErrorMet` fields remain available for the user.

- *Fast multipoint evaluation.* With `FlintNDEPathPlanning -> True`, FlintNDE plans nodes inside each segment; user points covered by one expansion node form an evaluation bucket and are evaluated from the cached vector series. Buckets with at least eight points use the subproduct/remainder tree of fast multipoint evaluation, and small buckets use the iterative algorithm to avoid tree overhead. User points in the same complex-affine group $x(s) = x_0 + s\,v$ share node coefficients whenever they lie in the same convergence disk; different segments never share local coefficients.

Singularity handling is explicit: the default `SingularityMode -> "Automatic"` maps to the backend jump mode, so a segment that meets a supported singularity is crossed by an explicit local-basis jump instead of being refused or detoured; `"Avoid"` refuses such a segment, and `"SingularityJump"` requests a jump unconditionally. Both jump modes are available only when `FlintNDEPathPlanning -> True`; with planning disabled the backend is forced into avoid mode and the user must supply a valid ordinary-point node chain. The branch reached by a jump is fixed by the local basis and by the winding of the path, and the user must check that it is the intended one. Exact Lee–Moser balances, strictly decoupled exponential times, and power-log recurrences are handled, while resonances whose local bases cannot be constructed, and any ramification or general Stokes requirement, fail closed. The internal working bits are $\lceil \mathrm{WP}\log_2 10\rceil + 32$ for `WorkingPrecision` = WP digits (the default WP is 200), and all path geometry (segment projections, match ratios, rotation factors, winding/monodromy) is kept in Arb/Acb precision. If the backend realification writes $k = -\mathrm{i}\kappa$, then $-k^2 = +\kappa^2$; a program token named `ik` then denotes the real variable κ, and `ik^2` is its square, not the imaginary unit times k^2.

4.4 Laurent-series reconstruction

When all vertex time powers share a common analytic regulator ε, e.g. $a_v = 1 + \varepsilon$, the master integrals develop a Laurent series

$$I(\varepsilon) = \sum_{p=n_{\min}}^{m} c_p\,\varepsilon^p + O(\varepsilon^{m+1}), \tag{4.2}$$

and `MSReconstructEpSeries[context, ep, pointTemplate, MaximumEpPower -> m]` reconstructs the coefficients $c_{n_{\min}}, \ldots, c_m$. The user supplies no ε values; the lowest power is not a numerical fit parameter.

This route follows the one used by `AMFlow` [54, 55]. The transport is numerical, while the regulator dependence of the result has to stay analytic, and carrying that analytic dependence through every matrix multiplication of a long path would dominate the cost. Each sample is therefore solved purely numerically at a fixed value of ε, and the coefficients of (4.2) are recovered afterwards from the final numbers of the whole set of runs, which is what keeps a multi-order expansion affordable.

Certification of the lowest power. Before any numerical solve at nonzero ε, MadStree constructs a support certificate from the symbolic boundary conditions and the d log-form DE of the production context. It requires:

1. regulator-independent path coordinates and no negative Laurent power at $\varepsilon = 0$ in any d log-form matrix, pulled-back connection, or residue;

2. finite-Laurent boundary branches, where branches with the same limiting Frobenius exponent and log degree are first combined with their physical weights and leading vectors, and only then are component valuations extracted;

3. at least one provably nonzero coefficient at the lowest power of the combined boundary; a noninteger power, non-Laurent structure, undecidable coefficient, or support entirely above the requested user range fails closed.

For $n_{\min} < 0$ the certificate describes a finite meromorphic Laurent boundary, not an analytic boundary at $\varepsilon = 0$; analyticity is recorded only when $n_{\min} \geq 0$. Along the current finite path an ε-analytic invertible fundamental matrix preserves the valuation of nonzero formal boundary conditions, so the certified $n_{\min}$ is passed directly to the sampling plan. A rank drop of an individual Frobenius recurrence operator at $\varepsilon = 0$ is retained as a resonance diagnostic only; it is not by itself interpreted as a $1/\varepsilon$ pole of the already combined physical boundary.

Sampling, fitting and validation. Given the highest requested power m and the certified lowest power $n_{\min}$, the plan sets

$$P = \max(0, -n_{\min}), \qquad \Delta = m - n_{\min}, \qquad N_{\text{fit}} = \max\left(\left\lceil \tfrac{5}{2}\Delta + P \right\rceil, \Delta + 1 + f\right), \quad (4.3)$$

where $f =$ `EpFitExtraOrder`, default 2, is the number of powers the fit keeps above the user maximum: the second branch of (4.3) would be $\Delta + 1$ if only $c_{n_{\min}}, \ldots, c_m$ were fitted, and the extra f is what makes the interpolation reach past c_m in every run. `SampleCount` may raise N_{fit} further, never below $\Delta + 1$. The plan then chooses the ε scale and precisions

$$\alpha_\varepsilon = \frac{P}{4} + \frac{g}{\Delta + 1}, \qquad \varepsilon_0 = 10^{-\alpha_\varepsilon}, \qquad p_0 = \max(\lceil (N_{\text{fit}} + P)\alpha_\varepsilon \rceil, 30), \quad (4.4)$$

where $g =$ `EpGoalDigits` is the requested decimal accuracy. The working precision is $\max(200, \lceil 2(1 + e)p_0 \rceil)$ with e the extra-precision fraction; the primary transport order is $4p_0$ and the reference chain adds $\lceil \max(50, p_0/5) \rceil$ orders. All N_{fit} production points enter one square Acb Vandermonde system interpolating through $\varepsilon^{n_{\min}+N_{\text{fit}}-1}$; only $c_{n_{\min}}, \ldots, c_m$ are returned, and no least squares or pseudoinverse is used. Independent validation points are placed at a fixed fraction of the production scale, never enter the fit, and the truncated series must pass the relative tolerance 10^{-g}; otherwise the run fails closed.

Incremental expansion and parallelism. The initial fit includes at least two powers above the user maximum (`EpFitExtraOrder -> 2`). If validation fails, each subsequent round adds `EpFitOrderIncrement -> 2` powers, solves only the new production points, and reuses every previous production and validation value, for at most `EpFitMaximumRounds -> 3` rounds; the tolerance is not relaxed and the caches are not discarded. Independent ε tasks run in an outer process pool bounded by `ParallelTaskCount`

`-> 12`, whose effective concurrency is the smaller of the limit and the number of tasks; this is separate from the in-process `ctx.threads` of python-flint. The whole route is run end to end by the third application example in Sec. 5.2.

4.5 Options, validation and common pitfalls

The principal options of the numerical workflow are summarized below.

option	meaning
`FlintNDEPathPlanning`	`True` (default): FlintNDE plans nodes and uses fast multipoint evaluation; `False`: user points as nodes in input order
`SingularityMode`	`"Automatic"` (default), `"Avoid"` or `"SingularityJump"`
`MessageLanguage`	`"EN"` (default) or `"CN"`, case-sensitive
`BoundaryScale`	scale setting the damping energies at the finite boundary anchor (> 1; default 8)
`BoundarySeriesOrder`	boundary-order metadata and regulator recurrence diagnostics; not the numerical Frobenius start order in the current generic route
`RankOrder`	vertex order by damping energy K_v (default `Automatic`: from the first target point)
`WorkingPrecision`	integer working precision in decimal digits, at least 20; default 200
`TransportOrder` / `ReferenceTransportOrder`	primary and reference local-series orders, including the numerical Frobenius start; defaults 48 and 64, with the reference order strictly larger
`TargetRelativeError`	target relative error of the primary/reference check; default `"1e-25"`
`MaximumEpPower`	highest returned regulator power of `MSReconstructEpSeries` (default 0)
`EpGoalDigits`	requested decimal accuracy of the reconstructed regulator coefficients (default 20)
`EpFitExtraOrder` / `EpFitOrderIncrement` / `EpFitMaximumRounds`	internal buffer powers, per-round increment and maximum refit rounds of the regulator fit (defaults 2, 2, 3)
`ParallelTaskCount`	maximum concurrent `ep` backend processes in `MSReconstructEpSeries` (default 12)
`PythonExecutable`	Python command for the FlintNDE adapter (default `Automatic`)
`MSRuntimeDirectory`	temporary-file root (default: `results_temp` in the calling script or notebook directory)

The transport orders and `TargetRelativeError` above apply to `MSEvaluatePath`. For `MSReconstructEpSeries`, these three options are not accepted; the numerical settings are

planned from `EpGoalDigits`. Its `BoundarySeriesOrder` defaults to `Automatic`, whereas `MSEvaluatePath` uses 24.

Saved and transient points are not a transport option: they are tags attached to coordinate rows below the symbol header (bare value lists and {{`values...`},`"tmp"`}). With `SingularityMode -> "Avoid"`, interior singularity crossings are refused and reported with the offending path pairs. The default `"Automatic"` mode allows supported singularity jumps when `FlintNDEPathPlanning -> True`; an explicit `"SingularityJump"` also requires this setting. When planning is disabled, `"Automatic"` uses `"Avoid"`. The user must check the selected multivalued branch. An explicitly requested terminal singular point is handled by local classification and evaluation, rather than by ordinary Taylor transport through the singularity.

Limitations and fail-closed guarantees:

- No loop-momentum integration, no sampled IBP, no Kira or external reduction.
- `dlogStatus` must be `"certifiedByFormulaChecks"`; otherwise the DE is not consumed.
- Uncertified boundary charts and singular points beyond the backend's certified capabilities all fail closed; resonances whose local bases cannot be constructed return structured refusals, and ordinary Taylor transport is never used across singular points.
- An incomplete `MSReduce` call may return an association with `"status" -> "partiallyReduced"` or `"status" -> "failed"`, rather than a top-level `Failure`. Inspect `"remainingShiftedIntegrals"` and `"nonMasterResidual"`.
- For `MSEvaluatePath`, check `result["flintNDE"]["targetRelativeErrorMet"]`, not only `result["status"]`. For `MSReconstructEpSeries`, check `"precisionTargetMet"`; coefficients may be returned with `"status" -> "computed_with_warning"` if the target is not met.
- The production boundary never uses defining integrals; finite-point defining integrals exist only inside independent validation programs.

Finally, we collect the most common mistakes.

Common pitfalls.

1. **Convention**: `NuConvention` is chosen at initialization only (`"Positive"` by default). To compare with the formulas of [39], initialize with `NuConvention -> "Negative"`, which selects the minus sign in (4.1).
2. **Sector keys**: fixed-length strings; leading zeros and the string type are part of the identity (Sec. 3.3).
3. **Single-stage workflow**: call `MSEvaluatePath` directly. `MadStree` in this version does not provide `MSGeneratePath`, `MSEvaluatePlannedPath`, public plan objects, old JSON schemas or any migration wrapper; the default `WorkingPrecision` is 200 and an explicit value overrides it.

4. **Singularity handling**: the default is `SingularityMode -> "Automatic"`, where `"Automatic"` is a string, not the symbol `Automatic`. Use `"Avoid"` to refuse interior crossings. Supported jumps require `FlintNDEPathPlanning -> True`, and the user must check the multivalued branch.

5. **Point tags**: after the coordinate-symbol header, only bare value lists and {{`values...`},`"tmp"`} are accepted; every other tag, including `"lo"`, is rejected.

6. **Backend environment**: MadStree uses zero probing; the interpreter is given by the `PythonExecutable` option or, when `Automatic`, by the `MADSTREE_PYTHON` environment variable (falling back to `"python"`), and it requires Python 3.10 or later with `python-flint>=0.6` and `sympy>=1.12`, and must be able to import the embedded `flintnde`. Backend failures report captured output through messages such as `backendLaunchFailed` and `backendRunFailed`; retained runtime logs contain the details. Successful calls and handled singular-path refusals remove their temporary input and log files.

7. **Message language**: `MessageLanguage` accepts only `"EN"` and `"CN"` (case-sensitive), with `"EN"` the default.

8. **Encoding**: loading through `Needs` uses explicit UTF-8 internally and needs no user encoding option; if loading the kernel file directly fails with syntax errors, set `$CharacterEncoding = "UTF-8"` before loading.

5 Examples

This section collects what ships with the release and what has been computed with it. Sec. 5.1 lists the seven examples distributed with the package, which exercise recurrence reduction, single-vertex families, time-order charts, massless and massive trees, zero external-leg energy, and the regulator reconstruction of Sec. 4.4. Sec. 5.2 presents three numerical applications: the tree with three vertices and two massive exchanges, whose transported results are compared with the multivariable hypergeometric representation of [25]; the bubble time kernel evaluated at fixed internal momenta; and the massless three-vertex chain evaluated in the regulator mode, which demonstrates the certified Laurent reconstruction of Sec. 4.4 on a resonant configuration. The comparisons standing behind the independent validation tasks recorded in Sec. 2, namely the single-vertex families and the massive two-vertex correlator against [39] and the IBP relations and differential equations against the `dSIBP` package [59], are not reproduced here.

5.1 Package examples

The distribution ships seven examples that exercise the complete workflow:

- `01_massless_full_edge.wl`: the massless shared two-state simplification of Sec. 3.2, masters, recurrence, d log-form DE and the automatic boundary/numerical entry; it also demonstrates the single-stage multipoint workflow, with CSV/JSON export;

- `02_single_vertex_family.wl`: single-vertex integral family input, local tensor inverses (Sec. 3.2), and finite-combination reduction;
- `03_time_only_cycle_chart.wl`: time-order cycle input, common-theta contacts (Sec. 3.6 and App. A), and all strict-rank chart certificates; contacts are discussed in Sec. 3.3;
- `04_three_vertex_tree.wl`: a three-vertex, two-propagator massless tree ($+++$ vertex structure), following the same flow as example 01, with a single-stage multipoint evaluation and CSV/JSON export;
- `05_massive_three_vertex_tree.wl`: a three-vertex, two-propagator massive tree ($+++$ vertex structure) with non-half-integer `nu12=3/4`, `nu23=1/3`, followed by a single-stage multipoint evaluation and CSV/JSON export;
- `06_massless_three_vertex_ep_regularization.wl`: a three-vertex massless tree with the common analytic regulator $a_1 = a_2 = a_3 = 1 + \varepsilon$; the user requests only the highest regulator power (`MaximumEpPower -> 0`), and the program certifies the lowest power from the symbolic boundary and DE before choosing production/validation points and extracting the pole and finite part; its numerical reconstruction is presented as the third application in Sec. 5.2;
- `07_zero_external_leg_energy.wl`: zero external-leg energy, represented through an auxiliary energy with zero target value.

In the massless three-vertex example (Example 06) with $a_1 = a_2 = a_3 = 1 + \varepsilon$, the physical boundary weights contain $\Gamma(2+\varepsilon)^3$, $\Gamma(2+\varepsilon)\Gamma(4+2\varepsilon)$ and $\Gamma(6+3\varepsilon)$, all regular at $\varepsilon = 0$, and the nine-dimensional DE has no negative ε power; the certificate therefore gives $n_{\min} = 0$ before any numerical solve, and no artificial $1/\varepsilon$ term is introduced.

Each example ends with a failure gate: it prints `Example PASSED` and exits 0 only when every checked result is a non-`Failure` object, and exits 1 with the reason otherwise. Run them in a Mathematica front end section by section, or with

```
wolframscript -file .../Examples/01_massless_full_edge.wl
```

5.2 Application examples

We illustrate the MadStree v1.0 workflow with two examples, using FlintNDE v1.0 for numerical transport. All timings measure elapsed time on a personal computer.

Tree with three vertices and two massive scalars exchanging The first example is the chain with three vertices in figure 2.




Figure 2. Tree with three vertices and two massive scalar propagators. The external energy sums are E_1, E_2, E_3. The internal propagators carry momentum magnitudes s_1, s_2 and mass labels $\widetilde{\nu}_1, \widetilde{\nu}_2$.

The vertices L, M, R have conformal times τ_L, τ_M, τ_R. The two internal propagators connect L to M and M to R, respectively. Each E_i is the sum of the external momentum magnitudes at vertex i. The internal momentum magnitudes are s_1, s_2, and the Hankel orders are ν_1, ν_2. For the masses in the principal series used here, $\nu_i = \mathrm{i}\widetilde{\nu}_i$, with real $\widetilde{\nu}_i$. Each vertex has a contour sign $\sigma_v = \pm 1$.

We first define the states at the two vertices of a propagator that label the master integrals. This application is evaluated in the `NuConvention -> "Negative"` option, i.e. with $s_\nu = -1$ in the unified definition of Sec. 3.1, because the mode functions that this application starts from follow [39], where they are written in that option:

$$\begin{aligned} h^{(r)}(\nu, 0; z) &= z^{-\nu} H_\nu^{(r)}(z), \\ h^{(r)}(\nu, 1; z) &= \partial_z h^{(r)}(\nu, 0; z) = -z^{-\nu} H_{\nu+1}^{(r)}(z), \qquad r = 1, 2. \end{aligned} \tag{5.1}$$

Nothing here is a third convention: the package default is $s_\nu = +1$, with $h^{(r)}(\nu, 0; z) = z^{+\nu} H_\nu^{(r)}(z)$ and $h^{(r)}(\nu, 1; z) = z^{+\nu} H_{\nu-1}^{(r)}(z)$, and the two readings are related by the single simultaneous replacement $\nu \to -\nu$ of Sec. 3.1, which must also be applied to the effective time powers below. Here r selects the Hankel function and the binary index selects the undifferentiated or differentiated state. We use the same Hankel order at both vertices of each propagator; constant factors from the physical mode normalisation are restored when forming the correlator. To display the time ordering explicitly, define

$$\begin{aligned} W_{>,\nu;ab}(s; \tau, \tau') &= h^{(1)}(\nu, a; -s\tau) h^{(2)}(\nu, b; -s\tau'), \\ W_{<,\nu;ab}(s; \tau, \tau') &= h^{(2)}(\nu, a; -s\tau) h^{(1)}(\nu, b; -s\tau'), \qquad a, b \in \{0, 1\}. \end{aligned} \tag{5.2}$$

These two products are $G_>$ and $G_<$ of (3.2) with the mode prefactors removed and the two states made explicit. The propagator kernel of contour type $\sigma\sigma'$, written $\mathcal{G}_{\sigma\sigma',\nu;ab}(s; \tau, \tau')$ below, is then the time-ordered combination of the two products exactly as (3.3) combines $G_>$ and $G_<$: $\mathcal{G}_{++}$ and $\mathcal{G}_{--}$ take $\theta(\tau - \tau') W_>$ plus $\theta(\tau' - \tau) W_<$ and $\theta(\tau - \tau') W_<$ plus $\theta(\tau' - \tau) W_>$, while $\mathcal{G}_{+-} = W_{<,\nu;ab}$ and $\mathcal{G}_{-+} = W_{>,\nu;ab}$ carry no theta at all. The binary indices act only on the Hankel factors, never on the step functions, and every kernel has

the arguments $(s;\tau,\tau')$.

For $\boldsymbol{\sigma}=(\sigma_L,\sigma_M,\sigma_R)$, the unpinched master integrals are

$$\begin{aligned}\mathrm{I}^{\boldsymbol{\sigma}}_{abcd} = \int_{-\infty}^{0} d\tau_L\, d\tau_M\, d\tau_R\, (-\tau_L)^{\nu_{0L}}(-\tau_M)^{\nu_{0M}}(-\tau_R)^{\nu_{0R}} \\ \times e^{\mathrm{i}(\sigma_L E_1\tau_L+\sigma_M E_2\tau_M+\sigma_R E_3\tau_R)} \\ \times \mathcal{G}_{\sigma_L\sigma_M,\nu_1;ab}(s_1;\tau_L,\tau_M)\mathcal{G}_{\sigma_M\sigma_R,\nu_2;cd}(s_2;\tau_M,\tau_R).\end{aligned} \tag{5.3}$$

Here $a,b,c,d\in\{0,1\}$. Each time integral runs from $-\infty$ to 0 with the Bunch–Davies prescription. The four indices refer to the states at the two vertices of each propagator (L_1,M_1,M_2,R_2), in that order. Thus the Top vector is

$$\mathrm{I}^{\boldsymbol{\sigma}}_{\mathrm{Top}} = \big(\mathrm{I}^{\boldsymbol{\sigma}}_{0000},\mathrm{I}^{\boldsymbol{\sigma}}_{0001},\dots,\mathrm{I}^{\boldsymbol{\sigma}}_{1111}\big)^{\mathsf{T}}. \tag{5.4}$$

It contains $2^4=16$ masters. The entry $\mathrm{I}^{\boldsymbol{\sigma}}_{0000}$ is the one with no derivative acting on the states and no additional integer shifts in the time powers. Equation (5.3) is the sector $\mathrm{Top}=(\xi_1\xi_2)=(11)$ of (3.14), with $c_{\mathrm{Top}}=c_V=3$ components by (3.15) and $n^{\mathrm{slot}}=4$ slots, and

$$\mathrm{I}^{\boldsymbol{\sigma}}_{abcd} = \mathrm{I}_{\mathrm{Top}}\big(\mathbf{0};(a,b,c,d)\big), \tag{5.5}$$

the bit order of (3.12) being the one used by $\mathrm{I}^{\boldsymbol{\sigma}}_{\mathrm{Top}}$.

The parameters $\nu_{0L},\nu_{0M},\nu_{0R}$ in the integrand are the effective vertex time powers. If p_1,p_2,p_3 are the powers before absorbing the massive mode prefactors, then

$$\begin{aligned}\nu_{0L} &= p_1+\frac{3}{2}+\nu_1, \qquad \nu_{0R} = p_3+\frac{3}{2}+\nu_2,\\ \nu_{0M} &= p_2+3+\nu_1+\nu_2.\end{aligned} \tag{5.6}$$

The vertex factors $\mathrm{i}\sigma_v$ and the external and mode normalisations are not included in (5.3); they are restored when constructing the physical correlator.

In the IBP reduction, derivatives of the functions that impose time ordering generate contact terms that merge adjacent vertices. These terms define the LeftPinch, RightPinch and DoublePinch sectors. The left pinch merges L and M and leaves the two indices of propagator 2, so its four masters are the child integrals of the contact term (3.48), written out as

$$\begin{aligned}\mathrm{I}^{\boldsymbol{\sigma},\mathrm{L}}_{cd} = \int_{-\infty}^{0}\mathrm{d}\tau_{LM}\int_{-\infty}^{0}\mathrm{d}\tau_R\,(-\tau_{LM})^{\nu_{0L}+\nu_{0M}-2\nu_1}(-\tau_R)^{\nu_{0R}} \\ \times e^{\mathrm{i}(\sigma_L E_1+\sigma_M E_2)\tau_{LM}+\mathrm{i}\sigma_R E_3\tau_R}\,\mathcal{G}_{\sigma_M\sigma_R,\nu_2;cd}(s_2;\tau_{LM},\tau_R), \qquad c,d\in\{0,1\},\end{aligned} \tag{5.7}$$

where τ_{LM} is the single time of the merged component and the merged power is $A_{C,\mathrm{L}}$ of (3.40): the raw contact power of the contracted propagator is $\varrho_1=-2\nu_1-1$ in the negative-prefactor convention of (5.1), and the component count $|C|=2$ adds the remaining $+1$, so that (5.6) gives $\nu_{0L}+\nu_{0M}-2\nu_1=p_1+p_2+\frac{9}{2}+\nu_2$. The right pinch, with masters $\mathrm{I}^{\boldsymbol{\sigma},\mathrm{R}}_{ab}$, merges M and R and keeps the two indices of propagator 1; it is obtained from (5.7) by the

exchange $(L, \nu_{0L}, \sigma_L E_1, s_1, \nu_1) \leftrightarrow (R, \nu_{0R}, \sigma_R E_3, s_2, \nu_2)$. The double pinch merges all three vertices and leaves no massive propagator, so its single master $\mathrm{I}^{\boldsymbol{\sigma},\mathrm{D}}$ is the one-time integral of $(-\tau)^{\nu_{0L}+\nu_{0M}+\nu_{0R}-2\nu_1-2\nu_2} e^{\mathrm{i}(\sigma_L E_1+\sigma_M E_2+\sigma_R E_3)\tau}$, whose power is $p_1+p_2+p_3+6$. In the index convention (3.13) the four sectors are Top $= (\xi_1\xi_2) = 11$ with three components and four slots, (LeftPinch, RightPinch) $= (01, 10)$ with two components and two slots each, and DoublePinch $= 00$ with one component and no slot, exactly as (3.15) requires of a graph with $c_V = 3$ vertices. When both internal propagators join vertices with equal contour signs, the full hierarchy is

$$\mathrm{Top}(16) \oplus \mathrm{LeftPinch}(4) \oplus \mathrm{RightPinch}(4) \oplus \mathrm{DoublePinch}(1), \qquad \dim = 25. \tag{5.8}$$

For other contour assignments, only the allowed pinched sectors are included: a propagator whose two vertices carry opposite signs is theta-free, so contracting it is not an available event.

Writing σ_{LM} and σ_{MR} for the contour signs of the two internal propagators, $+1$ for G_{++} and -1 for G_{--} as in (3.23), the lowering contact term of the Top IBP at L or M is the two-vertex relation (3.48) in this graph's indices,

$$\mathcal{R}^{(0)}_{\tilde{\boldsymbol{r}};\sigma_{LM}} = -\sigma_{LM}\,\delta_{a,1-b}\,(-1)^{a+1}\,\frac{4\mathrm{i}}{\pi}\,s_1^{-2\nu_1-1}\,\mathrm{I}_{\mathrm{L}}\big(-\boldsymbol{e}_{LM};(c,d)\big), \qquad \tilde{\boldsymbol{r}} = 1+2c+d, \tag{5.9}$$

where the child is taken at shift -1 on the merged component and 0 on τ_R, so that its merged power is one below (5.7), and the bits a, b of the contracted propagator are removed from the parent string, leaving $\tilde{\boldsymbol{r}}$ as the packed label of (c, d). The four sectors then carry the hidden constants

$$\begin{aligned} \mathcal{N}_{\mathrm{Top}} &= 1, & \mathcal{N}_{\mathrm{L}} &= -\sigma_{LM}\,\frac{4\mathrm{i}}{\pi}\,s_1^{-2\nu_1-1}, \\ \mathcal{N}_{\mathrm{R}} &= -\sigma_{MR}\,\frac{4\mathrm{i}}{\pi}\,s_2^{-2\nu_2-1}, & \mathcal{N}_{\mathrm{D}} &= -\sigma_{LM}\sigma_{MR}\,\frac{16}{\pi^2}\,s_1^{-2\nu_1-1}s_2^{-2\nu_2-1}, \end{aligned} \tag{5.10}$$

which is the product rule (3.16) with $\mathcal{N}_s^{(0)} = 1$, read in the negative-prefactor convention, where $q_e^{2\nu_e-1}$ becomes $q_e^{-2\nu_e-1}$. The object the package propagates in sector s is $J_s(\boldsymbol{n};\boldsymbol{a}) = \mathcal{N}_s\,\mathrm{I}_s(\boldsymbol{n};\boldsymbol{a})$, and for the $(+\,+\,+)$ assignment (5.10) is what the `"normalization"` field of the master records returns.

At real kinematic points, we compute the four assignments $(+\,+\,+)$, $(+\,+\,-)$, $(+\,-\,+)$ and $(-\,+\,+)$. After restoring the vertex factors and mode normalisations, each $\mathrm{I}^{\boldsymbol{\sigma}}_{0000}$ entry gives one contribution to the physical correlator. Reversing all three contour signs gives the complex conjugate contribution. The four computed terms and their conjugates determine $\mathcal{A}_{\mathrm{phys}}$. To match the phase convention above, the application code passes $-E_v$ as `externalLegEnergy`, uses `NuConvention -> "Negative"`, and supplies the corresponding pinch normalisations.

We set $p_1 = p_2 = p_3 = 0$ and express all energies and momenta in units of E_2. With

$q = s_1/E_2$, the scan follows

$$\begin{aligned}(E_1, E_2, E_3, s_1, s_2) &= \left(\frac{100q}{99}, 1, \frac{10}{99}, q, \frac{1}{10}\right), \qquad (\nu_1, \nu_2) = (\mathrm{i}, 2\mathrm{i}),\\ (\nu_{0L}, \nu_{0M}, \nu_{0R}) &= \left(\frac{3}{2} + \mathrm{i}, 3 + 3\mathrm{i}, \frac{3}{2} + 2\mathrm{i}\right).\end{aligned} \tag{5.11}$$

As the numerical benchmark, which is independent of the DE construction, we use the multivariable hypergeometric representation of this correlator derived in ref. [25]. It expresses the signal and background contributions as multiple series containing Appell F_2 functions. Their sum gives the complete physical correlator. These contributions are not the IBP sectors in (5.8). For this benchmark, DE transport evolves the full master vector for each contour assignment and then extracts the physical correlator from its $\mathrm{I}^{\boldsymbol{\sigma}}_{0000}$ entries. The hypergeometric calculation evaluates the same physical correlator directly at each target point; it does not separately evaluate all components of the master vectors. The reported comparison is therefore between physical correlators, not a componentwise comparison of all masters. The truncation order is selected by comparing adjacent orders. The reported hypergeometric time is the sum of the accepted pointwise evaluation times.

The DE calculation uses 80 decimal digits and local Taylor series truncated at order 80. An independent transport at order 96 provides an internal stability check. We compare batches of $N = 1$, 10, 50 and 100 target points on the slice in (5.11). For $N = 1$, 10 and 50, transport starts at $q = 0.1$ and ends at $q = 10^{-4}$. The single point is the end of the path, corresponding to $x = -\log_{10} q = 4$. The ten points are selected at approximately equal index intervals from the original 50 points, including both ends. The 100 point batch contains 80 points on this path and 20 additional points from $q = 0.105$ to 0.2. The additional points share the boundary at $q = 0.1$ but require transport in the opposite direction.

The hypergeometric times for these four batches are 22.78, 344.62, 1040.95 and 4242.89 s. The corresponding median DE times are 26.71, 48.82, 48.86 and 57.56 s (figure 3). Each DE time includes physical boundary construction, DE generation and the transport call, with the order 96 backend time subtracted. It includes data preparation and Python invocation. All four DE batches were measured three times, with the execution order varied between rounds. We report the median total time and show the full range of the three measurements. The observed ratios of hypergeometric time to DE time are 7.06, 21.31 and 73.71 for ten, 50 and 100 points, respectively. The isolated end of the path is slower with DE transport than with the hypergeometric representation.

The average DE time per output point decreases from 26.71 s to 4.88, 0.98 and 0.58 s across these batches. Boundary data and transport are shared, rather than recomputed independently for each target. Points encountered along the path use saved node values or evaluations of an already constructed local series. In the new tests, each branch used 29 internal nodes for one output point and 33 nodes for ten output points. Thus, additional outputs did not require a proportional increase in transport work.

Within a convergence disk, a shorter step suppresses higher Taylor terms. It can

therefore permit a lower truncation order at a fixed accuracy. This is a local property, not a guarantee of lower total cost, since shorter steps can require more segments.




Figure 3. Tree benchmark with MadStree v1.0 and FlintNDE v1.0. **a**, Hypergeometric evaluation time at individual kinematic points. **b**, Total evaluation times for $N = 1$, 10, 50 and 100 target points.

The 100 points in figure 3**a** comprise 80 points with $q = 10^{-x}$ and $x = 1 + 3j/79$ ($j = 0, \ldots, 79$), uniformly spaced in x, and 20 points with $q = 0.1 + 0.005j$ ($j = 1, \ldots, 20$), uniformly spaced in q. Each hypergeometric time uses the accepted truncation order. In panel **b**, the DE bars show medians of three runs, with error bars spanning the minimum and maximum. The times include boundary construction, DE generation and the transport call. The first three batches share the same start and end; the 100 point batch includes the additional path extension. Hypergeometric data are newly computed for one and ten points and retained for 50 and 100 points. The ratio labels give the hypergeometric time divided by the DE time. The figure footer reports the largest relative differences between transport orders and from the hypergeometric results across all four batches.

Across the scan of 100 points, the largest relative difference from the hypergeometric result is 1.712×10^{-32}. The comparison of transport orders 80 and 96 gives a maximum relative difference of 1.308×10^{-50}. At $x = 4$, the physical correlator is

$$\mathcal{A}_{\text{phys}}(4) = 9.87033514798954 \times 10^{-9}, \tag{5.12}$$

with a relative difference of 4.21×10^{-36} from the hypergeometric result.

On the same kinematic configuration, the direct Appell F_2 representation requires $q < 0.89$. The point $q = 0.8905$ lies outside this domain, whereas the nearest positive pole of the DE connection is at $q = 0.9$. To reach $q = 1$, we continue the chosen analytic branch

along

$$0.1 \to 0.86 \to 0.86 + 0.04\mathrm{i} \to 0.93 + 0.04\mathrm{i} \to 0.93 \to 1. \tag{5.13}$$

This contour avoids the pole through the upper half of the complex plane. The continuation records 81 points. The calculation at order 80 takes 35.08 s, or 45.76 s when the reference calculation at order 96 is included. The maximum relative difference between orders 80 and 96 is 1.802×10^{-44}. Within the overlap region, the maximum relative difference from the hypergeometric result is 1.712×10^{-32}. The DE method thus evaluates the selected analytic branch beyond the domain of the direct series representation.




Figure 4. Analytic continuation with MadStree v1.0. **a**, Dependence of the hypergeometric partial sums on the truncation order near the direct convergence boundary. **b**, DE continuation at order 80 along the contour in (5.13).

Massless three-vertex tree in the regulator mode The third example returns to the chain topology of the first example with both exchanges massless, and shifts all vertex time powers by the common analytic regulator (figure 5). At the unshifted powers $a_v = 1$ the boundary branch exponents of this graph resonate, which is the situation the regulator mode of Sec. 4.4 is made for: the computation is defined at $a_v = 1 + \varepsilon$ and the physical answer is the Laurent limit $\varepsilon \to 0$. The momenta are fixed by `ParameterRules` and the target point is $\{k_1, k_2, k_3\} = \{9\mathrm{i}, 3\mathrm{i}, 5\mathrm{i}\}$:




Figure 5. Massless three-vertex chain in the regulator mode. Both internal propagators are massless with momentum magnitudes q_{12}, q_{23}; each vertex v carries external energy k_v and the regularized time power $(-\tau_v)^{1+\varepsilon}$.

```
treeSpec = <|
  "vertices" -> {
    <|"id" -> 1, "externalLegEnergy" -> k1,
      "timePower" -> a1, "vertexType" -> "+"|>,
    <|"id" -> 2, "externalLegEnergy" -> k2,
      "timePower" -> a2, "vertexType" -> "+"|>,
    <|"id" -> 3, "externalLegEnergy" -> k3,
      "timePower" -> a3, "vertexType" -> "+"|>},
  "lines" -> {
    <|"type" -> "massless", "endpoints" -> {1, 2},
      "momentum" -> q12, "nu" -> 1/2|>,
    <|"type" -> "massless", "endpoints" -> {2, 3},
      "momentum" -> q23, "nu" -> 1/2|>}|>;
context = MSInitTree[treeSpec];
epSeries = MSReconstructEpSeries[context, ep,
  {{k1, k2, k3}, {9 I, 3 I, 5 I}},
  ParameterRules -> {q12 -> 1, q23 -> 2,
    a1 -> 1 + ep, a2 -> 1 + ep, a3 -> 1 + ep},
  MaximumEpPower -> 0, EpGoalDigits -> 12,
  EpSampleAngleRange -> {-Pi/3, Pi/3},
  RankOrder -> {1, 2, 3}]
```

The shipped example additionally sets `BoundaryScale -> 4`, `BoundarySeriesOrder -> 24` and the default pool of `ParallelTaskCount -> 12` workers. Each massless full propagator contributes one shared two-state slot (Sec. 3.2), so the sector hierarchy of this graph is $\mathrm{Top}(4) \oplus \mathrm{LeftPinch}(2) \oplus \mathrm{RightPinch}(2) \oplus \mathrm{DoublePinch}(1)$ with $\dim = 9$, the nine-dimensional DE recorded in Sec. 5.1; the massive chain of (5.8) carries four state slots and 25 masters. No contracted line contributes a pinch factor in (3.16), so every sector normalisation is 1 and the propagated vector is the master vector I itself, in the convention of (5.3) without vertex contour factors.

The certificate of Sec. 4.4 runs before any numerical solve and returns $n_{\min} = 0$: the three Γ combinations of Sec. 5.1 are regular at $\varepsilon = 0$ and the d log-form DE has no negative ε power. Three Frobenius recurrence operators do drop rank at $\varepsilon = 0$, with rank changes $9 \to 5$, $9 \to 8$ and $9 \to 8$ at the exponents $6(2+\varepsilon)$ and $7(2+\varepsilon)$; per Sec. 4.4 these are retained as resonance diagnostics and produce no pole, since the combined physical boundary decides the support. With $m = 0$ and $g = 12$, (4.3) and (4.4) give $N_{\text{fit}} = 3$, $\alpha_\varepsilon = 12$, $\varepsilon_0 = 10^{-12}$, $p_0 = 36$, a working precision of 200 digits and a primary transport order of 144 with 50 reference orders. The three production points are exact Gaussian rationals of moduli $\varepsilon_0(1 + j/100)$, $j = 1, 2, 3$, one on each of the rays $-\pi/6$, 0 and $+\pi/6$ of the open window $(-\pi/3, \pi/3)$; the two validation points sit at half that modulus on the first two rays and never enter the fit. All five solves are dispatched to one bounded pool (12 requested, 5 effective); the whole reconstruction, including the certificate, both validation solves and the fit, takes 257 s of wall time.

The returned pole coefficients are empty, as certified, and the ε^0 coefficient is the 9-entry master vector. Two representative entries are

$$\begin{aligned} [c_0]_{\text{Top},00} &= \left(1.70880370598 - 3.02691315048\,\mathrm{i}\right) \times 10^{-5}, \\ [c_0]_{\text{D}} &= 4.97150313687 \times 10^{-6} + 5.79 \times 10^{-55}\,\mathrm{i}, \end{aligned} \tag{5.14}$$

the first entry of (5.14) being the top-sector entry with both shared slots undifferentiated and the second the double-pinch master, a single one-time integral. At the two independent validation points the truncated series reproduces the directly transported values with a maximum relative residual of 3.86×10^{-36}, four orders above the requested tolerance 10^{-12}; the coefficients themselves are returned with about 199 significant digits. This example shows the regulator route end to end on a resonant configuration: the lowest power is certified symbolically before any numerical solve, the sampling grid and precisions follow from the certified powers, and the $\varepsilon \to 0$ limit is delivered together with an explicit validation residual.

Bubble time kernel at fixed loop momentum The second example has two time vertices joined by two massive propagators (figure 6). We evaluate the integrals over the two vertex times while holding the internal momenta fixed. We do not integrate over the spatial loop momentum.




Figure 6. Bubble time kernel with external energy sums P_1, P_2 and fixed internal momentum magnitudes x_1, x_2.

Let τ_1, τ_2 be the vertex times, P_1, P_2 the external energy sums, and x_1, x_2 the internal momentum magnitudes. The Hankel orders of the two internal propagators are ν_{B1}, ν_{B2}. With the products $\mathcal{G}_{\sigma\sigma',\nu;ab}$ of (5.2) and (3.3), the Top masters are

$$\begin{aligned} \mathrm{I}^{\sigma_1\sigma_2}_{abcd} = & \int_{-\infty}^{0} d\tau_1\, d\tau_2\, (-\tau_1)^{p_1} (-\tau_2)^{p_2} e^{\mathrm{i}(\sigma_1 P_1 \tau_1 + \sigma_2 P_2 \tau_2)} \\ & \times \mathcal{G}_{\sigma_1\sigma_2,\nu_{B1};ab}(x_1; \tau_1, \tau_2) \mathcal{G}_{\sigma_1\sigma_2,\nu_{B2};cd}(x_2; \tau_1, \tau_2). \end{aligned} \tag{5.15}$$

Here $a, b, c, d \in \{0, 1\}$. The indices a, b label the two states that propagator 1 contributes at vertices 1 and 2; c, d label those of propagator 2 in the same vertex order. Here p_1, p_2 are the effective time powers after absorbing the mode prefactors, unlike the bare powers used in (5.6). The contour signs are $\sigma_1, \sigma_2 = \pm 1$. We denote positive and negative signs by p and m, respectively; these labels are unrelated to the time powers p_1, p_2. The four contour assignments are pp, pm, mp and mm. In MadStree v1.0, assignments with equal contour signs have 22 masters after the exact state projection. Assignments with opposite contour signs have only the Top sector, with 16 masters. For each assignment, the projection onto the Top sector is $\mathrm{I}^{\sigma_1\sigma_2}_{\mathrm{Top}} = (\mathrm{I}^{\sigma_1\sigma_2}_{0000}, \mathrm{I}^{\sigma_1\sigma_2}_{0001}, \ldots, \mathrm{I}^{\sigma_1\sigma_2}_{1111})^{\mathsf{T}}$. The vertex factors $(\mathrm{i}\sigma_1)(\mathrm{i}\sigma_2)$ give

$$\boldsymbol{K}_{\mathrm{SK}} = -\mathrm{I}^{pp}_{\mathrm{Top}} + \mathrm{I}^{pm}_{\mathrm{Top}} + \mathrm{I}^{mp}_{\mathrm{Top}} - \mathrm{I}^{mm}_{\mathrm{Top}}. \tag{5.16}$$

The two pinched sectors of an equal-sign assignment have three masters each, and the hidden constants (3.16) of the three sectors are

$$\mathcal{N}_{\mathrm{Top}} = 1, \qquad \mathcal{N}_1 = -\sigma_1\sigma_2 \frac{4\mathrm{i}}{\pi} x_1^{2\nu_{B1}-1}, \qquad \mathcal{N}_2 = -\sigma_1\sigma_2 \frac{4\mathrm{i}}{\pi} x_2^{2\nu_{B2}-1}, \tag{5.17}$$

with $\mathcal{N}_s^{(0)} = 1$; this example runs in the package default convention, so the power is the $2\nu_e - 1$ of (3.16) rather than the $-2\nu_e - 1$ of the tree example. We use $k_s = x_1$ as the momentum unit and vary the dimensionless energy $\rho = P_1/k_s = P_2/k_s$. The parameter

values in (5.15) are

$$\begin{aligned}(k_s, x_1, x_2) &= (1, 1, \sqrt{2}), \qquad P_1 = P_2 = \rho, \\ (\nu_{B1}, \nu_{B2}) &= (\mathrm{i}/4, \mathrm{i}/4), \qquad p_1 = p_2 = 1 + \mathrm{i}/2.\end{aligned} \tag{5.18}$$

The normalised 0000 component of $\boldsymbol{K}_{\mathrm{SK}}$, written $\widehat{K}_{\mathrm{SK}}(\rho)$, is

$$\widehat{K}_{\mathrm{SK}}(\rho) = \frac{[\boldsymbol{K}_{\mathrm{SK}}(\rho)]_{0000}}{[\boldsymbol{K}_{\mathrm{SK}}(10)]_{0000}}. \tag{5.19}$$

The four zeros select the four undifferentiated states of the two massive propagators.

The scan contains 201 points over $2.8 \leq \rho \leq 10$. Calculations use decimal arithmetic with 80 digits. The transport orders are 80 and 96, and the boundary series orders are 17 and 15. The calculation time excluding reference transport at order 96 is 678.20 s. It includes all four contour assignments, DE construction and both boundary calculations. The reference transport takes 5.24 s, and the total recorded time is 683.43 s. The average cost for the scan, including setup, is 3.37 s per point. This average is not the cost of a calculation at a single point.

The maximum relative difference between boundary orders 17 and 15 is 4.740×10^{-24}. The maximum relative difference between transport orders 80 and 96 is 1.025×10^{-50}. These tests assess boundary and transport stability separately; the comparison of transport orders alone is not a bound on the total error. The largest relative difference from the independent scan is 3.70×10^{-14}.




Figure 7. Bubble time kernel evaluated with MadStree v1.0. **a**, The normalised kernel and the independent reference. **b**, The kernel in the complex plane, with colour indicating ρ.

Both panels in figure 7 use the same 201 points. The nearly radial distribution in panel **b** indicates an approximately constant phase. The maximum relative difference between

transport orders 80 and 96 is 1.025×10^{-50}. The average calculation time, including setup, is 3.37 s per point; this is the batch average, not an isolated point timing.

The tree example illustrates two uses of DE transport: efficient multipoint evaluation and continuation beyond the convergence domain of a direct series. The bubble example applies the same framework for time integrals to a graph containing a loop at fixed internal momenta. The regulator example certifies the Laurent support symbolically on a resonant massless chain and delivers the $\varepsilon \to 0$ limit from five numerical solves.

6 Summary and outlook

`MadStree` implements the general IBP and d log-form construction of [39, 57] as a single computation that starts from the diagram the user supplies and ends at kinematic points, with no intermediate formula written by hand. Three local blocks — the theta-free vertex exponential, the massive two-state block, and the massless block whose four states compress into one shared two-state factor — feed one registry, and that registry alone gives the iterative reduction, the uniform d log-form differential equation, and the contact terms that merge vertices under a common theta function. The boundary data come from the same objects: a strict ranking of the vertex energies freezes every theta function to 0 or 1, the factorized time integrals then return the leading data of all masters as products of Gamma functions, one per time component, and a vertex without an external leg is handled by a private auxiliary energy inside the same system rather than by a separate limit. Because neither the reduction nor the boundary needs the analytic form of the answer, one transport covers a whole path. The tree application of Sec. 5.2 shows the cost per output point falling as points are added to a batch, and shows the same machinery continuing a chosen branch beyond the convergence domain of the direct multivariable series representation.

Two decisions in the implementation matter as much as the formulas. Everything that could invalidate a result is settled before the numerics: the certification carried by `dlogStatus`, the boundary chart certificate without which `MSBoundaryData` refuses to proceed, and the support certificate that fixes the lowest regulator power $n_{\min}$ from the symbolic boundary and the DE, so that the $\varepsilon \to 0$ limit is a Laurent reconstruction and not a fit. When a check fails, the run fails closed and returns a `Failure` object with the captured backend output instead of a silently degraded number. The price of that discipline is a short list of obligations that stay with the user: after a jump across a supported singularity the branch reached by the transport has to be identified by the user, and the comparisons between boundary orders and between transport orders reported in Sec. 5.2 are stability checks, not a bound on the total error.

The scope is the one announced in the abstract: the package computes vertex time integrals, for a tree, or for a loop graph with its internal momenta held fixed, and it does not perform the momentum integrations. That is also the direction of the next step. The vertex blocks, the iterative reduction and the uniform d log-form DE derived here are the kernel of the parity-split IBP system and the d log-form DE of loop integrals in de Sitter space constructed in [58], so the algebraic layers have a loop-level counterpart already

written down, while the momentum integrations and the sectors they bring are the part that still has to be built.

A Common-theta bundles and the odd-subset expansion

Bundles and their theta argument. Fix a sector s and write $\mathrm{comp}_s(v)$ for the current component that contains the vertex v; a component is a maximal set of vertices already merged by contacts, and a lone vertex is one component. An uncontracted full propagator $e = (u_e, v_e)$ carries the theta $\theta(\tau_{u_e} - \tau_{v_e})$ of its two component times, so two propagators e, e' carry the *same* theta if and only if their component pairs coincide as ordered pairs,

$$\begin{gathered}\big(\mathrm{comp}_s(u_e), \mathrm{comp}_s(v_e)\big) = \big(\mathrm{comp}_s(u_{e'}), \mathrm{comp}_s(v_{e'})\big) = (U, V), \\ \text{so} \quad \Delta_e = \Delta_{e'} \equiv \Delta = \tau_U - \tau_V .\end{gathered} \tag{A.1}$$

A maximal set of propagators with this property is the common-theta bundle $\mathcal{B}$ of the pair (U, V), with $|\mathcal{B}| = n_{\mathcal{B}} \geq 2$; the case $n_{\mathcal{B}} = 1$ is an ordinary single-propagator contact and is contained in Sec. 3.4. Propagators whose theta arguments differ belong to different bundles, and the bundles are independent because their Δ's are independent variables: the contact events are therefore performed bundle by bundle, never across bundles. The vertex exponentials of Block (i) are theta-free and take no part in this bookkeeping, so only the two-state factors of (3.3) enter the steps below.

Step 1: multiplying the thetas out. For each propagator of one bundle, write the SK pair of products as in (3.3), α_e for the coefficient of $\theta(\Delta)$ and β_e for that of $\theta(-\Delta)$. At the level of the h blocks the two are the two orderings of the Hankel branches,

$$\alpha_e = h^{(1)}_{u_e}\, h^{(2)}_{v_e}, \qquad \beta_e = h^{(2)}_{u_e}\, h^{(1)}_{v_e}, \tag{A.2}$$

where the orders, states and arguments are those of (3.8); G_{--} interchanges the two names, and this is the only place the sign σ_e of the bundle enters. The product of the bundle is a product of $n_{\mathcal{B}}$ binomials in the two functions $\theta(\Delta)$ and $\theta(-\Delta)$. Expanding naively gives $2^{n_{\mathcal{B}}}$ terms, but inside one bundle $\theta(\Delta)$ is the *same* function for every factor, so idempotency $\theta(\Delta)^2 = \theta(\Delta)$ makes all powers of one argument collapse to that argument, and orthogonality $\theta(\Delta)\theta(-\Delta) = 0$ annihilates every term containing both arguments. Exactly the two terms of (3.63) survive. The factor $2^{n_{\mathcal{B}}}$ thus never appears within a bundle: it counts the events of $n_{\mathcal{B}}$ propagators with pairwise different theta arguments, which is the case of $n_{\mathcal{B}}$ different bundles, and those are enumerated bundle by bundle.

Step 2: one theta gives one delta. Differentiating the two-term form with respect to the single variable Δ gives

$$\partial_\Delta \left[\theta(\Delta) \prod_{e\in\mathcal{B}} \alpha_e + \theta(-\Delta) \prod_{e\in\mathcal{B}} \beta_e \right] = \delta(\Delta) \left(\prod_{e\in\mathcal{B}} \alpha_e - \prod_{e\in\mathcal{B}} \beta_e \right), \tag{A.3}$$

because $\partial_\Delta\theta(\Delta) = \delta(\Delta)$ and $\partial_\Delta\theta(-\Delta) = -\delta(\Delta)$. Three consequences are worth stating explicitly. First, the bundle produces *one* distribution, not $n_{\mathcal{B}}$ of them: the whole contact information of the bundle is concentrated in the single coefficient $\prod_e \alpha_e - \prod_e \beta_e$, which is antisymmetric under $\alpha \leftrightarrow \beta$. Second, that one $\delta(\Delta)$ forces the two component times to coincide *once*, so the components U and V merge once, independently of how many factors d_e a term of the coefficient contains; this is what fixes the merge count in (3.40). Third, for G_{--} the interchange of α and β turns (A.3) into its negative, which reproduces the single sign σ_e of (3.27) for a whole bundle.

Step 3: telescoping the antisymmetric difference. Order the propagators of the bundle as $1, \dots, n_{\mathcal{B}}$ and abbreviate $\alpha_e = A_e$, $\beta_e = B_e$. Inserting and subtracting successive partial products gives the telescoping identity

$$\begin{aligned} \prod_{e=1}^{n_{\mathcal{B}}} A_e - \prod_{e=1}^{n_{\mathcal{B}}} B_e &= \sum_{k=1}^{n_{\mathcal{B}}} \left(\prod_{e\le k} A_e \prod_{e>k} B_e - \prod_{e<k} A_e \prod_{e\ge k} B_e \right) \\ &= \sum_{k=1}^{n_{\mathcal{B}}} D_k \prod_{e<k} A_e \prod_{e>k} B_e, \end{aligned} \tag{A.4}$$

whose intermediate sum collapses because the k-th second summand and the $(k{+}1)$-th first summand are identical, while the first summand of $k = 1$ and the last summand of $k = n_{\mathcal{B}}$ supply the two products of (A.3). Each term of (A.4) contains exactly one difference factor $D_e = A_e - B_e$.

Step 4: collecting by subsets and the parity. Introduce, as in (3.64), the difference and the mean of the two products, $\mathsf{d}_e = D_e = A_e - B_e$ and $\mathsf{m}_e = (A_e + B_e)/2$, equivalently $A_e = \mathsf{m}_e + \mathsf{d}_e/2$ and $B_e = \mathsf{m}_e - \mathsf{d}_e/2$, and expand the spectators of (A.4) in this basis. In the k-th term the propagator k supplies the explicit d_k; a propagator $e \in S$ with $e < k$ contributes $+\mathsf{d}_e/2$, a propagator $e \in S$ with $e > k$ contributes $-\mathsf{d}_e/2$, and every propagator outside S contributes m_e. So the coefficient of the monomial $\prod_{e\in S} \mathsf{d}_e \prod_{e\notin S} \mathsf{m}_e$ is

$$c_S = 2^{1-|S|} \sum_{k\in S} (-1)^{\#\{e\in S:\, e>k\}}. \tag{A.5}$$

Writing $S = \{i_1 < \cdots < i_m\}$ with $m = |S|$, the exponent for $k = i_j$ is $m - j$, hence

$$\sum_{j=1}^{m} (-1)^{m-j} = \sum_{r=0}^{m-1} (-1)^r = \begin{cases} 1, & m \text{ odd}, \\ 0, & m \text{ even}, \end{cases} \qquad c_S = \big(1 - (-1)^{|S|}\big) 2^{-|S|}, \tag{A.6}$$

which is (3.64): only odd subsets survive, each with the pure numerical weight $2^{1-|S|}$. The empty set cannot occur, since every telescoping term carries at least the explicit d_k, in agreement with $c_\varnothing = 0$. The same parity follows from symmetry without any enumeration: the left-hand side of (A.4) changes sign under $\alpha \leftrightarrow \beta$, in which d_e is odd and m_e even, so a monomial with an even number of d factors must cancel; the explicit computation (A.5) only fixes the remaining coefficients.

Events, weights and the bookkeeping. Each non-empty odd subset S of one bundle is one simultaneous contact event. Its injection matrix $C_{st}^{(v,1)}$ of (3.49) is the tensor product of the local contact selectors of the propagators in S — $|1\rangle_e$ on a massless shared slot, the complementary-bit selector of (3.48) on a pair of massive-propagator states — multiplied by $2^{1-|S|}$ and by the spectator factors m_e of the unselected bundle propagators, then embedded in the global bit order with all spectator bits of other propagators preserved. The collected form (3.64), not the term-by-term form (A.4), is what the program enumerates: the k-th telescoping term stands for "contract only the k-th propagator," but after the spectators are expanded the same subset S is reached from several k, so an enumeration over k would generate the same sector transition more than once. After collecting, each event appears exactly once with a numerical weight, and the deduplication key of the reduction is the triple (parent sector, child sector, set of selected propagator identifiers), which is finite because every event strictly reduces the number of current components.

Child base power and coincident unselected propagators. For a component C produced by such events the base power is (3.40), $A_{C,s} = \sum_{v\in C} \nu_{0,v} + \sum_{e\in E_s(C)} \varrho_e + |C| - 1$. The three terms have three different origins and should not be conflated: the vertex powers add because the merged integrand is a product, each contracted propagator adds its own raw contact power ϱ_e because the corresponding δ brings one factor $(-\tau)^{\varrho_e}$, and the last term counts *merges of components*, which is $|C| - 1$ for any component of $|C|$ vertices and does not grow with the number of bundle propagators contracted at once. Since (3.40) depends only on the final component and on the contracted set, and not on the order or the clustering of the events, different legal event sequences that reach the same sector give the same child base power; the reduction is confluent at the level of the powers.

An unselected propagator whose two vertices fall into one component after a bundle event must also be reduced at equal times. For a massless full propagator the shared slot then contributes only its symmetric state, because the two Wightman combinations coincide at coincident times, so the odd shared state of the propagator gives zero; for a massive full propagator the two states carrying a single derivative become identified, $10 \sim 01$, which is the equal-time canonical quotient. Both cases are implemented the same way: the matrices are first built in the raw binary space of all massive and shared slots and then retracted to the true sector masters by the constant maps of (3.24), $P_s\, M^{\mathrm{raw}}\, S_s$, so the equal-time quotients require no further computation and no dense inversion. On the boundary side the same equal-time configuration makes the two branch arrangements of a coincident massive full propagator enter with the same power, and their endpoint coefficient is replaced by the mean of the two arrangements, one mean per coincident propagator, as stated in Sec. 3.5.

Relation to the single-contact case. For $|\mathcal{B}| = 1$ the two-term form (3.63) reduces to the ordinary SK pair of one propagator, (A.4) to the single difference d_e, and $c_{\{e\}} = 1$, so the bundle machinery reproduces exactly the single-propagator contacts of Sec. 3.4 and of [57], where every contact is attached to one full propagator. The new ingredients are the weight $2^{1-|S|}$, the parity selection, and the identification of the merge count with $|C| - 1$ rather than with the number of contracted propagators.

References

[1] S. Weinberg, *Quantum contributions to cosmological correlations*, *Phys. Rev. D* **72** (2005) 043514 [hep-th/0506236].

[2] X. Chen, Y. Wang and Z.-Z. Xianyu, *Schwinger-Keldysh Diagrammatics for Primordial Perturbations*, *JCAP* **12** (2017) 006 [1703.10166].

[3] N. Arkani-Hamed, D. Baumann, H. Lee and G.L. Pimentel, *The Cosmological Bootstrap: Inflationary Correlators from Symmetries and Singularities*, *JHEP* **04** (2020) 105 [1811.00024].

[4] D. Baumann, C. Duaso Pueyo, A. Joyce, H. Lee and G.L. Pimentel, *The cosmological bootstrap: weight-shifting operators and scalar seeds*, *JHEP* **12** (2020) 204 [1910.14051].

[5] D. Baumann, C. Duaso Pueyo, A. Joyce, H. Lee and G.L. Pimentel, *The Cosmological Bootstrap: Spinning Correlators from Symmetries and Factorization*, *SciPost Phys.* **11** (2021) 071 [2005.04234].

[6] E. Pajer, D. Stefanyszyn and J. Supeł, *The Boostless Bootstrap: Amplitudes without Lorentz boosts*, *JHEP* **12** (2020) 198 [2007.00027].

[7] E. Pajer, *Building a Boostless Bootstrap for the Bispectrum*, *JCAP* **01** (2021) 023 [2010.12818].

[8] G. Cabass, E. Pajer, D. Stefanyszyn and J. Supeł, *Bootstrapping large graviton non-Gaussianities*, *JHEP* **05** (2022) 077 [2109.10189].

[9] D. Baumann, D. Green, A. Joyce, E. Pajer, G.L. Pimentel, C. Sleight et al., *Snowmass White Paper: The Cosmological Bootstrap*, *SciPost Phys. Comm. Rep.* **2024** (2024) 1 [2203.08121].

[10] G.L. Pimentel and D.-G. Wang, *Boostless cosmological collider bootstrap*, *JHEP* **10** (2022) 177 [2205.00013].

[11] S. Jazayeri and S. Renaux-Petel, *Cosmological bootstrap in slow motion*, *JHEP* **12** (2022) 137 [2205.10340].

[12] Z. Qin and Z.-Z. Xianyu, *Helical inflation correlators: partial Mellin-Barnes and bootstrap equations*, *JHEP* **04** (2023) 059 [2208.13790].

[13] D.-G. Wang, G.L. Pimentel and A. Achúcarro, *Bootstrapping multi-field inflation: non-Gaussianities from light scalars revisited*, *JCAP* **05** (2023) 043 [2212.14035].

[14] Z. Qin and Z.-Z. Xianyu, *Closed-form formulae for inflation correlators*, *JHEP* **07** (2023) 001 [2301.07047].

[15] S. Aoki, L. Pinol, F. Sano, M. Yamaguchi and Y. Zhu, *Cosmological correlators with double massive exchanges: bootstrap equation and phenomenology*, *JHEP* **09** (2024) 176 [2404.09547].

[16] H. Liu and Z.-Z. Xianyu, *Massive inflationary amplitudes: differential equations and complete solutions for general trees*, *JHEP* **09** (2025) 183 [2412.07843].

[17] Z. Qin, S. Renaux-Petel, X. Tong, D. Werth and Y. Zhu, *The exact and approximate tales of boost-breaking cosmological correlators*, *JCAP* **09** (2025) 058 [2506.01555].

[18] Z.-Z. Xianyu and J. Zang, *Massive inflationary amplitudes: new representations and degenerate limits*, *JHEP* **03** (2026) 122 [2511.08677].

[19] C. Sleight, *A Mellin Space Approach to Cosmological Correlators*, *JHEP* **01** (2020) 090 [1906.12302].

[20] C. Sleight and M. Taronna, *Bootstrapping Inflationary Correlators in Mellin Space*, *JHEP* **02** (2020) 098 [1907.01143].

[21] C. Sleight and M. Taronna, *From AdS to dS exchanges: Spectral representation, Mellin amplitudes, and crossing*, *Phys. Rev. D* **104** (2021) L081902 [2007.09993].

[22] C. Sleight and M. Taronna, *From dS to AdS and back*, *JHEP* **12** (2021) 074 [2109.02725].

[23] A. Premkumar, *Regulating loops in de Sitter spacetime*, *Phys. Rev. D* **109** (2024) 045003 [2110.12504].

[24] Z. Qin and Z.-Z. Xianyu, *Phase information in cosmological collider signals*, *JHEP* **10** (2022) 192 [2205.01692].

[25] Z.-Z. Xianyu and J. Zang, *Inflation correlators with multiple massive exchanges*, *JHEP* **03** (2024) 070 [2309.10849].

[26] Z. Qin, *Cosmological correlators at the loop level*, *JHEP* **03** (2025) 051 [2411.13636].

[27] D. Marolf and I.A. Morrison, *The IR stability of de Sitter: Loop corrections to scalar propagators*, *Phys. Rev. D* **82** (2010) 105032 [1006.0035].

[28] Z.-Z. Xianyu and H. Zhang, *Bootstrapping one-loop inflation correlators with the spectral decomposition*, *JHEP* **04** (2023) 103 [2211.03810].

[29] M. Loparco, J. Penedones, K. Salehi Vaziri and Z. Sun, *The Källén-Lehmann representation in de Sitter spacetime*, *JHEP* **12** (2023) 159 [2306.00090].

[30] D. Werth, *Spectral representation of cosmological correlators*, *JHEP* **12** (2024) 017 [2409.02072].

[31] H. Zhang, *Dimensional regularization of bubble diagrams in de Sitter spacetime*, *JHEP* **02** (2026) 119 [2507.19318].

[32] B.L. Altshuler, *Kallen-Lehmann representation for Spinor-Scalar Loops in de Sitter Space-time. Spectral equations*, *JHEP* **12** (2025) 102 [2508.07467].

[33] H. Liu, Z. Qin and Z.-Z. Xianyu, *Dispersive bootstrap of massive inflation correlators*, *JHEP* **02** (2025) 101 [2407.12299].

[34] S. Das, D. Karan, B. Khatun and N. Kundu, *Reconstructing cosmological correlators via dispersion: from cutting to dressing rules*, 2602.05546.

[35] L.-T. Wang, Z.-Z. Xianyu and Y.-M. Zhong, *Precision calculation of inflation correlators at one loop*, *JHEP* **02** (2022) 085 [2109.14635].

[36] D. Werth, L. Pinol and S. Renaux-Petel, *Cosmological Flow of Primordial Correlators*, *Phys. Rev. Lett.* **133** (2024) 141002 [2302.00655].

[37] L. Pinol, S. Renaux-Petel and D. Werth, *The cosmological flow: a systematic approach to primordial correlators*, *JCAP* **02** (2025) 019 [2312.06559].

[38] D. Werth, L. Pinol and S. Renaux-Petel, *CosmoFlow: Python Package for Cosmological Correlators*, *Class. Quant. Grav.* **41** (2024) 175015 [2402.03693].

[39] J. Chen, B. Feng and Y.-X. Tao, *Multivariate hypergeometric solutions of cosmological (dS) correlators by d log-form differential equations*, *JHEP* **03** (2025) 075 [2411.03088].

[40] K.G. Chetyrkin and F.V. Tkachov, *Integration by parts: The algorithm to calculate β-functions in 4 loops*, *Nucl. Phys. B* **192** (1981) 159.

[41] A.V. Kotikov, *Differential equations method: New technique for massive Feynman diagrams calculation*, *Phys. Lett. B* **254** (1991) 158.

[42] A.V. Kotikov, *Differential equation method: The Calculation of N point Feynman diagrams*, *Phys. Lett. B* **267** (1991) 123.

[43] Z. Bern, L.J. Dixon and D.A. Kosower, *Dimensionally regulated pentagon integrals*, *Nucl. Phys. B* **412** (1994) 751 [hep-ph/9306240].

[44] T. Gehrmann and E. Remiddi, *Differential equations for two-loop four-point functions*, *Nucl. Phys. B* **580** (2000) 485 [hep-ph/9912329].

[45] J.M. Henn, *Multiloop integrals in dimensional regularization made simple*, *Phys. Rev. Lett.* **110** (2013) 251601 [1304.1806].

[46] A.V. Smirnov and F.S. Chukharev, *FIRE6: Feynman Integral REduction with modular arithmetic*, *Comput. Phys. Commun.* **247** (2020) 106877 [1901.07808].

[47] A.V. Smirnov and M. Zeng, *FIRE 7: Automatic Reduction with Modular Approach*, *Comput. Phys. Commun.* **325** (2026) 110200 [2510.07150].

[48] P. Maierhöfer, J. Usovitsch and P. Uwer, *Kira—A Feynman integral reduction program*, *Comput. Phys. Commun.* **230** (2018) 99 [1705.05610].

[49] J. Klappert, F. Lange, P. Maierhöfer and J. Usovitsch, *Integral reduction with Kira 2.0 and finite field methods*, *Comput. Phys. Commun.* **266** (2021) 108024 [2008.06494].

[50] F. Lange, J. Usovitsch and Z. Wu, *Kira 3: integral reduction with efficient seeding and optimized equation selection*, *Comput. Phys. Commun.* **322** (2026) 109999 [2505.20197].

[51] M. Hidding, *DiffExp, a Mathematica package for computing Feynman integrals in terms of one-dimensional series expansions*, *Comput. Phys. Commun.* **269** (2021) 108125 [2006.05510].

[52] X. Liu, Y.-Q. Ma and C.-Y. Wang, *A Systematic and Efficient Method to Compute Multi-loop Master Integrals*, *Phys. Lett. B* **779** (2018) 353 [1711.09572].

[53] X. Liu, Y.-Q. Ma, W. Tao and P. Zhang, *Calculation of Feynman loop integration and phase-space integration via auxiliary mass flow*, *Chin. Phys. C* **45** (2021) 013115 [2009.07987].

[54] X. Liu and Y.-Q. Ma, *AMFlow: A Mathematica package for Feynman integrals computation via auxiliary mass flow*, *Comput. Phys. Commun.* **283** (2023) 108565 [2201.11669].

[55] R.-J. Huang, X. Liu and Y.-Q. Ma, *AMFlow 2.0: significant algorithmic and software improvements for Feynman integral evaluation*, 2607.08477.

[56] Z.-F. Liu and Y.-Q. Ma, *Determining Feynman Integrals with Only Input from Linear Algebra*, *Phys. Rev. Lett.* **129** (2022) 222001 [2201.11637].

[57] J. Chen and B. Feng, *Towards systematic evaluation of de Sitter correlators via Generalized Integration-By-Parts relations*, *JHEP* **06** (2024) 199 [2401.00129].

[58] J. Chen, B. Feng, Z. Qin and Y.-X. Tao, *Loop integrals in de Sitter spacetime: The parity-split IBP system and* d log*-form differential equations*, 2604.14549.

[59] "dSIBP: a package for generating IBP seeds for QFT in de Sitter spacetime."