



Background
ROC
Placement
values
MW and AUC

Methodology
Parametric
Semiparametric
Beta

Examples
Binormal
CPAO
Texas Obesity

Future Work

References

Modeling the ROC as a Function of Covariates

Sarah Stanley

September 30, 2016



Motivation and Objective

- The ROC curve is a well-accepted measure of accuracy for diagnostic tests.
- In many applications, a test's performance is affected by covariates.
- Ignoring covariate effects can lead to faulty conclusions.
- Our goal is to investigate the effects of covariates on a test's ability to distinguish between a normal and an affected population.
- We present two existing methods (parametric and semiparametric) and introduce a new approach.

Background
ROC
Placement
values
MW and AUC

Methodology
Parametric
Semiparametric
Beta

Examples
Binormal
CPAO
Texas Obesity

Future Work

References



Outline

- Background
 - ROC and AUC
 - Placement Values
 - MW and AUC
- ROC regression methodology
 - Parametric Method
 - Semiparametric Method
 - Beta Method
- Examples
 - Binormal
 - CPAO
 - Texas Obesity
- Future Work

Background

ROC

Placement
values

MW and AUC

Methodology

Parametric

Semiparametric
Beta

Examples

Binormal

CPAO

Texas Obesity

Future Work

References



ROC and AUC

- Suppose
 Y_D = response of a subject from the diseased group
 $Y_{\bar{D}}$ = response of a subject from the non-diseased group.

- In terms of the survival function, we have

$$ROC(t) = S_D\left(S_{\bar{D}}^{-1}(t)\right), \quad t \in (0, 1)$$



- The AUC, a summary measure of the ROC, given by

$$P(Y_D > Y_{\bar{D}})$$

is the probability that a randomly selected subject is classified into the correct group.



Illustrating the AUC



- Low separation
- $ROC(t) = S_D\left(S_{\bar{D}}^{-1}(t)\right)$
 - Survival curves are nearly identical
 - ROC is close to the diagonal
- $AUC = P(Y_D > Y_{\bar{D}})$
 - Close to 0.5

Background
ROC
Placement
values
MW and AUC

Methodology
Parametric
Semiparametric
Beta



Examples
Binormal
CPAO
Texas Obesity

Future Work

References



Illustrating the AUC



- High separation
- $ROC(t) = S_D\left(S_{\bar{D}}^{-1}(t)\right)$
 - Survival curves are different
 - ROC rises more steeply
- $AUC = P(Y_D > Y_{\bar{D}})$
 - Close to 1

Background
ROC
Placement
values
MW and AUC

Methodology
Parametric
Semiparametric
Beta

Examples
Binomial
CPAO
Texas Obesity


Future Work

References



Placement Values

- We define $PV_D = S_{\bar{D}}(Y_D)$.



- The ROC is equivalent to the cdf of PV_D .

$$\begin{aligned} P[PV_D \leq t | \mathbf{X}] &= P[S_{\bar{D}\mathbf{X}}(Y_D) \leq t | \mathbf{X}] \\ &= P[Y_D \geq [S_{\bar{D}\mathbf{X}}^{-1}(t) | \mathbf{X}]] \\ &= ROC_{\mathbf{X}}(t) \end{aligned}$$

- Note also that the ROC curve can be thought of as the conditional expectation of $B_{Dt} = I[PV_D \leq t]$



Relationship between the Mann Whitney Statistic and the AUC

- The Mann-Whitney (MW) U-statistic for two independent random samples, \mathbf{x} and \mathbf{y} is given by

$$U = \sum_{i=1}^n \sum_{j=1}^m I(x_i > y_j)$$

- The MW statistic can be used as a nonparametric unbiased estimate of the AUC [Bamber(1975)].

Background

ROC

Placement
values

MW and AUC

Methodology

Parametric

Semiparametric
Beta

Examples

Binormal

CPAO

Texas Obesity

Future Work

References



Background

ROC

Placement
values

MW and AUC

Methodology

Parametric

Semiparametric

Beta

Examples

Binormal

CPAO

Texas Obesity

Future Work

References

Methodology



Direct ROC Regression Methodology

- Pepe (2002) proposed a generalized linear model (GLM) framework to directly model the ROC with covariates as follows

$$ROC_{\mathbf{X}}(t) = g^{-1}(h_0(t) + \mathbf{X}'\beta), \quad t \in (0, 1)$$

where g is a monotone link function, \mathbf{X} is a vector of covariates, h_0 is an unknown monotonic increasing function. and β is a vector of the model parameters.

- Note that the dependent variable is not directly observable, we thus estimate $ROC_{\mathbf{X}}(t)$ with either the cdf of the placement values or the conditional expectation of B_{Dt} .



Parametric ROC-GLM

- Alonzo and Pepe (2002) proposed a parametric form for $h_0(\cdot)$ such that

$$h_0(t) = \sum_{k=1}^K \alpha_k h_k(t),$$

where $\alpha = (\alpha_1, \dots, \alpha_K)$ is a vector of unknown parameters and $h(\cdot) = (h_1(\cdot), \dots, h_K(\cdot))$ are known functions.

- Thus, a parametric ROC-GLM model is

$$ROC_{\mathbf{X}}(t) = g^{-1} \left(\sum_{k=1}^K \alpha_k h_k(t) + \mathbf{X}'\beta \right), \quad t \in (0, 1).$$

Background

ROC

Placement
values

MW and AUC

Methodology

Parametric

Semiparametric
Beta

Examples

Binormal

CPAO

Texas Obesity


Future Work

References



Algorithm

- 1 Specify a set $T = \{t_\ell : \ell = 1, \dots, n_T\} \in (0, 1)$ of FPRs;
- 2 Estimate the covariate specific survival function $S_{\bar{D}X}$ for the reference population at each $t \in T$ using quantile regression.





Algorithm

Background
ROC
Placement
values
MW and AUC

Methodology
Parametric
Semiparametric
Beta

Examples
Binormal
CPAO
Texas Obesity

Future Work
References

- 1 Specify a set $T = \{t_\ell : \ell = 1, \dots, n_T\} \in (0, 1)$ of FPRs;
- 2 Estimate the covariate specific survival function $S_{\bar{D}\mathbf{X}}$ for the reference population at each $t \in T$ using quantile regression.
- 3 For each diseased observation y_{Dj} , calculate the placement values $PV_j = \hat{S}_{\bar{D}\mathbf{X}_{Dj}}(y_{Dj})$
- 4 Calculate the binary placement value indicator $\hat{B}_{jt} = I[PV_j \leq t], t \in T, j = 1, \dots, n_D;$
- 5 Fit the model $E[\hat{B}_{jt}] = g^{-1}\left(\sum_{k=1}^K \alpha_k h_k(t) + \mathbf{X}'\beta\right)$



Semiparametric ROC-GLM

- Developed by Cai(2004)
- Based on the idea that the ROC-GLM model

$$ROC_{\mathbf{X}}(t) = g^{-1}(h_0(t) + \mathbf{X}'\beta), \text{ for } t \in (0, 1)$$

is equivalent to

$$h_0(PV_D) = -\mathbf{X}'\beta + \epsilon,$$

where ϵ has known distribution g and $h_0(\cdot)$ is an unspecified increasing function.

- Essentially, pairwise comparisons of the diseased placement values are used to estimate β , and the estimates for β are then used as an offset in the estimation of $h_0(\cdot)$.



Algorithm

Background
ROC
Placement
values
MW and AUC

Methodology
Parametric
Semiparametric
Beta

Examples
Binormal
CPAO
Texas Obesity

Future Work

References

- 1 Specify a set $T = \{t_\ell : \ell = 1, \dots, n_T\} \in (0, 1)$ of FPRs;
- 2 Estimate the covariate specific survival function $S_{\bar{D}\mathbf{X}}$ via quantile regression.

- 3 Calculate the placement values $PV_j = \hat{S}_{\bar{D}\mathbf{X}_{Dj}}(y_{Dj})$

- 4 Calculate the binary placement value indicator

$$\hat{B}_{jt} = I[PV_j \leq t], t \in T, j = 1, \dots, n_D;$$

- 5 For each pair of observations in Y_D , calculate

$$\widehat{PV}_{j\ell} = I[PV_j \leq PV_\ell], \text{ and } x_{j\ell} = x_{Dj} - x_{D\ell}$$

with $j, \ell = 1, \dots, n_D, j \neq \ell$;

- 6 Fit the following GLM without an intercept to estimate β .

$$g(\widehat{PV}) = -\mathbf{X}'\beta.$$

- 7 Estimate $h_0(\cdot)$ using $\hat{\beta}$ and \hat{B}_{jt} as follows

$$g(\hat{B}_{jt}) = \text{intercept} + \text{offset}(\mathbf{X}'\hat{\beta}).$$



Consequences of parametric and semiparametric procedures

- Correlation is introduced when making pairwise comparisons.
- The resulting standard errors are thus incorrect.
- Recall, however, that the cdf of the placement values from the diseased population is equivalent to the ROC.
- A method that models the placement values directly avoids the above correlation problems.
- We implement a direct model of the placement values through Beta regression.

Background
ROC
Placement
values
MW and AUC

Methodology
Parametric
Semiparametric
Beta

Examples
Binormal
CPAO
Texas Obesity

Future Work

References



Beta Regression Model

We now introduce a Beta regression model (Ferrari, 2004). Recall that the mean and variance of $Y \sim \text{Beta}(a, b)$ are

$$E(Y) = \frac{a}{a+b} \text{ and } \text{Var}(Y) = \frac{ab}{(a+b)^2(a+b+1)}.$$

We will define the Beta regression model in terms of $\mu = E(Y)$ and a precision parameter $\phi = a + b$ so that the reparameterized beta distribution mean and variance are

$$E(Y) = \mu \text{ and } \text{Var}(Y) = \frac{\mu(1-\mu)}{1+\phi}.$$

Background
ROC
Placement
values
MW and AUC

Methodology
Parametric
Semiparametric
Beta

Examples
Binomial
CPAO
Texas Obesity

Future Work

References



Beta Regression Model

- Let y_1, \dots, y_n be independent random variables from a beta density with mean μ_t , $t = 1, \dots, n$ and precision ϕ .
- Then the beta regression model can be written as

$$g(\mu_t) = \sum_{i=1}^k x_{ti}\beta_i = \eta_t,$$

where β is a vector of regression parameters, x_{t1}, \dots, x_{tk} are observations on k covariates, and g is a monotonic link function.

- Using the logit link, we have $\mu_t = \frac{1}{1+e^{-x_t'\beta}}$. We can thus obtain the original parameters p and q from the beta distribution by calculating

$$\hat{a} = \frac{\hat{\phi}}{1 + e^{-x_t'\beta}} \text{ and } \hat{b} = \hat{\phi}\left(1 - \frac{1}{1 + e^{-x_t'\beta}}\right).$$



Beta Algorithm

Background
ROC
Placement
values
MW and AUC

Methodology
Parametric
Semiparametric
Beta

Examples
Binormal
CPAO
Texas Obesity

Future Work

References

- 1 Specify a set $T = \{t_\ell : \ell = 1, \dots, n_T\} \in (0, 1)$ of FPRs;
- 2 Estimate the covariate specific survival function $S_{\bar{D}\mathbf{X}}$ via quantile regression.
- 3 Calculate the placement values $PV_j = \hat{S}_{\bar{D}\mathbf{X}_{Dj}}(y_{Dj})$.
- 4 Perform a Beta regression on the placement values to obtain estimates of β and ϕ .
- 5 Transform to obtain $a = \mu\phi$ and $b = (1 - \mu)\phi$.
- 6 Calculate the cdf of the placement values using the Beta(a,b) distribution found above to obtain the ROC and the AUC.



Background
ROC
Placement
values
MW and AUC

Methodology
Parametric
Semiparametric
Beta

Examples

Binormal
CPAO
Texas Obesity

Future Work

References

Examples



Binormal ROC

Let

$$Y_D \sim N(\mu_D, \sigma_D^2), Y_{\bar{D}} \sim N(\mu_{\bar{D}}, \sigma_{\bar{D}}^2).$$

Then

$$ROC(t) = \Phi(a + b\Phi^{-1}(t)),$$

and

$$AUC = \Phi\left(\frac{a}{\sqrt{1+b^2}}\right),$$

where

$$a = \frac{\mu_D - \mu_{\bar{D}}}{\sigma_D}, b = \frac{\sigma_{\bar{D}}}{\sigma_D}.$$

Background
ROC
Placement
values
MW and AUC

Methodology
Parametric
Semiparametric
Beta

Examples
Binormal
CPAO
Texas Obesity

Future Work

References



Binormal Example

- Data simulated from

$$Y_D = 2 + 4X + \epsilon_D \text{ and } Y_{\bar{D}} = 1.5 + 3X + \epsilon_{\bar{D}},$$


where $X \sim U(0, 1)$ and $\epsilon_D, \epsilon_{\bar{D}} \sim N(0, 1.5^2)$.

- That is,
 $Y_D \sim N(2 + 4X, 1.5^2)$ and $Y_{\bar{D}} \sim N(1.5 + 3X, 1.5^2)$.
- Thus, the true AUC at covariate value $X = x_0$ is

$$AUC(x_0) = \Phi\left(\frac{\mu_D - \mu_{\bar{D}}}{(\sigma_D^2 + \sigma_{\bar{D}}^2)^{1/2}}\right) = \Phi\left(\frac{0.5 + x_0}{\sqrt{4.5}}\right).$$



Binormal Results



Values of x_0	0.5	0.6	0.7	0.8
Truth	0.68	0.70	0.71	0.73
Parametric	0.66	0.72	0.77	0.81
Semiparametric	0.66	0.70	0.74	0.77
Beta	0.65	0.71	0.76	0.80

Table : Comparison of AUC estimates for specified covariate values



CPAO Example

- Childhood Predictors of Adult Obesity Study (CPAO)
- Goal: determine how well childhood obesity can predict the likelihood of adult obesity.
- Questions of interest:
 - How well does childhood BMI discriminate between those who become obese in adulthood and those who do not?
 - Is the discrimination affected by gender?
- Covariates: age, gender, severity of adult obesity

Background

ROC

Placement
values

MW and AUC

Methodology

Parametric

Semiparametric
Beta

Examples

Binormal

CPAO

Texas Obesity


Future Work

References



CPAO Results

ROC for Specified Covariate Values for Females





CPAO Results

	5 yr	10 yr	15 yr	20 yr
Parametric	0.70	0.75	0.80	0.83
Semiparametric	0.86	0.88	0.90	0.91
Beta	0.74	0.78	0.81	0.84

Table : *Comparison of AUC estimates for specified ages (males)*

	5 yr	10 yr	15 yr	20 yr
Parametric	0.73	0.78	0.82	0.86
Semiparametric	0.81	0.83	0.85	0.87
Beta	0.78	0.81	0.84	0.86

Table : *Comparison of AUC estimates for specified ages (females)*

Background

ROC

Placement
values

MW and AUC

Methodology

Parametric

Semiparametric

Beta

Examples

Binormal

CPAO

Texas Obesity

Future Work

References



Texas Childhood BMI Example

Data Overview

- Five Texas counties: Cameron, Dimmit, Hidalgo, Bastrop, and McLennan
- Age, gender, height, and weight of preschool children recorded 2003 - 2008
- 2000 CDC tables will serve as the reference distribution
- Goal is to determine the effect of county as a covariate
- Placement values quantify the probability that a child's BMI exceeds a certain percentile
- We are interested in modeling these placement values with county as a covariate

Background

ROC

Placement
values

MW and AUC

Methodology

Parametric

Semiparametric
Beta

Examples

Binormal

CPAO

Texas Obesity


Future Work

References



Beta Model and Results

Background
ROC
Placement
values
MW and AUC
Methodology
Parametric
Semiparametric
Beta
Examples
Binormal
CPAO
Texas Obesity
Future Work
References



	Bastrop	Cameron	Dimmit	Hidalgo	McLennan
AUC	0.68	0.65	0.69	0.68	0.68



Conclusion

- Beta regression on the placement values yields comparable AUC estimates to those obtained via parametric and semiparametric approaches without inducing correlation.

Future Work

- Use of Historical Controls
- Meta-Analysis
- Bayesian Methods

Background
ROC
Placement
values
MW and AUC

Methodology
Parametric
Semiparametric
Beta

Examples
Binormal
CPAO
Texas Obesity

Future Work

References



References

- Alonzo, T. and M. Pepe (2002), "Distribution-free ROC analysis using binary regression techniques," *Biostatistics*, 3, 421-432.
- Bamber, D. (1975), "The area above the ordinal dominance graph and the area below the receiver operating characteristic graph," *Journal of Mathematical Psychology*, 12, 387-415.
- Cai, T. (2004), "Semi-parametric ROC regression analysis with placement values," *Biostatistics*, 5, 45-60.
- Ferrari, S. and Cribari-Neto, F. (2004), "Beta Regression for Modelling Rates and Proportions," *Journal of Applied Statistics*, 31k 799-815.
- Pepe, M. and T. Cai (2002), "The analysis of placement values for evaluating discriminatory measures," *UW Biostatistics Working Paper Series*. Working Paper 189.
- Rodriguez-Alvarez, M.X. et. al. (2011) "Comparative Study of ROC regression techniques," *Computational Statistics and Data Analysis*, 55, 888-902.