

The Exponential Distribution

Description

Density, distribution function, quantile function and random generation for the exponential distribution with rate `rate` (i.e., mean $1/\text{rate}$).

Usage

```
dexp(x, rate = 1, log = FALSE)
pexp(q, rate = 1, lower.tail = TRUE, log.p = FALSE)
qexp(p, rate = 1, lower.tail = TRUE, log.p = FALSE)
rexp(n, rate = 1)
```

Arguments

<code>x, q</code>	vector of quantiles.
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations. If <code>length(n) > 1</code> , the length is taken to be the number required.
<code>rate</code>	vector of rates.
<code>log, log.p</code>	logical; if TRUE, probabilities <code>p</code> are given as $\log(p)$.
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$, otherwise, $P[X > x]$.

Details

If `rate` is not specified, it assumes the default value of 1.

The exponential distribution with rate λ has density

$$f(x) = \lambda e^{-\lambda x}$$

for $x \geq 0$.

Value

`dexp` gives the density, `pexp` gives the distribution function, `qexp` gives the quantile function, and `rexp` generates random deviates.

The length of the result is determined by `n` for `rexp`, and is the maximum of the lengths of the numerical arguments for the other functions.

The numerical arguments other than `n` are recycled to the length of the result. Only the first elements of the logical arguments are used.

Note

The cumulative hazard $H(t) = -\log(1 - F(t))$ is `-pexp(t, r, lower = FALSE, log = TRUE)`.

Source

`dexp`, `pexp` and `qexp` are all calculated from numerically stable versions of the definitions.

`rexp` uses

Ahrens, J. H. and Dieter, U. (1972). Computer methods for sampling from the exponential and normal distributions. *Communications of the ACM*, **15**, 873–882.

References

Becker, R. A., Chambers, J. M. and Wilks, A. R. (1988) *The New S Language*. Wadsworth & Brooks/Cole.

Johnson, N. L., Kotz, S. and Balakrishnan, N. (1995) *Continuous Univariate Distributions*, volume 1, chapter 19. Wiley, New York.

See Also

[exp](#) for the exponential function.

[Distributions](#) for other standard distributions, including [dgamma](#) for the gamma distribution and [dweibull](#) for the Weibull distribution, both of which generalize the exponential.

Examples

[Run examples](#)

```
dexp(1) - exp(-1) #-> 0
```

```
## a fast way to generate *sorted* U[0,1] random numbers:
rsunif <- function(n) { n1 <- n+1
  cE <- cumsum(rexp(n1)); cE[seq_len(n)]/cE[n1] }
plot(rsunif(1000), ylim=0:1, pch=".")
abline(0,1/(1000+1), col=adjustcolor(1, 0.5))
```