

NCT99887766 — A Phase 3, Randomized, Double-Blind Study of Zoravatinib Plus Pembrolizumab vs. Placebo Plus Pembrolizumab in Treatment-Naïve Advanced Non-Small Cell Lung Cancer

Background: Zoravatinib is an oral small-molecule inhibitor of VEGFR-2 and c-MET under investigation in combination with the anti-PD-1 checkpoint inhibitor pembrolizumab. This poster reports the primary efficacy analysis from the ZORA-301 trial (data cutoff: 15-Jan-2026).

Table 1. Primary and Secondary Efficacy Endpoints by Treatment Arm

Endpoint	Arm A: Zoravatinib + Pembrolizumab (n=214)			Arm B: Placebo + Pembrolizumab (n=211)		
	ORR (%)	PFS (mo)	OS (mo)	ORR (%)	PFS (mo)	OS (mo)
Overall population	58.4	11.2	24.6	41.2	7.8	18.9
PD-L1 TPS $\geq$ 50%	67.1	13.5	27.3*	44.0	8.1	19.4
PD-L1 TPS 1-49%	52.9	9.6	22.8	39.5	7.2	17.6
EGFR/ALK wild-type	59.0	11.4	24.9	40.8	7.6	18.5

* OS in the PD-L1 TPS $\geq$ 50% subgroup was not yet mature at the data cutoff (61% of events); the 27.3-month estimate carries wide confidence intervals and should be interpreted with caution.

Safety: The most common Grade $\geq$ 3 treatment-related adverse events in Arm A were hypertension (14.2%), diarrhea (8.9%), and palmar-plantar erythrodysesthesia (6.5%). No new safety signals were identified.