EE24BTECH11030 - KEDARANANDA

Question:

On comparing the ratios $\frac{a_1}{a_2}$, $\frac{b_1}{b_2}$ and $\frac{c_1}{c_2}$, find out whether the following pair of linear equations are consistent, or inconsistent.

$$3x + 2y = 5; 2x - 3y = 7 \tag{1}$$

Theoritical Solution: To determine whether the given pair of linear equations is consistent or inconsistent, we compare the ratios $\frac{a_1}{a_2}$, $\frac{b_1}{b_2}$, and $\frac{c_1}{c_2}$, where:

$$a_1x + b_1y = c_1$$
 and $a_2x + b_2y = c_2$ (2)

From the equations:

$$3x + 2y = 5$$
 and $2x - 3y = 7$, (3)

we identify:

$$a_1 = 3, b_1 = 2, c_1 = 5, a_2 = 2, b_2 = -3, c_2 = 7.$$
 (4)

Now calculate the ratios:

$$\frac{a_1}{a_2} = \frac{3}{2}, \quad \frac{b_1}{b_2} = \frac{2}{-3}, \quad \frac{c_1}{c_2} = \frac{5}{7}.$$
 (5)

Since:

$$\frac{a_1}{a_2} \neq \frac{b_1}{b_2},\tag{6}$$

the given pair of equations is **consistent** and the lines represented by the equations are **intersecting**. Therefore, the system of equations has a unique solution.

Computational Solution:

SOLUTION USING LU FACTORIZATION

Given the system of linear equations:

$$3x + 2y = 5, (7)$$

$$2x - 3y = 7. (8)$$

We rewrite the equations as:

$$x_1 = x, (9)$$

$$x_2 = y, (10)$$

1

giving the system:

$$3x_1 + 2x_2 = 5, (11)$$

$$2x_1 - 3x_2 = 7. (12)$$

Step 1: Convert to Matrix Form

We write the system as:

$$A\mathbf{x} = \mathbf{b},\tag{13}$$

where:

$$A = \begin{bmatrix} 3 & 2 \\ 2 & -3 \end{bmatrix},\tag{14}$$

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix},\tag{15}$$

$$\mathbf{b} = \begin{bmatrix} 5 \\ 7 \end{bmatrix}. \tag{16}$$

Step 2: LU Factorization of Matrix A

We decompose A as:

$$A = LU, (17)$$

where L is a lower triangular matrix and U is an upper triangular matrix. Let:

$$L = \begin{bmatrix} 1 & 0 \\ \ell_{21} & 1 \end{bmatrix},\tag{18}$$

$$U = \begin{bmatrix} u_{11} & u_{12} \\ 0 & u_{22} \end{bmatrix}. \tag{19}$$

First, we compute U:

$$u_{11} = 3,$$
 (20)

$$u_{12} = 2, (21)$$

$$\ell_{21} = \frac{2}{3},\tag{22}$$

$$u_{22} = -3 - \left(\frac{2}{3} \cdot 2\right) = -\frac{13}{3}.\tag{23}$$

Thus, the LU decomposition is:

$$L = \begin{bmatrix} 1 & 0 \\ \frac{2}{3} & 1 \end{bmatrix},\tag{24}$$

$$U = \begin{bmatrix} 3 & 2\\ 0 & -\frac{13}{3} \end{bmatrix}. \tag{25}$$

Step 3: Solve $L\mathbf{y} = \mathbf{b}$ (Forward Substitution)

We solve:

$$L\mathbf{y} = \mathbf{b} \quad \text{or} \quad \begin{bmatrix} 1 & 0 \\ \frac{2}{3} & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} 5 \\ 7 \end{bmatrix}.$$
 (26)

From the first row:

$$y_1 = 5.$$
 (27)

From the second row:

$$\frac{2}{3}y_1 + y_2 = 7 \implies \frac{2}{3} \cdot 5 + y_2 = 7 \implies y_2 = \frac{11}{3}.$$
 (28)

Thus:

$$\mathbf{y} = \begin{bmatrix} 5 \\ \frac{11}{3} \end{bmatrix}. \tag{29}$$

Step 4: Solve $U\mathbf{x} = \mathbf{y}$ (Backward Substitution)

We solve:

$$U\mathbf{x} = \mathbf{y} \quad \text{or} \quad \begin{bmatrix} 3 & 2\\ 0 & -\frac{13}{3} \end{bmatrix} \begin{bmatrix} x_1\\ x_2 \end{bmatrix} = \begin{bmatrix} 5\\ \frac{11}{3} \end{bmatrix}.$$
 (30)

From the second row:

$$-\frac{13}{3}x_2 = \frac{11}{3} \implies x_2 = -\frac{11}{13}.$$
 (31)

From the first row:

$$3x_1 + 2x_2 = 5 \implies 3x_1 + 2\left(-\frac{11}{13}\right) = 5,$$
 (32)

$$3x_1 - \frac{22}{13} = 5 \implies 3x_1 = 5 + \frac{22}{13} = \frac{65}{13} + \frac{22}{13} = \frac{87}{13},$$
 (33)

$$x_1 = \frac{87}{39} = \frac{29}{13}. (34)$$

Thus:

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} \frac{29}{13} \\ -\frac{11}{13} \end{bmatrix}. \tag{35}$$

Final Solution

The solution is:

$$x = \frac{29}{13},\tag{36}$$

$$y = -\frac{11}{13}. (37)$$

As we can clearly see that there is solution for the given lines these are consistent

