

CONIC SECTIONS

EE24BTECH11030 - J.KEDARANANDA

14. If $a > 2b > 0$ then the positive value of m for which $y = mx - b\sqrt{1+m^2}$ is a common tangent to $x^2 + y^2 = b^2$ and $(x-a)^2 + y^2 = b^2$ is (2002S)
- $\frac{2b}{a^2-4b^2}$
 - $\frac{\sqrt{a^2-4b^2}}{2b}$
 - $\frac{2b}{a-2b}$
 - $\frac{b}{a-2b}$
15. The locus of the mid-point of the line segment joining the focus to a moving point on the parabola $y^2 = 4ax$ is another parabola with directrix (2002S)
- $x = -a$
 - $x = -a/2$
 - $x = a$
 - $x = a/2$
16. The equation of the common tangent to the curves $y^2 = 8x$ and $xy = -1$ is (2002S)
- $3y = 9x + 2$
 - $y = 2x + 1$
 - $2y = x + 8$
 - $y = x + 2$
17. The area of the quadrilateral formed by the tangents at the end points of the latus rectum to the ellipse $\frac{x^2}{9} + \frac{y^2}{5} = 1$, is (2003S)
- 27/4 sq.units
 - 9 sq.units
 - 27/2 sq.units
 - 27 sq.units
18. The focal chord to $y^2 = 16x$ is tangent to $(x-6)^2 + y^2 = 2$, then the possible values of the slope of this chord, are (2003S)
- 1, 1
 - 2, 2
 - 2, -1/2
 - 2, -1/2
19. For hyperbola $\frac{x^2}{\cos^2 \alpha} - \frac{y^2}{\sin^2 \alpha} = 1$ which of the following remains constant with change in ' α ' (2003S)
- abscissae of vertices
 - abscissae of foci
 - eccentricity
 - directrix
20. If tangents are drawn to ellipse $x^2 + 2y^2 = 2$, then the locus of the mid-point of the intercept made by the tangents between the coordinate axes is (2004S)
- $\frac{1}{2x^2} + \frac{1}{4y^2}$
 - $\frac{1}{4x^2} + \frac{1}{2x^2}$
 - $\frac{x^2}{2} + \frac{y^2}{4} = 1$
 - $\frac{x^2}{4} + \frac{y^2}{2} = 1$
21. The angle between the tangents drawn from the point (1,4) to the parabola $y^2 = 4x$ is (2004S)
- $\pi/6$

- b) $\pi/4$
- c) $\pi/3$
- d) $\pi/2$
22. If the line $2x + \sqrt{6}y = 2$ touches the hyperbola $x^2 - 2y^2 = 4$, then the point of contact is (2004S)
- a) $(-2, \sqrt{6})$
- b) $(-5, 2\sqrt{6})$
- c) $(\frac{1}{2}, \frac{1}{\sqrt{6}})$
- d) $(4, -\sqrt{6})$
23. The minimum area of the triangle formed by the tangent to the $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ & coordinate axes is (2005S)
- a) ab sq. units
- b) $\frac{a^2+b^2}{2}$ sq. units
- c) $\frac{(a+b)^2}{2}$ sq. units
- d) $\frac{a^2+ab+b^2}{3}$ sq. units
24. Tangent to the curve $y = x^2 + 6$ at a point $(1, 7)$ touches the circle $x^2 + y^2 + 16x + 12y + c = 0$ at a point Q. Then the coordinates of Q are (2005S)
- a) $(-6, -11)$
- b) $(-9, -13)$
- c) $(-10, -15)$
- d) $(-6, -7)$
25. The axis of the parabola is along the line $y=x$ and the distance of its vertex and focus from origin are $\sqrt{2}$ and $2\sqrt{2}$ respectively. If the vertex and focus both lie in the first quadrant, then the equation of the parabola is (2006-3M, -1)
- a) $(x+y)^2 = (x-y-2)$
- b) $(x-y)^2 = (x+y-2)$
- c) $(x-y)^2 = 4(x+y-2)$
- d) $(x-y)^2 = 8(x+y-2)$
26. A hyperbola, having the transverse axis of length $2\sin\theta$, is confocal with the ellipse $3x^2 + 4y^2 = 12$. Then its equation is (2007-3 marks)
- a) $x^2 \operatorname{cosec}^2\theta - y^2 \sec^2\theta = 1$
- b) $x^2 \sec^2\theta - y^2 \operatorname{cosec}^2\theta = 1$
- c) $x^2 \sin^2\theta - y^2 \cos^2\theta = 1$
- d) $x^2 \cos^2\theta - y^2 \sin^2\theta = 1$
27. Let a and b be non-zero real numbers. Then, the equation $(ax^2 + by^2 + c)(x^2 - 5xy + 6y^2) = 0$ represents (2008)
- a) four straight lines, when $c=0$ and a, b are of the same sign.
- b) two straight lines and a circle, when $a=b$, and c is of sign opposite to that of a
- c) two straight lines and a hyperbola, when a and b are of the same sign and c is of opposite to that of a
- d) a circle and an ellipse, when a and b are of the same sign and c is of sign opposite to that of a
28. Consider a branch of the hyperbola $x^2 - 2y^2 - 2\sqrt{2}x - 4\sqrt{2}y - 6 = 0$ with vertex at a point A. Let B be one of the end points of its latus rectum. If C is the focus of the hyperbola nearest to the point A, then the area of the triangle ABC is (2008)
- a) $1 - \sqrt{\frac{2}{3}}$
- b) $\sqrt{\frac{3}{2}} - 1$
- c) $1 + \sqrt{\frac{2}{3}}$
- d) $\sqrt{\frac{3}{2}} + 1$
29. The line passing through the extremity A of the major axis and extremity B of the minor

axis of the ellipse

$$x^2 + 9y^2 = 9$$

meets its auxillary circle at the point M. Then the area of the triangle with the vertices at A,M and the origin O is (2009)

a) $\frac{31}{10}$

b) $\frac{29}{10}$

c) $\frac{21}{10}$

d) $\frac{27}{10}$