

# Assignment 1

Jaswanth Chowdary Madala

- 1) If a line intersects two concentric circles (circles with the same centre) with centre  $\mathbf{O}$  at  $\mathbf{A}$ ,  $\mathbf{B}$ ,  $\mathbf{C}$  and  $\mathbf{D}$ , prove that  $AB = CD$ .




Fig. 1: Graph

**Solution:** Let the equations of two concentric circles be,

$$\|\mathbf{x}\|^2 = 4 \quad (0.0.1)$$

$$\|\mathbf{x}\|^2 = 9 \quad (0.0.2)$$

Let the equation of the line be,

$$\mathbf{x} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \mu \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (0.0.3)$$

The parameter  $\mu$  of the points of intersection of line (0.0.4) with the conic section (0.0.5)

$$\mathbf{x} = \mathbf{h} + \mu \mathbf{m} \quad (0.0.4)$$

$$g(\mathbf{x}) = \mathbf{x}^T \mathbf{V} \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f = 0 \quad (0.0.5)$$

is given by the equation

$$\mu^2 \mathbf{m}^T \mathbf{V} \mathbf{m} + 2\mu \mathbf{m}^T (\mathbf{V} \mathbf{h} + \mathbf{u}) + g(\mathbf{h}) = 0 \quad (0.0.6)$$

The points of intersection of circle (0.0.1) and the line (0.0.4)  $\mathbf{B}$ ,  $\mathbf{C}$  are given by,

$$\mathbf{V} = \mathbf{I}, \mathbf{u} = \mathbf{O}, f = -4 \quad (0.0.7)$$

$$\mathbf{h} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \mathbf{m} = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (0.0.8)$$

Substituting the above expressions (0.0.7), (0.0.8) in (0.0.6), we get

$$\mathbf{m}^T \mathbf{V} \mathbf{m} = 1 \quad (0.0.9)$$

$$\mathbf{m}^T (\mathbf{V} \mathbf{h} + \mathbf{u}) = 0 \quad (0.0.10)$$

$$g(\mathbf{h}) = -3 \quad (0.0.11)$$

$$\mu^2 - 3 = 0 \quad (0.0.12)$$

$$\mu = \pm \sqrt{3} \quad (0.0.13)$$

$$\mathbf{B} = \begin{pmatrix} 1 \\ \sqrt{3} \end{pmatrix}, \mathbf{C} = \begin{pmatrix} 1 \\ -\sqrt{3} \end{pmatrix} \quad (0.0.14)$$

The points of intersection of circle (0.0.2) and the line (0.0.4)  $\mathbf{A}$ ,  $\mathbf{D}$  are given by,

$$\mathbf{V} = \mathbf{I}, \mathbf{u} = \mathbf{O}, f = -9 \quad (0.0.15)$$

$$\mathbf{h} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \mathbf{m} = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (0.0.16)$$

Substituting the above expressions (0.0.15), (0.0.16) in (0.0.6), we get

$$\mathbf{m}^T \mathbf{V} \mathbf{m} = 1 \quad (0.0.17)$$

$$\mathbf{m}^T (\mathbf{V} \mathbf{h} + \mathbf{u}) = 0 \quad (0.0.18)$$

$$g(\mathbf{h}) = -8 \quad (0.0.19)$$

$$\mu^2 - 8 = 0 \quad (0.0.20)$$

$$\mu = \pm 2\sqrt{2} \quad (0.0.21)$$

$$\mathbf{A} = \begin{pmatrix} 1 \\ 2\sqrt{2} \end{pmatrix}, \mathbf{D} = \begin{pmatrix} 1 \\ -2\sqrt{2} \end{pmatrix} \quad (0.0.22)$$

$$\|\mathbf{A} - \mathbf{B}\| = \left\| \begin{pmatrix} 0 \\ 2\sqrt{2} - \sqrt{3} \end{pmatrix} \right\| \quad (0.0.23)$$

$$= 2\sqrt{2} - \sqrt{3} \quad (0.0.24)$$

$$\|\mathbf{C} - \mathbf{D}\| = \left\| \begin{pmatrix} 0 \\ 2\sqrt{2} - \sqrt{3} \end{pmatrix} \right\| \quad (0.0.25)$$

$$= 2\sqrt{2} - \sqrt{3} \quad (0.0.26)$$

Hence  $AB = CD$ . The parameters used in the construction are shown in the below table 1

Parameter	Value
$\mathbf{C}_1$	$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$
$r_1$	2
$\mathbf{C}_2$	$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$
$r_2$	3
$\mathbf{h}$	$\begin{pmatrix} 1 \\ 0 \end{pmatrix}$
$\mathbf{m}$	$\begin{pmatrix} 0 \\ 1 \end{pmatrix}$

TABLE 1