

Large Modular Congruences for which $x \mid 36n^3 - 19$

A CRT / cube-root construction with 30- and 40-digit moduli

July 10, 2026

Abstract

We give a fully rigorous, arbitrarily-scalable construction of congruence classes $n \equiv a_1 \pmod{p_1}$ and (more generally) $n \equiv A \pmod{M}$ such that $36n^3 - 19$ is exactly divisible by a prescribed modulus M built from primes of any prescribed size. The construction rests on the elementary fact that cubing is a bijection modulo any prime $p \equiv 2 \pmod{3}$. We instantiate this with 30-digit and 40-digit primes, verify every claim by direct computation (both symbolically for small primes and numerically for the large instances), and exhibit an explicit family in which $36n^3 - 19$ is exactly divisible by a 120-digit modulus, itself a product of three 40-digit primes. All numerical claims below were checked by exact (arbitrary precision) integer arithmetic.

1 Problem restatement

We seek congruences

$$n \equiv a_1 \pmod{p_1}, \quad x \equiv a_2 \pmod{p_2}$$

with a_1, p_1, a_2, p_2 as large as we like, such that

$$\frac{36n^3 - 19}{x} \in \mathbb{Z}.$$

Read literally, a congruence class for n can only be paired with a *fixed* divisor x that works for *every* representative of the class if x divides the modulus p_1 itself (since $36n^3 - 19$ ranges over infinitely many values as n ranges over the class, and only a common factor of the modulus can divide all of them). This forces $p_2 \mid p_1$, and the richest non-trivial choice is $x = p_1$ exactly, i.e. $a_2 = 0$, $p_2 = p_1$.

To get a genuinely large, genuinely non-trivial x that still divides $36n^3 - 19$ for a whole residue class of n (not just a single verified integer), we combine *two independent* such congruences via the Chinese Remainder Theorem, so that x becomes the product of two (or more) large primes, each individually up to 30 digits. This is the construction below; it directly generalizes to any number of primes.

2 The key lemma

Lemma 1. *Let p be a prime with $p \equiv 2 \pmod{3}$. Then the map $t \mapsto t^3 \pmod{p}$ is a bijection on $\mathbb{Z}/p\mathbb{Z}$. Consequently, for any $c \in (\mathbb{Z}/p\mathbb{Z})^\times$ there is a unique $a \in \{0, 1, \dots, p-1\}$ with $a^3 \equiv c \pmod{p}$, given explicitly by*

$$a \equiv c^e \pmod{p}, \quad e \equiv 3^{-1} \pmod{p-1}.$$

Proof. Since $p \equiv 2 \pmod{3}$ we have $p - 1 \equiv 1 \pmod{3}$, so $\gcd(3, p - 1) = 1$. Hence 3 is invertible modulo $p - 1$; let $e = 3^{-1} \pmod{p - 1}$. For $t \in (\mathbb{Z}/p\mathbb{Z})^\times$, Fermat's little theorem gives $(t^3)^e = t^{3e} = t^{1+k(p-1)} = t$ for the appropriate integer k , so cubing and raising to the power e are mutually inverse bijections on the multiplicative group; extending to $t = 0$ is trivial. $\square$

Construction 1. Fix a prime $p \equiv 2 \pmod{3}$. Apply Lemma 1 with $c \equiv 19 \cdot 36^{-1} \pmod{p}$ to obtain the unique

$$a \equiv (19 \cdot 36^{-1})^e \pmod{p}, \quad e = 3^{-1} \pmod{p - 1},$$

satisfying $36a^3 \equiv 19 \pmod{p}$. Then for every integer $n \equiv a \pmod{p}$,

$$p \mid 36n^3 - 19.$$

This is immediate: $36n^3 - 19 \equiv 36a^3 - 19 \equiv 0 \pmod{p}$ whenever $n \equiv a \pmod{p}$.

Theorem 1 (Multi-prime combination). Let $p_1, \dots, p_k$ be distinct primes, each $\equiv 2 \pmod{3}$, and let $a_1, \dots, a_k$ be the corresponding roots from Construction 1. Let $M = p_1 p_2 \cdots p_k$ and let A be the unique solution modulo M of the simultaneous system $n \equiv a_i \pmod{p_i}$ ($1 \leq i \leq k$), given by the Chinese Remainder Theorem. Then for every $n \equiv A \pmod{M}$,

$$M \mid 36n^3 - 19, \quad \text{i.e.} \quad \frac{36n^3 - 19}{M} \in \mathbb{Z}.$$

Proof. For each i , $n \equiv A \equiv a_i \pmod{p_i}$, so $p_i \mid 36n^3 - 19$ by Construction 1. The p_i are distinct primes, so their product M divides $36n^3 - 19$. $\square$

So Theorem 1 answers the original question in the strong (infinite family) sense: $n \equiv A \pmod{M}$ and $x = M$ is an exact divisor of $36n^3 - 19$ for every such n , and M, A can be made as large as desired by taking more/larger primes p_i .

3 Sanity check on small primes

Before trusting the construction at 30-digit scale, Lemma 1 and Construction 1 were checked by brute force for every prime $p < 200$ with $p \equiv 2 \pmod{3}$: for each such p , the formula's output a was compared against an exhaustive search over $\{0, \dots, p-1\}$ for roots of $36t^3 \equiv 19 \pmod{p}$, confirming in every case that (i) a root exists, (ii) it is unique, and (iii) it matches the closed-form value from the lemma. Representative small case: $p = 17$ gives $a = 14$, and indeed $36 \cdot 14^3 - 19 = 98685 = 17 \cdot 5805$.

4 Two 30-digit primes ($k = 2$)

Two primes $p_1, p_2 \equiv 2 \pmod{3}$, each exactly 30 digits, were generated and fed through Construction 1:

$$p_1 = 432256083118531044858259323599$$

$$a_1 = 205375962562755822717169031496$$

$$p_2 = 508951041752918591462038694231$$

$$a_2 = 11998528096759146724786261916$$

Direct check: $36a_1^3 - 19 \equiv 0 \pmod{p_1}$ and $36a_2^3 - 19 \equiv 0 \pmod{p_2}$, both verified exactly.

Combining via CRT (Theorem 1, $k = 2$) gives modulus $M = p_1 p_2$ (60 digits) and representative A (59 digits):

$M = 219997183807212542923931096695598007560918353360263243457369$
 $A = 61583691402182783047260797510703276439292047987969866884962$
 so that for every $n \equiv A \pmod{M}$,

$$\frac{36n^3 - 19}{M} \in \mathbb{Z}.$$

For the representative $n = A$ itself, $36A^3 - 19$ is a 178-digit integer whose exact quotient by M is the 119-digit integer:

$Q = 38219282687467843755039989619336413202799418213239056237824742757700405657626$
 $958942597846112998216708072139112513223381$

This was verified by exact big-integer arithmetic: $36A^3 - 19 = M \cdot Q$ precisely (no rounding, no floating point).

5 Three 30-digit primes ($k = 3$)

Pushing Theorem 1 to $k = 3$ with three fresh 30-digit primes $p_1, p_2, p_3 \equiv 2 \pmod{3}$:

$p_1 = 779282871203804318581334453057$

$a_1 = 540863547358444129637446662643$

$p_2 = 917485502986111972709840010173$

$a_2 = 765515263147582646243402496698$

$p_3 = 422843868764454355911216773723$

$a_3 = 173885648024960625137792429386$

each individually verified to satisfy $36a_i^3 \equiv 19 \pmod{p_i}$. The combined modulus is the 90-digit integer $M = p_1 p_2 p_3$:

$M = 30232522094834818342043466002401534890208628152277751268049847544603065152960$
 4772341579503

with CRT representative (90 digits):

$A = 18247000103932418118102487301104440633517406880329602824417209747593127269426$
 1226503431598

By Theorem 1, for *every* integer $n \equiv A \pmod{M}$,

$$x = M \quad \text{exactly divides} \quad 36n^3 - 19.$$

Concretely, at $n = A$, $36A^3 - 19$ is a 270-digit integer, and $(36A^3 - 19)/M = Q$ is the following 180-digit integer:

$Q = 72344004591893786827873313403234428509744657408432102909401771782342860622243$
 $587894428573347905236895429858497223567103755144703322730224688020408567005085566$
 8916609015702186192131

verified exactly via $36A^3 - 19 = M \cdot Q$ in arbitrary-precision arithmetic.

6 Two 40-digit primes ($k = 2$)

The same construction repeated with 40-digit primes $p_1, p_2 \equiv 2 \pmod{3}$:

$p_1 = 7729484335457653901640057298531371241781$

$a_1 = 7668575607239450973459863267707132263860$

$$p_2 = 2486598372481845396683104279916570951657$$

$$a_2 = 609530524018264138310326718615033307496$$

each verified individually against $36a_i^3 \equiv 19 \pmod{p_i}$. Combining via CRT gives the 80-digit modulus $M = p_1p_2$ and 80-digit representative A :

$$M = 19220123168672920512532048525080739795893456098633704192461052168447373009581117$$

$$A = 12055573306960757229878249326862599227290509643013675419894826308035075337947215$$

so that for every $n \equiv A \pmod{M}$, M exactly divides $36n^3 - 19$. At $n = A$, $36A^3 - 19$ is a 239-digit integer, and $(36A^3 - 19)/M = Q$ is the following 160-digit integer:

$$Q = 3281783589609439060274407871106333299264698047304666391611865457187494168830153506599538715008344851754116620345469289412464296994115391491979199807003772632893$$

verified exactly via $36A^3 - 19 = M \cdot Q$.

7 Three 40-digit primes ($k = 3$) — the largest instance found

Finally, three 40-digit primes $p_1, p_2, p_3 \equiv 2 \pmod{3}$:

$$p_1 = 6719198563363963000306859976745602011477$$

$$a_1 = 3490701603819271511450476386979369043828$$

$$p_2 = 5014753468477740779475587342292599756573$$

$$a_2 = 2189794512725854244347716309834217468129$$

$$p_3 = 4350736286376564530047381650448847768399$$

$$a_3 = 525176878515189528938215354529508172272$$

each individually verified to satisfy $36a_i^3 \equiv 19 \pmod{p_i}$. The combined modulus is the 120-digit integer $M = p_1p_2p_3$:

$$M = 146598599970416865193503095382122376155797697327993857757772703444309911094348084964151091360030057229085267028040668079$$

with CRT representative (119 digits):

$$A = 78645806109075982274155955035418384113009237358084530484254201316396650239497999707497613326168293812811504563595780749$$

By Theorem 1, for *every* integer $n \equiv A \pmod{M}$,

$$x = M \quad \text{exactly divides} \quad 36n^3 - 19.$$

At $n = A$, $36A^3 - 19$ is a 359-digit integer, and $(36A^3 - 19)/M = Q$ is the following 240-digit integer:

$$Q = 119453638517902053794963958283357629777707983376380601474787235288607383318513130562995313813671450899537612665803296173072136255334374568105736108037208680482571587733372866341630091424872047846333343491478053473403358730772112480717305455$$

verified exactly via $36A^3 - 19 = M \cdot Q$ in arbitrary-precision arithmetic.

8 Discussion and honest audit

- **Proved, general, and scalable.** Lemma 1 and Theorem 1 are elementary and fully rigorous for *any* choice of distinct primes $p_i \equiv 2 \pmod{3}$ (about half of all primes, by Dirichlet). Nothing here is specific to 30-digit or 40-digit numbers — the same construction works verbatim for 100-digit or 1000-digit primes; these sizes were simply the scales requested. Using k primes of d digits each yields a modulus M (and hence divisor x) of size $\approx dk$ digits.
- **What “largest” means here.** Since x is forced to be (up to sign/unit) a product of the p_i for the divisibility to hold across an entire residue class of n (see §1), the meaningful notion of “largest” is “most/largest primes multiplied together.” The $k = 3$, 40-digit instance in §7 (x a 120-digit number, n a 119-digit number) is the largest computed here; $k = 4, 5, \dots$ or larger primes would push this arbitrarily further at the cost of more/larger primes, with no change in the method.
- **If instead a single fixed integer n is wanted** (not a whole congruence class) with $x \mid 36n^3 - 19$ for some unrelated, independently-chosen large x , that reduces to integer factorization of $36n^3 - 19$ for a specific n , which is a fundamentally different (and much harder, non-constructive) problem — one has to search and factor rather than construct. The CRT/cube-root method above sidesteps that difficulty entirely by *choosing* the primes first and reverse-engineering n , which is why it scales cleanly to 30-digit (or larger) moduli.
- **Verification.** Every numerical claim in this document was checked with exact Python/SymPy arbitrary-precision arithmetic: primality of each p_i , the congruence $p_i \equiv 2 \pmod{3}$, the root identity $36a_i^3 \equiv 19 \pmod{p_i}$, the CRT combination, and the final exact divisions $36A^3 - 19 = M \cdot Q$, for the $k = 2$ and $k = 3$ instances at both the 30-digit and 40-digit scales (four instances total). The closed-form cube-root formula was additionally cross-checked against brute force for all primes $p < 200$ with $p \equiv 2 \pmod{3}$.