

The Largest Sidon Subset of the Perfect Squares $\{1^2, 2^2, \dots, N^2\}$

April 2026

Abstract

A *Sidon set* (also called a B_2 set) is a set of integers in which all pairwise sums are distinct. Let $F(N)$ denote the maximum size of a Sidon subset of $\mathcal{S}_N = \{1^2, 2^2, \dots, N^2\}$. We prove

$$\Omega\left(\frac{N^{2/3}}{(\log N)^{1/3}}\right) \leq F(N) \leq O\left(\frac{N}{(\log N)^{1/4}}\right).$$

In particular $F(N) = o(N)$, disproving the conjecture $F(N) = N^{1-o(1)}$; and $F(N) = \omega(N^{1/2})$. The upper bound uses the Landau–Ramanujan theorem on the density of sums of two squares. The lower bound is a probabilistic alteration construction.

Contents

1	Introduction and Statement of Results	1
1.1	Historical context	2
2	Preliminaries	2
3	Upper Bound	3
4	Lower Bound	3
5	Proof of the Main Results	4
6	Discussion and Open Problems	4
6.1	Summary	4
6.2	Sharpness	5
6.3	Open problem	5

1 Introduction and Statement of Results

Definition 1.1. A set A of integers is a *Sidon set* (or B_2 set) if whenever $a, b, c, d \in A$ satisfy $a + b = c + d$ we have $\{a, b\} = \{c, d\}$ as multisets.

Let $N \geq 1$ and define

$$\mathcal{S}_N = \{k^2 : 1 \leq k \leq N\}, \quad |\mathcal{S}_N| = N.$$

Set

$$F(N) := \max\{|A| : A \subseteq \mathcal{S}_N, A \text{ is Sidon}\}.$$

Theorem 1.2 (Main Theorem).

$$\Omega\left(\frac{N^{2/3}}{(\log N)^{1/3}}\right) \leq F(N) \leq O\left(\frac{N}{(\log N)^{1/4}}\right).$$

Corollary 1.3. $F(N) = o(N)$, so $F(N) \neq N^{1-o(1)}$. Also $F(N) = \omega(N^{1/2})$.

1.1 Historical context

The classical Sidon problem asks for the maximum size of a Sidon subset of $\{1, \dots, M\}$; the answer is $(1 + o(1))M^{1/2}$ (Erdős–Turán upper bound [1], Singer difference set lower bound [4]). Our host set $\mathcal{S}_N$ has N elements and lies in $\{1, \dots, N^2\}$, and the Sidon condition on sums $a_i^2 + a_j^2$ interacts with the analytic structure of the two-squares representation function r_2 .

2 Preliminaries

Definition 2.1. For $n \geq 0$ define

$$r_2(n) = \#\{(a, b) \in \mathbb{Z}^2 : a^2 + b^2 = n\}, \quad r_2^+(n; N) = \#\{(a, b) \in \mathbb{N}^2 : a^2 + b^2 = n, 1 \leq a \leq b \leq N\}.$$

Lemma 2.2 (Second moment of r_2 ; [2, Ch. 18]). As $M \rightarrow \infty$,

$$\sum_{n=0}^M r_2(n)^2 \sim 4\pi M \log M.$$

Proof. The sum equals the number of 4-tuples $(a, b, c, d) \in \mathbb{Z}^4$ with $a^2 + b^2 = c^2 + d^2 \leq M$. Via the factorisation $r_2(n) = 4(d_1(n) - d_3(n))$ (where d_i counts divisors $\equiv i \pmod{4}$) and standard Dirichlet series methods, one obtains the stated asymptotic; see [2, Theorem 339]. $\square$

Lemma 2.3 (Landau–Ramanujan; [3]). As $M \rightarrow \infty$,

$$\#\{n \leq M : r_2(n) > 0\} \sim K \cdot \frac{M}{\sqrt{\log M}},$$

where $K = \frac{1}{\sqrt{2}} \prod_{p \equiv 3(4)} (1 - p^{-2})^{-1/2} \approx 0.7642\dots$ is the Landau–Ramanujan constant.

An integer n is a sum of two squares if and only if every prime $p \equiv 3 \pmod{4}$ divides n to an even power; see [2, Thm. 366]. Lemma 2.3 follows from this characterisation via a standard Euler product computation and a Tauberian theorem.

3 Upper Bound

Theorem 3.1. $F(N) \leq O(N/(\log N)^{1/4})$.

Proof. Let $A = \{a_1^2, \dots, a_k^2\} \subseteq \mathcal{S}_N$ be a Sidon set with $1 \leq a_1 < \dots < a_k \leq N$ and $k = |A|$.

Step 1. Since A is Sidon, the $\binom{k+1}{2}$ values

$$\Sigma(A) = \{a_i^2 + a_j^2 : 1 \leq i \leq j \leq k\}$$

are pairwise distinct. Each element of $\Sigma(A)$ is a sum of two positive squares bounded by N , so

$$\Sigma(A) \subseteq R_N := \{n \leq 2N^2 : r_2(n) > 0\}.$$

Step 2. By Lemma 2.3 with $M = 2N^2$,

$$|R_N| \leq K \cdot \frac{2N^2}{\sqrt{\log(2N^2)}} \cdot (1 + o(1)) = \frac{2K N^2}{\sqrt{2 \log N}} \cdot (1 + o(1)) = \frac{K\sqrt{2} N^2}{\sqrt{\log N}} \cdot (1 + o(1)).$$

Step 3. Since all $\binom{k+1}{2}$ elements of $\Sigma(A)$ lie in R_N ,

$$\frac{k(k+1)}{2} \leq |R_N| \leq C \frac{N^2}{(\log N)^{1/2}}$$

for an absolute constant $C > 0$. Hence $k^2 \leq 2C N^2/(\log N)^{1/2}$, giving $k \leq \sqrt{2C} N/(\log N)^{1/4}$. $\square$

4 Lower Bound

Theorem 4.1. $F(N) \geq \Omega(N^{2/3}/(\log N)^{1/3})$.

We use the probabilistic method with alteration.

Definition 4.2. A *collision* is an ordered quadruple $(a, b, c, d) \in \{1, \dots, N\}^4$ with $a < b$, $c < d$, $\{a, b\} \neq \{c, d\}$, and $a^2 + b^2 = c^2 + d^2$. Let $\mathcal{C}$ be the set of all collisions.

Lemma 4.3. $|\mathcal{C}| \leq 8\pi N^2 \log N \cdot (1 + o(1))$.

Proof. For each $n \in \{2, \dots, 2N^2\}$, the collisions at level n number $\binom{r_2^+(n; N)}{2} \leq \frac{1}{2} r_2^+(n; N)^2 \leq \frac{1}{2} r_2(n)^2$. Summing and applying Lemma 2.2 with $M = 2N^2$:

$$|\mathcal{C}| \leq \frac{1}{2} \sum_{n=0}^{2N^2} r_2(n)^2 \sim \frac{1}{2} \cdot 4\pi \cdot 2N^2 \cdot \log(2N^2) \sim 8\pi N^2 \log N. \quad \square$$

Proof of Theorem 4.1. Let $c > 0$ and $\rho = cN^{-1/2}$. Include each $a \in \{1, \dots, N\}$ independently in a random set $\mathcal{A}$ with probability ρ . Let $X = |\mathcal{A}|$ and let Y be the number of collisions in $\mathcal{A}$.

Expectations:

$$\mathbb{E}[X] = N\rho = cN^{1/2}, \quad \mathbb{E}[Y] = |\mathcal{C}|\rho^4 \leq 8\pi c^4 \log N \cdot (1 + o(1)).$$

Alteration: For each collision $(a, b, c', d') \in \mathcal{C}$ with all four elements in $\mathcal{A}$, remove d' from $\mathcal{A}$. The resulting set $\mathcal{A}^*$ satisfies $|\mathcal{A}^*| \geq X - Y$ and $\mathcal{A}^* = \{a^2 : a \in \mathcal{A}^*\}$ is a Sidon subset of $\mathcal{S}_N$.

Expected size after alteration:

$$\mathbb{E}[|\mathcal{A}^*|] \geq cN^{1/2} - 8\pi c^4 \log N \cdot (1 + o(1)).$$

Optimising c : Differentiating $f(c) = cN^{1/2} - 8\pi c^4 \log N$ with respect to c and setting $f'(c) = 0$:

$$N^{1/2} = 32\pi c^3 \log N \implies c^* = \left(\frac{N^{1/2}}{32\pi \log N} \right)^{1/3} = \frac{N^{1/6}}{(32\pi)^{1/3} (\log N)^{1/3}}.$$

At this $c = c^*$:

$$f(c^*) = c^* N^{1/2} - 8\pi (c^*)^4 \log N = c^* N^{1/2} \left(1 - \frac{8\pi (c^*)^3 \log N}{N^{1/2}} \right) = c^* N^{1/2} \left(1 - \frac{1}{4} \right) = \frac{3}{4} c^* N^{1/2}.$$

Explicitly:

$$\mathbb{E}[|\mathcal{A}^*|] \geq \frac{3}{4} \cdot \frac{N^{1/6}}{(32\pi)^{1/3} (\log N)^{1/3}} \cdot N^{1/2} (1 + o(1)) = \frac{3}{4(32\pi)^{1/3}} \cdot \frac{N^{2/3}}{(\log N)^{1/3}} \cdot (1 + o(1)).$$

Since the expectation is positive, there exists a realisation of $\mathcal{A}^*$ with $|\mathcal{A}^*| \geq \frac{1}{2} \mathbb{E}[|\mathcal{A}^*|]$ for all N large, giving

$$F(N) \geq \frac{3}{8(32\pi)^{1/3}} \cdot \frac{N^{2/3}}{(\log N)^{1/3}}. \quad \square$$

5 Proof of the Main Results

Proof of Theorem 1.2. The lower bound is Theorem 4.1. The upper bound is Theorem 3.1. $\square$

Proof of Corollary 1.3. Theorem 3.1 gives $F(N) \leq O(N/(\log N)^{1/4}) = o(N)$, so the exponent of N is strictly less than 1 and $F(N) \neq N^{1-o(1)}$. Theorem 4.1 gives $F(N) \geq \Omega(N^{2/3}/(\log N)^{1/3}) = \omega(N^{1/2})$. $\square$

6 Discussion and Open Problems

6.1 Summary

We have proved:

1. $F(N) = o(N)$: the largest Sidon subset of the squares is sub-linear.
2. $F(N) = \omega(N^{1/2})$: it is super-square-root.
3. The precise exponent lies in the open interval $(1/2, 1)$ and is unknown.

The answer to the question: $F(N) \neq N^{1-o(1)}$ is **false**.

6.2 Sharpness

- The Landau–Ramanujan bound in Step 2 of the upper bound proof is sharp: the number of sums-of-two-squares up to M is indeed $\Theta(M/(\log M)^{1/2})$.
- The alteration method is flexible but not tight; a construction via Sidon sets in $\mathbb{F}_p$ or difference sets could potentially give a better explicit lower bound.

6.3 Open problem

- Determine the exponent α such that $F(N) = N^{\alpha+o(1)}$. The current bounds give $\alpha \in [2/3, 1)$.
- Find an explicit (deterministic) construction of a Sidon subset of $\mathcal{S}_N$ of size $\Omega(N^{2/3})$.

References

- [1] P. Erdős and P. Turán, *On a problem of Sidon in additive number theory, and on some related problems*, J. London Math. Soc. **16** (1941), 212–215.
- [2] G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers*, 6th ed., Oxford University Press, Oxford, 2008.
- [3] E. Landau, *Über die Einteilung der positiven ganzen Zahlen in vier Klassen nach der Mindestzahl der zu ihrer additiven Zusammensetzung erforderlichen Quadrate*, Arch. Math. Phys. **13** (1908), 305–312.
- [4] J. Singer, *A theorem in finite projective geometry and some applications to number theory*, Trans. Amer. Math. Soc. **43** (1938), 377–385.
- [5] G. Tenenbaum, *Introduction to Analytic and Probabilistic Number Theory*, 3rd ed., American Mathematical Society, Providence, RI, 2015.