

GENERALIZATION OF THE MULTIPLE ORTHOGONALITY TO THE BIVARIATE CASE

J. ANTONIO VILLEGAS, LIDIA FERNÁNDEZ

ABSTRACT

Polynomials known as Multiple Orthogonal Polynomials (MOPs) in a single variable are polynomials that satisfy orthogonality conditions concerning multiple measures and play significant role in several applications such as Hermite-Padé approximation, random matrix theory or integrable systems. However, this theory has only been studied in the univariate case. In this poster, some generalized definitions of the two main types of multiple orthogonality are given, together with some examples and extended results.

Keywords: Orthogonal Polynomials, Approximation Theory, Applications, Multiple orthogonality.

AMS Classification: 33C45, 33C50, 42C05.

BIBLIOGRAPHY

- [1] M. E. H. Ismail and W. Van Assche, *Classical and quantum orthogonal polynomials in one variable*, Encyclopedia of mathematics and its applications, Cambridge University Press (2005).
- [2] C. F. Dunkl and Y. Xu *Orthogonal Polynomials of Several Variables*, Cambridge University Press (2014).
- [3] W. Van Assche *Orthogonal and multiple orthogonal polynomials, random matrices, and painlevé equations*, “Orthogonal Polynomials” (M. Foupouagnigni, W. Koepf, eds), Tutorials, Schools and Workshops in the Mathematical Sciences, Springer Nature Switzerland (2020) 629–683.

J. Antonio Villegas,
Instituto de Matemáticas (IMAG) and Departamento de Matemática Aplicada,
Universidad de Granada.
`jantoniovr@ugr.es`

Lidia Fernández,
Instituto de Matemáticas (IMAG) and Departamento de Matemática Aplicada,
Universidad de Granada.
`lidiafr@ugr.es`